

WHITTAKER COINVARIANTS FOR $GL(m|n)$

JONATHAN BRUNDAN AND SIMON M. GOODWIN

ABSTRACT. Let $W_{m|n}$ be the (finite) W -algebra attached to the principal nilpotent orbit in the general linear Lie superalgebra $\mathfrak{gl}_{m|n}(\mathbb{C})$. In this paper we study the *Whittaker coinvariants functor*, which is an exact functor from category \mathcal{O} for $\mathfrak{gl}_{m|n}(\mathbb{C})$ to a certain category of finite-dimensional modules over $W_{m|n}$. We show that this functor has properties similar to Soergel's functor \mathbb{V} in the setting of category \mathcal{O} for a semisimple Lie algebra. We also use it to compute the center of $W_{m|n}$ explicitly, and deduce consequences for the classification of blocks of \mathcal{O} up to Morita/derived equivalence.

1. INTRODUCTION

This article is a sequel to [BBG], in which we began a study of the *principal W -algebra* $W = W_{m|n}$ associated to the general linear Lie superalgebra $\mathfrak{g} = \mathfrak{gl}_{m|n}(\mathbb{C})$. This associative superalgebra is a quantization of the Slodowy slice to the principal nilpotent orbit in \mathfrak{g} ; see e.g. [P, GG, L1] for more about (finite) W -algebras in the purely even case.

There are several different approaches to the construction of W . We begin by briefly recalling one of these in more detail. Since $\mathfrak{gl}_{m|n}(\mathbb{C}) \cong \mathfrak{gl}_{n|m}(\mathbb{C})$, there is no loss in generality in assuming throughout the article that $m \leq n$. Pick a nilpotent element $e \in \mathfrak{g}_{\bar{0}}$ with just two Jordan blocks (necessarily of sizes m and n), and let $\mathfrak{g} = \bigoplus_{d \in \mathbb{Z}} \mathfrak{g}(d)$ be a good grading for $e \in \mathfrak{g}(1)$. Let $\mathfrak{p} := \bigoplus_{d \geq 0} \mathfrak{g}(d)$ and $\mathfrak{m} := \bigoplus_{d < 0} \mathfrak{g}(d)$. We get a generic character $\chi : \mathfrak{m} \rightarrow \mathbb{C}$ by taking the supertrace from d with e . Setting $\mathfrak{m}_{\chi} := \{x - \chi(x) \mid x \in \mathfrak{m}\} \subseteq U(\mathfrak{m})$, we then have by definition that

$$W := \{u \in U(\mathfrak{p}) \mid u \mathfrak{m}_{\chi} \subseteq \mathfrak{m}_{\chi} U(\mathfrak{g})\}.$$

In [BBG], we obtained a presentation for W by generators and relations, showing that it is a certain truncated shifted version of the Yangian $Y(\mathfrak{gl}_{1|1})$. In particular, it is quite close to being supercommutative. We also classified its irreducible representations via highest weight theory. Every irreducible representation arises as a quotient of an appropriately defined Verma module, all of which have dimension 2^m . Then there is another more explicit construction of the irreducible representations, implying that they have dimension 2^{m-t} for some *atypicality* $0 \leq t \leq m$.

By a *Whittaker vector*, we mean a vector v in some right \mathfrak{g} -module such that $vx = \chi(x)v$ for each $x \in \mathfrak{m}$; equivalently, $v \mathfrak{m}_{\chi} = 0$. This is the appropriate analog for \mathfrak{g} of the notion of a Whittaker vector for a semisimple Lie algebra as studied in Kostant's classic paper [Ko]. From the definition of W , we see

2010 *Mathematics Subject Classification*: 17B10, 17B37.

First author supported in part by NSF grant nos. DMS-1161094 and DMS-1700905.

Second author supported in part by EPSRC grant no. EP/R018952/1.

that the space of Whittaker vectors, which we denote by $H^0(M)$, is a right W -module. We refer to H^0 as the *Whittaker invariants* functor. On the other hand, for a left \mathfrak{g} -module M , it is clear from the definition of W that the space $H_0(M) := M/\mathfrak{m}_\chi M$ of *Whittaker coinvariants* is a left W -module. The restriction of this functor to the BGG category \mathcal{O} for \mathfrak{g} (defined with respect to the standard Borel subalgebra \mathfrak{b} of \mathfrak{g} such that $\mathfrak{b}_0 = \mathfrak{p}_0$) gives an exact functor from \mathcal{O} to the category of finite-dimensional left W -modules.

The main goal in the first part of this article is to describe the effect of H_0 on various natural families of modules in \mathcal{O} . In particular, in Theorem 3.14, we show that it sends Verma modules in \mathcal{O} (induced from the standard Borel \mathfrak{b}) to the Verma modules for W . Our proof of this is elementary but surprisingly technical, and it turns out to be the key ingredient needed for many things after that. We use it to show that H_0 sends irreducible modules in \mathcal{O} of maximal Gelfand–Kirillov dimension to irreducible W -modules, and it sends all other irreducibles in \mathcal{O} to zero. Moreover, every irreducible W -module arises in this way. We also compute the composition multiplicities of Verma modules for W . They always have composition length 2^t where t is the atypicality mentioned earlier, but are not necessarily multiplicity-free. As another more surprising application, we deduce that the center of W is canonically isomorphic to the center of $U(\mathfrak{g})$; see Theorem 3.21. Thus central characters for \mathfrak{g} and W are identified.

After that, we restrict attention just to the subcategory $\mathcal{O}_{\mathbb{Z}}$ of \mathcal{O} that is the sum of all of its blocks with integral central character. Let $\overline{\mathcal{O}}_{\mathbb{Z}}$ be the full subcategory of W -mod consisting of the W -modules isomorphic to $H_0(M)$ for $M \in \mathcal{O}_{\mathbb{Z}}$. We show that $\overline{\mathcal{O}}_{\mathbb{Z}}$ is Abelian, and the Whittaker coinvariants functor restricts to an exact functor

$$H_0 : \mathcal{O}_{\mathbb{Z}} \rightarrow \overline{\mathcal{O}}_{\mathbb{Z}}$$

which satisfies the universal property of the quotient of $\mathcal{O}_{\mathbb{Z}}$ by the Serre subcategory $\mathcal{T}_{\mathbb{Z}}$ consisting of all the modules of less than maximal Gelfand–Kirillov dimension; see Theorem 4.8. Thus, $\overline{\mathcal{O}}_{\mathbb{Z}}$ is an explicit realization of the Serre quotient $\mathcal{O}_{\mathbb{Z}}/\mathcal{T}_{\mathbb{Z}}$. By [BLW, Theorem 4.10], the quotient functor $\mathcal{O}_{\mathbb{Z}} \rightarrow \mathcal{O}_{\mathbb{Z}}/\mathcal{T}_{\mathbb{Z}}$ is fully faithful on projectives, hence, so too is H_0 . This is reminiscent of a result of Backelin [Ba] in the setting of category \mathcal{O} for a semisimple Lie algebra. Backelin’s result was based ultimately on the *Struktursatz* from [S]. In that case, Soergel’s *Endomorphismensatz* shows moreover that the blocks of the quotient category can be realized explicitly in terms of the cohomology algebras of some underlying partial flag varieties.

It would be very interesting to establish some sort of analog of Soergel’s *Endomorphismensatz* in the super case. Ideally, this would give an explicit combinatorial description (e.g. by quiver and relations) of the basic algebras B_ξ that are Morita equivalent to the various blocks $\overline{\mathcal{O}}_\xi$ of our category $\overline{\mathcal{O}}_{\mathbb{Z}}$. Note these algebras are not commutative in general; e.g. see [B5, Example 4.7 and Remark 4.8] for some baby examples. In Soergel’s proof of the *Endomorphismensatz*, the cohomology algebras of partial flag varieties arise as quotients of $Z(\mathfrak{g})$, which is also the principal W -algebra in that setting according to [Ko]. Paralleling this in the super case, we show that all the maximally atypical B_ξ ’s

can be realized as quotients of a certain idempotent form \dot{W} of W ; see Theorems 4.22 and 4.25. We also compute explicitly the Cartan matrix of B_ξ ; see Theorem 4.14 for an elementary proof based on properties of the Whittaker coinvariants functor, and Theorem 4.27 for a proof based on the super Kazhdan-Lusztig conjecture of [CLW, BLW] (which has the advantage of incorporating the natural grading).

We end the article by discussing some applications to the classification of blocks of $\mathcal{O}_\mathbb{Z}$, both up to Morita equivalence and up to gradable derived equivalence in the sense of [CM, Definition 4.2]; see Theorems 4.33 and 4.35 and Conjectures 4.34 and 4.37.

Finally in the introduction, we draw attention to the work of Losev in [L2]. This paper includes a study of Whittaker coinvariants functors associated to arbitrary nilpotent orbits in semisimple Lie algebras. Lie superalgebras are not considered in detail, though some remarks are made about how the theory may apply in this situation in [L2, §6.3.2]. His theory includes many of the features discussed above. In particular, he also views these functors as some generalized Soergel functors. The approach in this paper is quite different and leads to more explicit results, which we require for the subsequent applications.

Acknowledgements. This article was written in part during the programme ‘‘Local Representation Theory and Simple Groups’’ held at the Bernoulli Center, EPFL, Lausanne, Switzerland in Autumn 2016. We also thank Kevin Coulemier for pointing out the counterexample mentioned after Conjecture 4.37.

Notation. We fix once and for all some choice of *parity function* $\text{par} : \mathbb{C} \rightarrow \mathbb{Z}/2$ such that $\text{par}(0) = \bar{0}$ and $\text{par}(z + 1) = \text{par}(z) + \bar{1}$ for all $z \in \mathbb{C}$. Also, \leq denotes the partial order on \mathbb{C} defined by $z \leq w$ if $w - z \in \mathbb{N}$.

2. CATEGORY \mathcal{O} FOR THE GENERAL LINEAR LIE SUPERALGEBRA

In this section, we set up our general combinatorial notation, then review various standard facts about category \mathcal{O} for $\mathfrak{gl}_{m|n}(\mathbb{C})$. We assume from the outset that $m \leq n$ as this will be essential when we introduce the W -algebra in the next section, but note that the general results in this section do not depend on this hypothesis.

2.1. Combinatorics. We fix integers $0 \leq m \leq n$ and a two-rowed *pyramid* π with m boxes in the first (top) row and n boxes in the second (bottom) row. We require that the top row does not jut out past the bottom row. For example, here are the possible pyramids for $m = 2$ and $n = 5$:

$$\begin{array}{|c|c|} \hline 1 & 2 \\ \hline \end{array} \begin{array}{|c|c|c|c|c|} \hline 3 & 4 & 5 & 6 & 7 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline \end{array} \begin{array}{|c|c|c|c|} \hline 3 & 4 & 5 & 6 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline \end{array} \begin{array}{|c|c|c|c|} \hline 3 & 4 & 5 & 6 \\ \hline \end{array}, \quad \begin{array}{|c|c|} \hline 1 & 2 \\ \hline \end{array} \begin{array}{|c|c|c|} \hline 3 & 4 & 5 \\ \hline \end{array}.$$

As in these examples, we number the boxes of π by $1, \dots, m + n$, so that the boxes in the first (resp. second) row are indexed $1, \dots, m$ (resp. $m + 1, \dots, m + n$) from left to right. Then we write $\text{row}(i)$ and $\text{col}(i)$ for the row and column numbers of the i th box of π , numbering columns by $1, \dots, n$ in order from left to right. Also we denote the number of columns of height 1 on the left (resp. right) side of π by s_- (resp. s_+); in the degenerate case $m = 0$, one should instead pick any $s_-, s_+ \geq 0$ with $s_- + s_+ = n$.

A π -*tableau* is a filling of the boxes of the pyramid π by complex numbers. Let Tab denote the set of all such π -tableaux. Sometimes we will represent $A \in \text{Tab}$ as an array $A = \begin{smallmatrix} a_1 \cdots a_m \\ b_1 \cdots b_n \end{smallmatrix}$ of complex numbers. Here are a few combinatorial notions about tableaux.

- For $A = \begin{smallmatrix} a_1 \cdots a_m \\ b_1 \cdots b_n \end{smallmatrix}$, we let $a(A) := a_1 + \cdots + a_m$ and $b(A) := b_1 + \cdots + b_n$ be the sum of the entries on its top and bottom rows, respectively.
- We say that $A = \begin{smallmatrix} a_1 \cdots a_m \\ b_1 \cdots b_n \end{smallmatrix}$ is *dominant* if $a_1 > \cdots > a_m$ and $b_1 < \cdots < b_n$.
- We say that $A = \begin{smallmatrix} a_1 \cdots a_m \\ b_1 \cdots b_n \end{smallmatrix}$ is *anti-dominant* if $a_i \succ a_j$ for each $1 \leq i < j \leq m$ and $b_i \prec b_j$ for each $1 \leq i < j \leq n$.
- A *matched pair* in A is a pair of equal entries from the same column.
- The *defect* $\text{def}(A)$ is the number of matched pairs in A .
- Two π -tableaux A, B are *row equivalent*, denoted $A \sim B$, if B can be obtained from A by rearranging entries within each row.
- The *degree of atypicality* $\text{atyp}(A)$ is the maximal defect of any $B \sim A$.
- We write $B \cup A$ if B can be obtained from A by picking several of the matched pairs in A and subtracting 1 from each of them. There are $2^{\text{def}(A)}$ such tableaux B .
- The *Bruhat order* \leq on Tab is the smallest partial order such that $B < A$ whenever one of the following holds:
 - B is obtained from $A = \begin{smallmatrix} a_1 \cdots a_m \\ b_1 \cdots b_n \end{smallmatrix}$ by interchanging a_i and a_j , assuming $a_i > a_j$ for some $1 \leq i < j \leq m$;
 - B is obtained from $A = \begin{smallmatrix} a_1 \cdots a_m \\ b_1 \cdots b_n \end{smallmatrix}$ by interchanging b_i and b_j , assuming $b_i < b_j$ for some $1 \leq i < j \leq n$;
 - B is obtained from $A = \begin{smallmatrix} a_1 \cdots a_m \\ b_1 \cdots b_n \end{smallmatrix}$ by subtracting one from both a_i and b_j , assuming $a_i = b_j$ for some $1 \leq i \leq m$ and $1 \leq j \leq n$.
- Let \approx be the equivalence relation generated by the Bruhat order \leq . We refer to the \approx -equivalence classes as *linkage classes*. All π -tableaux in a given linkage class ξ have the same atypicality.

The representation theoretic significance of these definitions will be made clear later in the article.

2.2. Modules and supermodules. In the introduction we have ignored the distinction between modules and supermodules. We will be more careful in the remainder of the article. For an associative algebra A , we write $A\text{-mod}$ or $A\text{-mod}_{\text{fd}}$ for the categories of left A -modules or finite-dimensional left A -modules, respectively.

Superalgebras and supermodules are objects in the symmetric monoidal category of vector superspaces. We denote the parity of a homogeneous vector v in a vector superspace by $|v| \in \mathbb{Z}/2$, and recall that the tensor flip $V \otimes W \xrightarrow{\sim} W \otimes V$ is given on homogeneous vectors by $v \otimes w \mapsto (-1)^{|v||w|} w \otimes v$. The notation $[\cdot, \cdot]$ always denotes the *supercommutator* $[x, y] = xy - (-1)^{|x||y|} yx$ of homogeneous elements of a superalgebra.

Let A be an associative superalgebra. A left A -*supermodule* is a superspace $M = M_{\bar{0}} \oplus M_{\bar{1}}$ equipped with a linear left action of A such that $A_i M_j \subseteq M_{i+j}$. A *supermodule homomorphism* is a parity-preserving linear map that is a homomorphism in the usual sense. We write $A\text{-smod}$ for the Abelian category of all left A -supermodules and supermodule homomorphisms, and $A\text{-smod}_{\text{fd}}$ for

the subcategory of finite-dimensional ones. We denote the usual *parity switching functor* on all of these categories by Π .

2.3. Super category \mathcal{O} . Let \mathfrak{g} be the Lie superalgebra $\mathfrak{gl}_{m|n}(\mathbb{C})$. We write $e_{i,j}$ for the ij -matrix unit in \mathfrak{g} , which is of parity $|i| + |j|$ where

$$|i| = \begin{cases} \bar{0} & \text{for } 1 \leq i \leq m, \\ \bar{1} & \text{for } m+1 \leq i \leq m+n. \end{cases}$$

Let \mathfrak{t} be the Cartan subalgebra of \mathfrak{g} consisting of all diagonal matrices and $\{\delta_i\}_{1 \leq i \leq m+n}$ be the basis for \mathfrak{t}^* dual to the basis $\{e_{i,i}\}_{1 \leq i \leq m+n}$ of \mathfrak{t} . The usual *supertrace form* (\cdot, \cdot) on \mathfrak{g} induces a non-degenerate super-symmetric bilinear form (\cdot, \cdot) on \mathfrak{t}^* such that $(\delta_i, \delta_j) = (-1)^{|i|} \delta_{i,j}$.

Now suppose that \triangleleft is a total order on the set $\{1, \dots, m+n\}$. Let $\mathfrak{b}^{\triangleleft}$ be the Borel subalgebra of \mathfrak{g} spanned by $\{e_{i,j}\}_{i \triangleleft j}$. Then define $\mathcal{O}^{\triangleleft}$ to be the full category of $U(\mathfrak{g})$ -smod consisting of all \mathfrak{g} -supermodules M such that

- M is finitely generated over \mathfrak{g} ;
- M is locally finite-dimensional over $\mathfrak{b}^{\triangleleft}$;
- M is semisimple over \mathfrak{t} ;
- the λ -weight space M_λ of M is concentrated in parity¹

$$\text{par}(\lambda) := \text{par}((\lambda, \delta_{m+1} + \dots + \delta_{m+n})) + [(n-m)/2] + m s_- \in \mathbb{Z}/2. \quad (2.1)$$

The parity assumption means that one can simply forget the $\mathbb{Z}/2$ -grading on objects of $\mathcal{O}^{\triangleleft}$, since it can be recovered uniquely from the weights. The reader should not be concerned about the dependence on the choice of the function $\text{par} : \mathbb{C} \rightarrow \mathbb{Z}/2$ (which was made at the end of the introduction): the categories $\mathcal{O}^{\triangleleft}$ arising from two different choices are obviously equivalent. We note that all objects of $\mathcal{O}^{\triangleleft}$ are of finite length.

Introduce the weight $\rho^{\triangleleft} \in \mathfrak{t}^*$ so that

$$(\rho^{\triangleleft}, \delta_j) = \# \{i \triangleleft j \mid |i| = \bar{1}\} - \# \{i \triangleleft j \mid |i| = \bar{0}\} \quad (2.2)$$

for $j = 1, \dots, m+n$. For $\mathbf{A} \in \text{Tab}$, let $\lambda_{\mathbf{A}}^{\triangleleft} \in \mathfrak{t}^*$ be the unique weight such that $(\lambda_{\mathbf{A}}^{\triangleleft} + \rho^{\triangleleft}, \delta_j)$ is the entry in the j th box of \mathbf{A} . Then we let $M^{\triangleleft}(\mathbf{A})$ denote the *Verma supermodule* of $\mathfrak{b}^{\triangleleft}$ -highest weight $\lambda_{\mathbf{A}}^{\triangleleft}$, i.e.

$$M^{\triangleleft}(\mathbf{A}) := U(\mathfrak{g}) \otimes_{U(\mathfrak{b}^{\triangleleft})} \mathbb{C}_{\mathbf{A}}^{\triangleleft} \quad (2.3)$$

where $\mathbb{C}_{\mathbf{A}}^{\triangleleft}$ is a one-dimensional $\mathfrak{b}^{\triangleleft}$ -supermodule of weight $\lambda_{\mathbf{A}}^{\triangleleft}$ concentrated in parity $\text{par}(\lambda_{\mathbf{A}}^{\triangleleft})$. Note the parity choice here is forced upon us since we want $M^{\triangleleft}(\mathbf{A})$ to belong to $\mathcal{O}^{\triangleleft}$. The Verma supermodule $M^{\triangleleft}(\mathbf{A})$ has a unique irreducible quotient $L^{\triangleleft}(\mathbf{A})$, and the supermodules $\{L^{\triangleleft}(\mathbf{A}) \mid \mathbf{A} \in \text{Tab}\}$ give a complete set of inequivalent irreducible objects in $\mathcal{O}^{\triangleleft}$.

By a *normal order* we mean a total order \triangleleft such that $1 \triangleleft \dots \triangleleft m$ and $m+1 \triangleleft \dots \triangleleft m+n$. For any normal order \triangleleft , the underlying even subalgebra $\mathfrak{b}_0^{\triangleleft}$ is equal simply to the usual standard Borel subalgebra of $\mathfrak{g}_0 = \mathfrak{gl}_m(\mathbb{C}) \oplus \mathfrak{gl}_n(\mathbb{C})$ consisting of upper triangular matrices. Observing that a \mathfrak{g} -supermodule is locally finite over $\mathfrak{b}^{\triangleleft}$ if and only if it is locally finite over $\mathfrak{b}_0^{\triangleleft}$, it is clear for

¹We have made this particular choice so that (2.4) holds; it is important also in the proof of Lemma 3.8.

any two normal orders \triangleleft and \blacktriangleleft that $\mathcal{O}^{\triangleleft} = \mathcal{O}^{\blacktriangleleft}$. Henceforth, we denote this category coming from a normal order simply by \mathcal{O} .

Let us explain how to translate between the various labellings of the irreducible objects of \mathcal{O} arising from different choices of normal order. The basic technique to pass from \triangleleft to \blacktriangleleft is to apply a sequence of *odd reflections* connecting \triangleleft to \blacktriangleleft . A single odd reflection connects normal orders \triangleleft and \blacktriangleleft which agree except at $i \in \{1, \dots, m\}$ and $j \in \{m+1, \dots, m+n\}$, with i, j being consecutive in both orders. Assuming that $i \triangleleft j$, we have that $L^{\triangleleft}(\mathbf{A}) \cong L^{\blacktriangleleft}(\mathbf{A}')$ where \mathbf{A}' is obtained from \mathbf{A} by adding 1 to its i th and j th entries if these entries are equal, or $\mathbf{A}' := \mathbf{A}$ if these entries are different. This was observed originally by Serganova in her PhD thesis.

The following fundamental lemma is well known, e.g. see [CLW, Lemma 6.1]. The proof involves noting that it suffices to consider the case when \triangleleft and \blacktriangleleft are connected by a single odd reflection, and then it can be observed from explicit formulas for the characters.

Lemma 2.1. *For any two normal orders \triangleleft and \blacktriangleleft and $\mathbf{A} \in \text{Tab}$, the formal characters of $M^{\triangleleft}(\mathbf{A})$ and $M^{\blacktriangleleft}(\mathbf{A})$ are equal. Thus the symbols $[M^{\triangleleft}(\mathbf{A})]$ and $[M^{\blacktriangleleft}(\mathbf{A})]$ are equal in the Grothendieck group of \mathcal{O} .*

We will mostly work just with the *natural order* $<$ on $\{1, \dots, m+n\}$, meaning of course that $1 < 2 < \dots < m+n$; for this order we denote $\mathfrak{b}^<, \lambda_{\mathbf{A}}^<, \rho^<, M^<(\mathbf{A})$ and $L^<(\mathbf{A})$ simply by $\mathfrak{b}, \lambda_{\mathbf{A}}, \rho, M(\mathbf{A})$ and $L(\mathbf{A})$. In particular \mathfrak{b} is the *standard Borel subalgebra* of upper triangular matrices in \mathfrak{g} . The resulting labelling of the irreducible objects of \mathcal{O} is the best choice for several other purposes. For example, the irreducible object $L(\mathbf{A})$ is *finite-dimensional* if and only if \mathbf{A} is *dominant*; hence, the irreducible objects $L(\mathbf{A})$ for all dominant tableaux \mathbf{A} give a complete set of inequivalent finite-dimensional irreducible \mathfrak{g} -supermodules. This was established originally by Kac in [K] by an argument involving parabolic induction from $\mathfrak{g}_{\bar{0}}$. In a similar way, one sees that $L(\mathbf{A})$ is of *maximal Gelfand–Kirillov dimension* amongst all supermodules in \mathcal{O} if and only if \mathbf{A} is *anti-dominant*.

The natural order on $\{1, \dots, m+n\}$ corresponds to the ordering of the boxes of the pyramid π induced by the lexicographic order of coordinates $(\text{row}(i), \text{col}(i))$, i.e. $i < j$ if and only if $\text{row}(i) < \text{row}(j)$, or $\text{row}(i) = \text{row}(j)$ and $\text{col}(i) < \text{col}(j)$. There is another normal order which plays a significant role for us, namely, the order $<'$ arising from the reverse lexicographic order on coordinates, i.e. $i < j$ if and only if $\text{col}(i) < \text{col}(j)$, or $\text{col}(i) = \text{col}(j)$ and $\text{row}(i) < \text{row}(j)$. For this order, we denote $\mathfrak{b}^<', \lambda_{\mathbf{A}}^<', \rho^<', M^<'(\mathbf{A})$ and $L^<'(\mathbf{A})$ by $\mathfrak{b}', \lambda'_{\mathbf{A}}, \rho', M'(\mathbf{A})$ and $L'(\mathbf{A})$. Note that in [BBG] the weight ρ' was denoted $\tilde{\rho}$.

Whereas the Borel subalgebra \mathfrak{b} arising from the natural ordering has a unique odd simple root, the Borel subalgebra \mathfrak{b}' has a maximal number of odd simple roots. This leads to some significant differences when working with the ordering $<'$ compared to the natural ordering. For instance, it is not so easy to describe the tableaux \mathbf{A} such that $L'(\mathbf{A})$ is either finite-dimensional or of maximal Gelfand–Kirillov dimension in purely combinatorial terms.

Using the description of ρ' given by (2.2) a direct calculation gives

$$(\rho', \delta_{m+1} + \dots + \delta_{m+n}) \equiv [(n-m)/2] + m s_- \pmod{2}. \quad (2.4)$$

Hence, recalling (2.1), we have that $\text{par}(\lambda'_A) = \text{par}(b(A))$ for $A \in \text{Tab}$.

2.4. The Harish-Chandra homomorphism. Let $Z(\mathfrak{g})$ denote the center of $U(\mathfrak{g})$. This can be understood via the Harish-Chandra homomorphism, which gives an isomorphism between $Z(\mathfrak{g})$ and a certain subalgebra $I(\mathfrak{t})$ of $S(\mathfrak{t})$. Let $x_i := e_{i,i}$ for $i = 1, \dots, m$ and $y_j := -e_{m+j, m+j}$ for $j = 1, \dots, n$, so that $S(\mathfrak{t}) = \mathbb{C}[x_1, \dots, x_m, y_1, \dots, y_n]$. The Weyl group of \mathfrak{g} with respect to \mathfrak{t} is the product of symmetric groups $S_m \times S_n$, which acts naturally on $S(\mathfrak{t})$ so that S_m permutes x_1, \dots, x_m and S_n permutes y_1, \dots, y_n . Then

$$I(\mathfrak{t}) := \left\{ f \in S(\mathfrak{t})^{S_m \times S_n} \mid \begin{array}{l} \frac{\partial f}{\partial x_i} + \frac{\partial f}{\partial y_j} \equiv 0 \pmod{x_i - y_j} \\ \text{for any } 1 \leq i \leq m, 1 \leq j \leq n \end{array} \right\}. \quad (2.5)$$

A distinguished set of generators for $I(\mathfrak{t})$ is given by the *elementary supersymmetric polynomials*

$$e_r(x_1, \dots, x_m / y_1, \dots, y_n) := \sum_{s+t=r} (-1)^t e_s(x_1, \dots, x_m) h_t(y_1, \dots, y_n) \quad (2.6)$$

for all $r \geq 1$, where $e_s(x_1, \dots, x_m)$ is the s th elementary symmetric polynomial and $h_t(y_1, \dots, y_n)$ is the t th complete symmetric polynomial; see e.g. [Se, §0.6.1].

To define the Harish-Chandra homomorphism itself we fix a total order \triangleleft on $\{1, \dots, m+n\}$. Recall that $\mathfrak{b}^\triangleleft$ is the Borel subalgebra spanned by $\{e_{i,j}\}_{i \triangleleft j}$; let $\mathfrak{n}^\triangleleft$ be its nilradical. Writing $U(\mathfrak{g})_0$ for the centralizer of \mathfrak{t} in $U(\mathfrak{g})$, let $\phi^\triangleleft : U(\mathfrak{g})_0 \rightarrow S(\mathfrak{t})$ be the algebra homomorphism defined by the projection along the direct sum decomposition $U(\mathfrak{g})_0 = S(\mathfrak{t}) \oplus (U(\mathfrak{g})_0 \cap U(\mathfrak{g})\mathfrak{n}^\triangleleft)$. Let

$$\text{HC} := S_{-\rho^\triangleleft} \circ \phi^\triangleleft : Z(\mathfrak{g}) \rightarrow S(\mathfrak{t}), \quad (2.7)$$

where the shift automorphism $S_{-\rho^\triangleleft}$ is the automorphism of $S(\mathfrak{t})$ defined by $x \mapsto x - \rho^\triangleleft(x)$ for each $x \in \mathfrak{t}$. Now we can state the key theorem here; see [M, §13.2] for a recent exposition of the proof.

Theorem 2.2 (Kac, Sergeev). *The homomorphism HC is an isomorphism between $Z(\mathfrak{g})$ and $I(\mathfrak{t})$.*

Our definition of the Harish-Chandra homomorphism involves the choice of the total order \triangleleft . But in fact one obtains the same isomorphism $Z(\mathfrak{g}) \xrightarrow{\sim} I(\mathfrak{t})$ no matter which order is chosen:

Theorem 2.3. *The map $\text{HC} : Z(\mathfrak{g}) \rightarrow S(\mathfrak{t})$ does not depend on the particular choice of the total order \triangleleft used in its definition.*

Proof. Suppose first that \triangleleft and \blacktriangleleft are two orders that are conjugate under $S_m \times S_n$, i.e. there exists a permutation $\sigma \in S_m \times S_n$ such that

$$i \triangleleft j \Leftrightarrow \sigma(i) \blacktriangleleft \sigma(j),$$

where S_m permutes $\{1, \dots, m\}$ and S_n permutes $\{m+1, \dots, m+n\}$. Let HC and HC' be the Harish-Chandra homomorphisms defined via \triangleleft and \blacktriangleleft , respectively. Take any $z \in Z(\mathfrak{g})$ and write it as $z = z_0 + z_1$ for $z_0 \in S(\mathfrak{t})$, $z_1 \in U(\mathfrak{g})_0 \cap U(\mathfrak{g})\mathfrak{n}^\triangleleft$. Identifying the Weyl group $S_m \times S_n$ with permutation matrices in the group $\text{GL}_m(\mathbb{C}) \times \text{GL}_n(\mathbb{C})$ in the obvious way, we get an action of $S_m \times S_n$ on \mathfrak{g} by conjugation. Since this is an action by inner automorphisms it fixes z ,

so we have that $z = \sigma(z) = \sigma(z_0) + \sigma(z_1)$ with $\sigma(z_0) \in S(\mathfrak{t})$ and $\sigma(z_1) \in U(\mathfrak{g})_0 \cap U(\mathfrak{g})\mathfrak{n}^\blacktriangleleft$. Now compute:

$$\mathrm{HC}'(z) = S_{-\rho^\blacktriangleleft}(\sigma(z_0)) = S_{-\sigma(\rho^\blacktriangleleft)}(\sigma(z_0)) = \sigma(S_{-\rho^\blacktriangleleft}(z_0)) = \sigma(\mathrm{HC}(z)) = \mathrm{HC}(z),$$

where the last equality follows because $\mathrm{HC}(z)$ is symmetric by Theorem 2.2.

Since any order is $S_m \times S_n$ -conjugate to a normal order, we have thus reduced the problem to showing that the Harish-Chandra homomorphisms arising from any two normal orders \triangleleft and \blacktriangleleft are equal. Again, let HC and HC' be the Harish-Chandra homomorphisms defined from the orders \triangleleft and \blacktriangleleft , respectively, assuming now that both orders are normal. For any $z \in Z(\mathfrak{g})$ and $\mathbf{A} \in \mathrm{Tab}$, the element z acts on the Verma supermodule $M^\triangleleft(\mathbf{A})$ (resp. $M^\blacktriangleleft(\mathbf{A})$) by the scalar $\mathrm{HC}(z)(\lambda_{\mathbf{A}})$ (resp. $\mathrm{HC}'(z)(\lambda_{\mathbf{A}})$). So to prove that $\mathrm{HC}(z) = \mathrm{HC}'(z)$ it suffices to show that $M^\triangleleft(\mathbf{A})$ and $M^\blacktriangleleft(\mathbf{A})$ have the same central character, which follows from Lemma 2.1. \square

There is an explicit formula for the elements $z_r \in Z(\mathfrak{g})$ lifting the elementary supersymmetric polynomials $e_r(x_1, \dots, x_m/y_1, \dots, y_n)$. To formulate this, recall from [GKLLRT] that the kk -quasideterminant of a $k \times k$ -matrix M is

$$|M|_{k,k} := d - ca^{-1}b,$$

assuming M is decomposed into block matrices as $M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ so that a is an invertible $(k-1) \times (k-1)$ matrix and d is a scalar. Working in the algebra $U(\mathfrak{g})[[u^{-1}]]$ where u is an indeterminate, let

$$\zeta_k(u) := u|T_k(u)|_{k,k},$$

where $T_k(u)$ is the $k \times k$ matrix with ij -entry $\delta_{i,j} + (-1)^{|i|}u^{-1}e_{i,j}$. The coefficients of these formal Laurent series for $k = 1, \dots, m+n$ generate a commutative subalgebra of $U(\mathfrak{g})$. Then set

$$z(u) = \sum_{r \geq 0} z_r u^{-r} := \prod_{k=1}^m \zeta_k(u+1-k) \Big/ \prod_{k=1}^n \zeta_{m+k}(u+k-m). \quad (2.8)$$

This defines elements $z_1, z_2, z_3, \dots \in U(\mathfrak{g})$. For example for $\mathfrak{gl}_{|1|}(\mathbb{C})$ one gets $z_1 = e_{1,1} + e_{2,2}$.

Theorem 2.4. *The elements $\{z_r\}_{r \geq 1}$ generate the center $Z(\mathfrak{g})$. Moreover,*

$$\mathrm{HC}(z_r) = e_r(x_1, \dots, x_m/y_1, \dots, y_n).$$

Proof. Let $Y(\mathfrak{g})$ be the Yangian of \mathfrak{g} and $b_{m|n}(u) \in Y(\mathfrak{g})[[u^{-1}]]$ be Nazarov's quantum Berezinian from [N]; see also [Gw, Definition 3.1]. In [Gw, Theorem 1], Gow establishes a remarkable factorization of this quantum Berezinian, from which we see that $z(u)$ is the image of $(u+1)^{-1}(u+2)^{-1} \cdots (u+n-m)^{-1}b_{m|n}(u)$ under the usual evaluation homomorphism $Y(\mathfrak{g}) \rightarrow U(\mathfrak{g})$. The coefficients of $b_{m|n}(u)$ are central in $Y(\mathfrak{g})$ by [N] (or [Gw, Theorem 2]). Hence, the coefficients z_1, z_2, \dots of our $z(u)$ are central in $U(\mathfrak{g})$.

It remains to compute $\mathrm{HC}(z_r)$. For this we use the definition of HC coming from the natural order $<$, since for this order it is clear how to apply the

projection $\phi = \phi^\triangleleft$ to each of the power series $\zeta_i(u)$. One gets that

$$\text{HC}(z(u)) = \prod_{k=1}^m (1 + u^{-1}x_k) \Big/ \prod_{k=1}^n (1 + u^{-1}y_k).$$

The u^{-r} -coefficient of this expression is equal to $e_r(x_1, \dots, x_m/y_1, \dots, y_n)$. \square

Remark 2.5. In fact there exist other factorizations of $z(u)$ analogous to (2.8) which are adapted to more general total orders \triangleleft . To explain, let \triangleleft be an arbitrary total order on $\{1, \dots, m+n\}$. Let $\sigma \in S_{m+n}$ be the permutation such that $\sigma(1) \triangleleft \sigma(2) \triangleleft \dots \triangleleft \sigma(m+n)$. Let

$$\zeta_{\sigma(k)}^\triangleleft(u) := u|T_k^\triangleleft(u)|_{k,k},$$

where $T_k^\triangleleft(u)$ is the $k \times k$ matrix with ij -entry $\delta_{i,j} + (-1)^{|\sigma(i)|} u^{-1} e_{\sigma(i), \sigma(j)}$. Then we have that

$$z(u) = \prod_{k=1}^m \zeta_k^\triangleleft(u + (\rho^\triangleleft, \delta_k)) \Big/ \prod_{k=1}^n \zeta_k^\triangleleft(u + (\rho^\triangleleft, \delta_k)).$$

This (and some analogous factorizations of the quantum Berezinian in $Y(\mathfrak{g})$) may be derived from [Gw, Theorem 1] by some explicit commutations in the super Yangian; we omit the details.

2.5. Projectives, prinjectives and blocks. The following *linkage principle* gives some rough information about the composition multiplicities of the Verma supermodules $M(\mathbf{A})$. This involves the Bruhat order \leq from §2.1.

Lemma 2.6. $[M(\mathbf{A}) : L(\mathbf{B})] \neq 0 \Rightarrow \mathbf{B} \leq \mathbf{A}$.

Proof. This is a consequence of the superalgebra analog of the Jantzen sum formula from [M, §10.3] or [Gk]; see [B5, Lemma 2.5] for details. \square

For $\mathbf{A} \in \text{Tab}$ we denote the projective cover of $L(\mathbf{A})$ in \mathcal{O} by $P(\mathbf{A})$. The supermodule $P(\mathbf{A})$ has a Verma flag, that is, a finite filtration with sections of the form $M(\mathbf{B})$ for $\mathbf{B} \in \text{Tab}$. Moreover, by *BGG reciprocity*, the multiplicity $(P(\mathbf{A}) : M(\mathbf{B}))$ of $M(\mathbf{B})$ in a Verma flag of $P(\mathbf{A})$ is equal to the composition multiplicity $[M(\mathbf{B}) : L(\mathbf{A})]$; see e.g. [B2]. Combined with Lemma 2.6, this implies that the category \mathcal{O} is a *highest weight category* with weight poset (Tab, \leq) . Of course its standard objects are the Verma supermodules $\{M(\mathbf{A})\}_{\mathbf{A} \in \text{Tab}}$.

Remark 2.7. In fact each choice of normal order \triangleleft on $\{1, \dots, m+n\}$ gives rise to a *different* structure of highest weight category on \mathcal{O} , with standard objects being the corresponding Verma supermodules $M^\triangleleft(\mathbf{A})$. In this article we only need the highest weight structure that comes from the natural order.

By a *prinjective object* we mean one that is both projective and injective. The following lemma classifies the prinjective objects in \mathcal{O} , showing that they are the projective covers of the irreducible objects of maximal Gelfand–Kirillov dimension.

Lemma 2.8. *Let $\mathbf{A} \in \text{Tab}$. Then $P(\mathbf{A})$ is prinjective if and only if \mathbf{A} is anti-dominant. In that case $P(\mathbf{A})$ is both the projective cover and the injective hull of $L(\mathbf{A})$.*

Proof. This is a consequence of [BLW, Theorem 2.22] (bearing in mind also [BLW, Theorem 3.10]); see also [B5, Lemma 4.3, Remark 4.4] and [CS, Corollary 6.2(ii)]. \square

Recall finally that \approx is the equivalence relation on Tab generated by the Bruhat order. For a linkage class $\xi \in \text{Tab}/\approx$, we let \mathcal{O}_ξ be the Serre subcategory of \mathcal{O} generated by $\{L(\mathbf{A})\}_{\mathbf{A} \in \xi}$. Lemma 2.6 implies immediately that this is a sum of blocks of \mathcal{O} . In fact each \mathcal{O}_ξ is an indecomposable block, thanks to [CMW, Theorem 3.12]. Thus the blocks of \mathcal{O} are in bijection with the linkage classes.

Lemma 2.9. *Let $S_m \times S_n$ act on Tab by permuting entries within rows. Suppose we are given a linkage class $\xi \in \text{Tab}/\approx$ and simple transposition $\sigma \in S_m \times S_n$ such that $\sigma(\xi) := \{\sigma(\mathbf{A}) \mid \mathbf{A} \in \xi\}$ is a different linkage class to ξ . Then, there is an equivalence of categories $T_\sigma : \mathcal{O}_\xi \rightarrow \mathcal{O}_{\sigma(\xi)}$ such that $T_\sigma(M(\mathbf{A})) \cong M(\sigma(\mathbf{A}))$ and $T_\sigma(L(\mathbf{A})) \cong L(\sigma(\mathbf{A}))$ for each $\mathbf{A} \in \xi$.*

Proof. This is a reformulation of [CMW, Proposition 3.9], where the equivalence T_σ is constructed explicitly as a certain twisting functor. \square

3. PRINCIPAL W -ALGEBRAS AND WHITTAKER FUNCTORS

After reviewing some basic definitions and results from [BBG], we proceed to introduce the Whittaker coinvariants functor H_0 , which takes representations of \mathfrak{g} to representations of its *principal W -algebra*, i.e. the (finite) W -algebra associated to a principal nilpotent orbit $e \in \mathfrak{g}$. We will mainly be concerned with the restriction of this functor to the category \mathcal{O} . The main result of the section shows that H_0 sends $M(\mathbf{A})$ to the corresponding Verma supermodule $\overline{M}(\mathbf{A})$ for W ; up to a parity shift, the latter was already introduced in [BBG]. This has several important consequences: we use it to determine the composition multiplicities of each $\overline{M}(\mathbf{A})$, to show that H_0 sends irreducibles in \mathcal{O} to irreducibles or zero, and to describe the center of W explicitly.

3.1. The principal W -superalgebra. We continue with $\mathfrak{g} = \mathfrak{gl}_{m|n}(\mathbb{C})$ as in the previous section. Consider the principal nilpotent element

$$e := e_{1,2} + e_{2,3} + \cdots + e_{m-1,m} + e_{m+1,m+2} + e_{m+2,m+3} + \cdots + e_{m+n-1,m+n} \in \mathfrak{g}.$$

Define a good grading $\mathfrak{g} = \bigoplus_{r \in \mathbb{Z}} \mathfrak{g}(r)$ for $e \in \mathfrak{g}(1)$ by declaring that each matrix unit $e_{i,j}$ is of degree

$$\deg(e_{i,j}) := \text{col}(j) - \text{col}(i). \quad (3.1)$$

Set

$$\mathfrak{p} := \bigoplus_{r \geq 0} \mathfrak{g}(r), \quad \mathfrak{h} := \mathfrak{g}(0), \quad \mathfrak{m} := \bigoplus_{r < 0} \mathfrak{g}(r).$$

Let $\chi \in \mathfrak{g}^*$ be defined by $\chi(x) := (x, e)$. The restriction of χ to \mathfrak{m} is a character of \mathfrak{m} . Then define $\mathfrak{m}_\chi := \{x - \chi(x) \mid x \in \mathfrak{m}\}$, which is a shifted copy of \mathfrak{m} inside $U(\mathfrak{m})$. The *principal W -superalgebra* may then be defined as

$$W := \{u \in U(\mathfrak{p}) \mid u \mathfrak{m}_\chi \subseteq \mathfrak{m}_\chi U(\mathfrak{g})\}, \quad (3.2)$$

which is a subalgebra of $U(\mathfrak{p})$. Although this definition depends implicitly on the choice of pyramid π , the isomorphism type of W depends only on m and

n not π , see [BBG, Remark 4.8]. The following theorem shows that W is isomorphic to a truncated shifted version of the Yangian $Y(\mathfrak{gl}_{1|1})$.

Theorem 3.1 ([BBG, Theorem 4.5]). *The superalgebra W contains distinguished even elements $\{d_1^{(r)}, d_2^{(r)}\}_{r \geq 0}$ and odd elements $\{e^{(r)}\}_{r > s_+} \cup \{f^{(r)}\}_{r > s_-}$. These elements generate W subject only to the following relations:*

$$\begin{aligned} d_i^{(0)} &= 1, & d_1^{(r)} &= 0 \text{ for } r > m, \\ [d_i^{(r)}, d_j^{(s)}] &= 0, & [e^{(r)}, f^{(s)}] &= \sum_{a=0}^{r+s-1} \tilde{d}_1^{(a)} d_2^{(r+s-1-a)}, \\ [e^{(r)}, e^{(s)}] &= 0, & [d_i^{(r)}, e^{(s)}] &= \sum_{a=0}^{r-1} d_i^{(a)} e^{(r+s-1-a)}, \\ [f^{(r)}, f^{(s)}] &= 0, & [d_i^{(r)}, f^{(s)}] &= - \sum_{a=0}^{r-1} f^{(r+s-1-a)} d_i^{(a)}, \end{aligned}$$

where $\tilde{d}_i^{(r)}$ is defined recursively from $\sum_{a=0}^r \tilde{d}_i^{(a)} d_i^{(r-a)} = \delta_{r,0}$.

We will occasionally need to appeal to the explicit formulae² for the generators $d_i^{(r)}$, $e^{(s)}$ and $f^{(s)}$ from [BBG, §4]. In particular, these formulae show that $d_1^{(1)}$ and $d_2^{(1)}$ are the unique elements of $S(\mathfrak{t}) = \mathbb{C}[\mathfrak{t}^*]$ such that

$$d_1^{(1)}(\lambda) = (\lambda + \rho', \delta_1 + \cdots + \delta_m), \quad (3.3)$$

$$d_2^{(1)}(\lambda) = (\lambda + \rho', \delta_{m+1} + \cdots + \delta_{m+n}). \quad (3.4)$$

It is also often useful to work with the generating functions

$$d_i(u) := \sum_{r \geq 0} d_i^{(r)} u^{-r} \in W[[u^{-1}]], \quad \tilde{d}_i(u) := \sum_{r \geq 0} \tilde{d}_i^{(r)} u^{-r} \in W[[u^{-1}]],$$

so that $\tilde{d}_i(u) = d_i(u)^{-1}$. Using these, we can define more elements $\{c^{(r)}, \tilde{c}^{(r)}\}_{r \geq 0}$ by setting

$$c(u) = \sum_{r \geq 0} c^{(r)} u^{-r} := \tilde{d}_1(u) d_2(u), \quad \tilde{c}(u) = \sum_{r \geq 0} \tilde{c}^{(r)} u^{-r} := d_1(u) \tilde{d}_2(u). \quad (3.5)$$

In particular, by the defining relations, we have that $c^{(r+s-1)} = [e^{(r)}, f^{(s)}]$ for $r > s_+$, $s > s_-$. The elements $\{c^{(r)}\}_{r \geq 1}$ are known to belong to the center $Z(W)$; see [BBG, Remark 2.3]. Hence, so too do the elements $\{\tilde{c}^{(r)}\}_{r \geq 1}$. We will show in Corollary 3.22 below that either of these families of elements give generators for $Z(W)$.

Recall finally by [BBG, Theorem 6.1] that W has a triangular decomposition: let W^0 , W^+ and W^- be the subalgebras generated by $\{d_1^{(r)}, d_2^{(s)}\}_{1 \leq r \leq m, 1 \leq s \leq n}$, $\{e^{(r)}\}_{s_+ < r \leq s_+ + m}$ and $\{f^{(r)}\}_{s_- < r \leq s_- + m}$, respectively; then the multiplication map $W^- \otimes W^0 \otimes W^+ \rightarrow W$ is a vector space isomorphism. Moreover, by the PBW theorem for W , the subalgebra W^0 is a free polynomial algebra of rank $m + n$, while W^+ and W^- are Grassmann algebras of dimension 2^m .

²There is a typo in [BBG, (4.12)–(4.13)]; both of these formulae need an extra minus sign. Similarly the formulae for $\tilde{d}_i^{(r)}$ in [BBG, (4.19)–(4.20)] need to be changed by a sign.

3.2. Highest weight theory for W . Next, we review some results about the representation theory of W established in [BBG]. The triangular decomposition allows us to define Verma supermodules for W as follows. Let $W^\# := W^0W^+$. This is a subalgebra of W , and there is a surjective homomorphism $W^\# \rightarrow W^0$ which is the identity on W^0 and zero on each $e^{(r)} \in W^+$.

Given $\mathbf{A} = \begin{smallmatrix} a_1 \cdots a_m \\ b_1 \cdots b_n \end{smallmatrix} \in \text{Tab}$, let $\mathbb{C}_{\mathbf{A}}$ be the one-dimensional W^0 -supermodule spanned by a vector $\bar{1}_{\mathbf{A}}$ of parity $\text{par}(b(\mathbf{A}))$, such that

$$d_1^{(r)}\bar{1}_{\mathbf{A}} = e_r(a_1, \dots, a_m)\bar{1}_{\mathbf{A}}, \quad d_2^{(s)}\bar{1}_{\mathbf{A}} = e_s(b_1, \dots, b_n)\bar{1}_{\mathbf{A}} \quad (3.6)$$

for $1 \leq r \leq m, 1 \leq s \leq n$. View $\mathbb{C}_{\mathbf{A}}$ as a $W^\#$ -supermodule via the surjection $W^\# \rightarrow W^0$. Then induce to form the *Verma supermodule*

$$\bar{M}(\mathbf{A}) := W \otimes_{W^\#} \mathbb{C}_{\mathbf{A}}, \quad (3.7)$$

setting $\bar{m}_{\mathbf{A}} := 1 \otimes \bar{1}_{\mathbf{A}}$. Of course, $\bar{M}(\mathbf{A})$ only depends on the row equivalence class of \mathbf{A} . The PBW theorem for W implies that $\dim \bar{M}(\mathbf{A}) = 2^m$.

We say that $M \in W\text{-smod}$ is a *highest weight supermodule* of *highest weight* $\mathbf{A} = \begin{smallmatrix} a_1 \cdots a_m \\ b_1 \cdots b_n \end{smallmatrix} \in \text{Tab}$ if there exists a homogeneous vector $v \in M$ that generates M as a W -supermodule with $e^{(r)}v = 0$ for $r > s_+$, $d_1^{(r)}v = e_r(a_1, \dots, a_m)v$ for $1 \leq r \leq m$, and $d_2^{(s)}v = e_s(b_1, \dots, b_n)v$ for $1 \leq s \leq n$. The Verma supermodule $\bar{M}(\mathbf{A})$ is the *universal highest weight supermodule* of highest weight \mathbf{A} : given any highest weight supermodule M of highest weight \mathbf{A} as above, there exists a unique surjective homomorphism from either $\bar{M}(\mathbf{A})$ or $\Pi\bar{M}(\mathbf{A})$ onto M such that $\bar{m}_{\mathbf{A}} \mapsto v$; the homomorphism is from $\bar{M}(\mathbf{A})$ if and only if $|v| = \text{par}(b(\mathbf{A}))$.

By [BBG, Lemma 7.1], each $\bar{M}(\mathbf{A})$ has a unique irreducible quotient $\bar{L}(\mathbf{A})$.

Theorem 3.2 ([BBG, Theorem 7.2]). *The supermodules $\{\bar{L}(\mathbf{A})\}_{\mathbf{A} \in \text{Tab}}$ give all of the irreducible W -supermodules (up to isomorphism and parity switch). Moreover, $\bar{L}(\mathbf{A}) \cong \bar{L}(\mathbf{B})$ if and only if $\mathbf{A} \sim \mathbf{B}$.*

In particular, the theorem shows that all irreducible W -supermodules are finite-dimensional. Henceforth, we will restrict our attention to the full subcategory $W\text{-smod}_{\text{fd}}$ of $W\text{-smod}$ consisting of finite-dimensional supermodules.

There is actually a very simple way to realize $\bar{L}(\mathbf{A})$ explicitly. Recall that

$$\mathfrak{h} \cong \mathfrak{gl}_1(\mathbb{C})^{\oplus s_-} \oplus \mathfrak{gl}_{1|1}(\mathbb{C})^{\oplus m} \oplus \mathfrak{gl}_1(\mathbb{C})^{\oplus s_+}. \quad (3.8)$$

For any $\mathbf{A} = \begin{smallmatrix} a_1 \cdots a_m \\ b_1 \cdots b_n \end{smallmatrix} \in \text{Tab}$, let $K(\mathbf{A})$ be the \mathfrak{h} -supermodule induced from a one-dimensional $\mathfrak{b}' \cap \mathfrak{h}$ -supermodule of weight $\lambda'_{\mathbf{A}}$ and parity $\text{par}(\lambda'_{\mathbf{A}}) = \text{par}(b(\mathbf{A}))$, cf. (2.4); we use the letter K here because it is a Kac supermodule for \mathfrak{h} (as well as being a Verma supermodule). Note that $\dim K(\mathbf{A}) = 2^m$. We denote the highest weight vector in $K(\mathbf{A})$ by $k_{\mathbf{A}}$. Observe that

$$M'(\mathbf{A}) \cong U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} K(\mathbf{A}). \quad (3.9)$$

Also let $V(\mathbf{A})$ be the unique irreducible quotient of $K(\mathbf{A})$. Thus $V(\mathbf{A})$ is an irreducible \mathfrak{h} -supermodule of $\mathfrak{b}' \cap \mathfrak{h}$ -highest weight $\lambda'_{\mathbf{A}}$, and $\dim V(\mathbf{A}) = 2^{m - \text{def}(\mathbf{A})}$. Finally, using the (injective!) homomorphism $W \hookrightarrow U(\mathfrak{p}) \twoheadrightarrow U(\mathfrak{h})$ derived from the natural inclusion and projection maps, we restrict these supermodules to W to obtain

$$\bar{K}(\mathbf{A}) := K(\mathbf{A}) \downarrow_W^{U(\mathfrak{h})}, \quad \bar{V}(\mathbf{A}) := V(\mathbf{A}) \downarrow_W^{U(\mathfrak{h})}. \quad (3.10)$$

We sometimes denote $k_A \in \overline{K}(A)$ instead by \overline{k}_A .

Theorem 3.3. *If $\text{def}(A) = \text{atyp}(A)$ then $\overline{L}(A) \cong \overline{V}(A)$.*

Proof. This is essentially [BBG, Theorem 8.4], but we should note also that the isomorphism constructed is necessarily even since it sends \overline{m}_A to \overline{k}_A , which are both of the same parity $\text{par}(b(A))$. \square

Lemma 3.4. *For any $A \in \text{Tab}$, we have that $[\overline{K}(A)] = \sum_{B \cup A} [\overline{V}(B)]$, equality written in the Grothendieck group $K_0(W\text{-smod}_{\text{fd}})$.*

Proof. By the representation theory of $\mathfrak{gl}_{1|1}(\mathbb{C})$, we have that

$$[K(A)] = \sum_{B \cup A} [V(B)].$$

The lemma follows from this on restricting to W . \square

3.3. Invariants and coinvariants. Given a right \mathfrak{g} -supermodule M , it is easy to check from (3.2) that the subspace

$$H^0(M) := H^0(\mathfrak{m}_\chi, M) = \{v \in M \mid v \mathfrak{m}_\chi = 0\} \quad (3.11)$$

is stable under right multiplication by elements of W . Hence, we obtain the *Whittaker invariants functor*

$$H^0 : \text{smod-}U(\mathfrak{g}) \rightarrow \text{smod-}W. \quad (3.12)$$

Let $\text{smod}_\chi\text{-}U(\mathfrak{g})$ be the full subcategory of $\text{smod-}U(\mathfrak{g})$ consisting of all the supermodules on which \mathfrak{m}_χ acts locally nilpotently. The super analog of *Skryabin's theorem* asserts that the restriction of H^0 defines an equivalence of categories from $\text{smod}_\chi\text{-}U(\mathfrak{g})$ to $\text{smod-}W$. Let Q denote the $(W, U(\mathfrak{g}))$ -superbimodule

$$Q := U(\mathfrak{g})/\mathfrak{m}_\chi U(\mathfrak{g}), \quad (3.13)$$

denoting the canonical image of $1 \in U(\mathfrak{g})$ in Q by 1_χ . Then the functor

$$- \otimes_W Q : \text{smod-}W \rightarrow \text{smod}_\chi\text{-}U(\mathfrak{g}) \quad (3.14)$$

is the inverse functor to H^0 in Skryabin's theorem. As observed already in [Z, Remarks 3.9–3.10], Skryabin's proof of this result in the purely even setting from [Sk] extends routinely to the super case. Along the way, one sees that Q is a free left W -supermodule with an explicitly constructed basis, from which we see that there exists a W -supermodule homomorphism

$$p : Q \twoheadrightarrow W \quad (3.15)$$

such that $p(1_\chi) = 1$. We fix such a choice for later use.

Instead, suppose that M is a left \mathfrak{g} -supermodule. Then again it is clear from (3.2) that the left action of W leaves the subspace $\mathfrak{m}_\chi M$ invariant, hence, we get induced a well-defined left action of W on

$$H_0(M) := H_0(\mathfrak{m}_\chi, M) = M/\mathfrak{m}_\chi M. \quad (3.16)$$

This gives us the *Whittaker coinvariants functor*

$$H_0 : U(\mathfrak{g})\text{-smod} \rightarrow W\text{-smod}. \quad (3.17)$$

Equivalently, this is the functor $Q \otimes_W -$.

The first lemma below connects Whittaker invariants and coinvariants. To formulate it we need some duals: if M is a left supermodule over some superalgebra then we write M^* for the full linear dual of M considered as a right supermodule with the obvious action $(fv)(a) = f(va)$ (no signs!). Similarly, we write *M for the dual of a right supermodule, which is a left supermodule. There are natural supermodule homomorphisms $M \rightarrow ({}^*M)^*$ and $M \rightarrow {}^*(M^*)$ (which involve a sign!). Note also that if V is a finite-dimensional superspace and M is arbitrary then the canonical maps

$$M^* \otimes V^* \xrightarrow{\sim} (V \otimes M)^*, \quad {}^*M \otimes {}^*V \xrightarrow{\sim} {}^*(V \otimes M) \quad (3.18)$$

are isomorphisms.

Lemma 3.5. *Let M be a left \mathfrak{g} -supermodule. Then there is a functorial isomorphism $H_0(M)^* \cong H^0(M^*)$. In particular, if $H_0(M)$ is finite-dimensional, then $H_0(M) \cong {}^*H^0(M^*)$.*

Proof. The natural inclusion $H_0(M)^* \hookrightarrow M^*$ induced by $M \rightarrow H_0(M)$ has image contained in $H^0(M^*)$. This gives a W -supermodule homomorphism $H_0(M)^* \hookrightarrow H^0(M^*)$. To see that it is surjective, we observe that any element of $H^0(M^*) \subseteq M^*$ annihilates $\mathfrak{m}_\chi M$, hence, comes from an element of $H_0(M)^*$. \square

The next lemma is an analog of another well-known result in the even setting.

Lemma 3.6. *The functor H_0 sends short exact sequences of left \mathfrak{g} -supermodules that are finitely generated over \mathfrak{m} to short exact sequences of finite-dimensional left W -supermodules.*

Proof. For any left \mathfrak{m} -supermodule M , we introduce its χ -restricted dual

$$M^\# := \{f \in M^* \mid f(\mathfrak{m}_\chi^r M) = 0 \text{ for } r \gg 0\}.$$

This defines a functor $(-)^{\#} : U(\mathfrak{m})\text{-smod} \rightarrow \text{smod-}U(\mathfrak{m})$. We claim that this functor is exact. To see this, we note as in the proof of [B4, Lemma 3.10] that the functor $(-)^{\#}$ is isomorphic to $\text{Hom}_{\mathfrak{m}}(-, E_\chi)$, where $E_\chi := U(\mathfrak{m})^{\#}$ viewed as an $(\mathfrak{m}, \mathfrak{m})$ -superbimodule in the obvious way. The proof of [Sk, Assertion 2] shows that E_χ is injective as a left \mathfrak{m} -supermodule; this follows ultimately from the non-commutative Artin-Rees lemma. The desired exactness follows.

If M is a left \mathfrak{g} -supermodule then $M^\#$ is actually a \mathfrak{g} -submodule of M^* , and this submodule belongs to $\text{smod}_\chi\text{-}U(\mathfrak{g})$. Hence, $(-)^{\#}$ can also be viewed as an exact functor $U(\mathfrak{g})\text{-smod} \rightarrow \text{smod}_\chi\text{-}U(\mathfrak{g})$. As in [B4, Lemma 3.11], we have quite obviously for any left \mathfrak{g} -supermodule that $H^0(M^*) = H^0(M^\#)$ as subspaces of M^* . Since H^0 is exact on $\text{smod}_\chi\text{-}U(\mathfrak{g})$ by Skryabin's theorem, we have now proved that the functor $U(\mathfrak{g})\text{-smod} \rightarrow \text{smod-}W$ given by $M \mapsto H^0(M^*)$ is exact. Finally, if M is a left \mathfrak{g} -supermodule that is finitely generated over \mathfrak{m} , then it is clear that $H_0(M)$ is finite-dimensional, so that $H_0(M) \cong {}^*H^0(M^*)$ by Lemma 3.5. The lemma follows. \square

Corollary 3.7. *The restriction of the functor H_0 to the category \mathcal{O} from §2.3 is exact and has image contained in $W\text{-smod}_{\text{fd}}$.*

Proof. In view of the lemma, it just remains to observe that all supermodules in \mathcal{O} are finitely generated over \mathfrak{m} . This follows because the Verma supermodules

$M(\mathbf{A})$ are finitely generated over \mathfrak{m} , which is easily seen from the definition since $\mathfrak{g}_{\bar{0}} = \mathfrak{b}_{\bar{0}} \oplus \mathfrak{m}_{\bar{0}}$. \square

We also need the following lemma which takes care of all the necessary book-keeping regarding parities.

Lemma 3.8. *For $M \in \mathcal{O}$, the elements $d_1^{(1)}, d_2^{(1)} \in W^0$ act semisimply on $H_0(M)$. Moreover, the z -eigenspace of $d_2^{(1)}$ is concentrated in parity $\text{par}(z)$ for each $z \in \mathbb{C}$.*

Proof. For $M \in \mathcal{O}$ and any vector $v \in M$ of \mathfrak{t} -weight λ , we know from (3.3)–(3.4) that $d_1^{(1)}$ and $d_2^{(1)}$ act on v (hence, $v + \mathfrak{m}_{\chi}M$) by the scalars $(\lambda + \rho', \delta_1 + \cdots + \delta_n)$ and $(\lambda + \rho', \delta_{m+1} + \cdots + \delta_{m+n})$, respectively. Hence, they both act semisimply on all of $H_0(M)$. For the last part, let $z := (\lambda + \rho', \delta_{m+1} + \cdots + \delta_{m+n})$. Then, by (2.1) and (2.4), the vector v is of parity $\text{par}(\lambda) = \text{par}(z)$. \square

3.4. On tensoring with finite-dimensional representations. Let $\text{Rep}(\mathfrak{g})$ be the symmetric monoidal category of *rational representations* of \mathfrak{g} , that is, finite-dimensional left \mathfrak{g} -supermodules which are semisimple over \mathfrak{t} with weights lying in $\mathfrak{t}_{\mathbb{Z}}^* := \bigoplus_{i=1}^{m+n} \mathbb{Z}\delta_i$. Tensoring with $V \in \text{Rep}(\mathfrak{g})$ defines a *projective functor*

$$V \otimes - : U(\mathfrak{g})\text{-smod} \rightarrow U(\mathfrak{g})\text{-smod}.$$

This is a rigid object of the strict monoidal category $\text{End}(U(\mathfrak{g})\text{-smod})$ of \mathbb{C} -linear endofunctors of $U(\mathfrak{g})\text{-smod}$: it has a biadjoint defined by tensoring with the usual dual V^\vee of V in the category $\text{Rep}(\mathfrak{g})$. In this subsection, we introduce an analogous biadjoint pair of endofunctors $V \circledast -$ and $V^\vee \circledast -$ of $W\text{-smod}_{\text{fd}}$. We will use the language of module categories over monoidal categories, see e.g. [EGNO, Chapter 7].

It is convenient to start by working with right supermodules. From the previous subsection, we recall the notation M^* and $*M$ for duals of left (resp. right) supermodules, which are right (resp. left) supermodules. In particular, for V as in the previous paragraph, V^* is a right $U(\mathfrak{g})$ -supermodule. Tensoring with it gives us an exact functor $- \otimes V^* : \text{smod-}U(\mathfrak{g}) \rightarrow \text{smod-}U(\mathfrak{g})$. In fact, writing $\text{End}(\text{smod-}U(\mathfrak{g}))$ for the strict monoidal category of all \mathbb{C} -linear endofunctors of $\text{smod-}U(\mathfrak{g})$, this defines a monoidal functor

$$\text{Rep}(\mathfrak{g})^{\text{op}} \rightarrow \text{End}(\text{smod-}U(\mathfrak{g})), \quad V \mapsto - \otimes V^*.$$

In other words, $\text{smod-}U(\mathfrak{g})$ is a right module category over the monoidal category $\text{Rep}(\mathfrak{g})$. The main coherence map that is needed for this comes from the natural isomorphisms $(- \otimes W^*) \circ (- \otimes V^*) \cong - \otimes (V^* \otimes W^*) \cong - \otimes ((W \otimes V)^*)$.

It is clear that $- \otimes V^*$ takes objects of $\text{smod}_{\chi}\text{-}U(\mathfrak{g})$ to objects of $\text{smod}_{\chi}\text{-}U(\mathfrak{g})$. Hence, we can consider $- \otimes V^*$ also as an endofunctor of $\text{smod}_{\chi}\text{-}U(\mathfrak{g})$. Transporting this through Skryabin's equivalence from (3.14), we obtain an exact functor

$$- \circledast V^* := H^0((- \otimes_W Q) \otimes V^*) : \text{smod-}W \rightarrow \text{smod-}W$$

Like in the previous paragraph, this actually defines a monoidal functor

$$\text{Rep}(\mathfrak{g})^{\text{op}} \rightarrow \text{End}(\text{smod-}W),$$

making $\text{smod-}W$ into a right module category over $\text{Rep}(\mathfrak{g})$. To construct the coherence map $(- \circledast W^*) \circ (- \circledast V^*) \cong - \circledast ((W \otimes V)^*)$ for this, one needs to use

the canonical adjunction between $-\otimes_W Q$ and H^0 . Perhaps the most important fact about this functor is that there is an isomorphism of vector superspaces

$$M \circledast V^* \xrightarrow{\sim} M \otimes V^* \quad (3.19)$$

which is natural in both M and V . In particular, $-\circledast V^*$ takes finite-dimensional W -supermodules to finite-dimensional W -supermodules. By definition, the isomorphism (3.19) is defined by the restriction of the map

$$\begin{aligned} (M \otimes_W Q) \otimes V^* &\rightarrow M \otimes V^*, \\ (m \otimes 1_\chi u) \otimes f &\mapsto m p(u) \otimes f \end{aligned}$$

where p is the map from (3.15). The proof of this assertion goes back to the PhD thesis of Lynch. For this and other details about this construction, we refer to [BK1, §8.2]; the super case is essentially the same.

Lemma 3.9. *There is an isomorphism $H^0(M) \circledast V^* \cong H^0(M \otimes V^*)$ which is natural in M and V . It makes $H^0 : \text{smod-}U(\mathfrak{g}) \rightarrow \text{smod-}W$ into a morphism of right $\text{Rep}(\mathfrak{g})$ -module categories.*

Proof. We start from the canonical isomorphism $M \cong H^0(M) \otimes_W Q$ defined by the canonical adjunction from Skryabin's theorem. Then apply $H^0 \circ (- \otimes V^*)$ to both sides. \square

We are ready to switch the discussion to left supermodules. For $V \in \text{Rep}(\mathfrak{g})$ as before and $M \in W\text{-smod}_{\text{fd}}$, we define

$$V \circledast M := {}^*(M^* \circledast V^*), \quad (3.20)$$

noting that $M^* \circledast V^*$ is also finite-dimensional thanks to (3.19). Again, we have that $(W \circledast -) \circ (V \circledast -) \cong (W \otimes V) \circledast -$, so that we obtain a monoidal functor

$$\text{Rep}(\mathfrak{g}) \rightarrow \text{End}(W\text{-smod}_{\text{fd}}), \quad V \mapsto V \circledast - \quad (3.21)$$

making $W\text{-smod}_{\text{fd}}$ into a (left) module category over $\text{Rep}(\mathfrak{g})$. Also, applying ${}^*(-)$ to (3.19) with M replaced by M^* then using (3.18), we get a canonical isomorphism

$$V \otimes M \cong {}^*(V^*) \otimes {}^*(M^*) \cong {}^*(M^* \otimes V^*) \xrightarrow{\sim} {}^*(M^* \circledast V^*) = V \circledast M \quad (3.22)$$

as vector superspaces.

In general, due to the parity condition prescribed by (2.1), the endofunctor $V \otimes -$ does not leave \mathcal{O} invariant. However, it does providing the λ -weight space of V is concentrated in parity $\text{par}((\lambda, \delta_{m+1} + \cdots + \delta_{m+n}))$ for all $\lambda \in \mathfrak{t}_{\mathbb{Z}}^*$. Let $\text{Rep}_0(\mathfrak{g})$ be the full monoidal subcategory of $\text{Rep}(\mathfrak{g})$ consisting of all such V . Then, for $V \in \text{Rep}_0(\mathfrak{g})$, we do get a monoidal functor

$$\text{Rep}_0(\mathfrak{g}) \rightarrow \text{End}(\mathcal{O}), \quad V \mapsto V \otimes -. \quad (3.23)$$

So \mathcal{O} is a module category over $\text{Rep}_0(\mathfrak{g})$.

Theorem 3.10. *There is a natural isomorphism $H_0(V \otimes M) \cong V \circledast H_0(M)$ making $H_0 : \mathcal{O} \rightarrow W\text{-smod}_{\text{fd}}$ into a morphism of $\text{Rep}_0(\mathfrak{g})$ -module categories.*

Proof. Take $M \in W\text{-smod}_{\text{fd}}$. Since $H_0(V \otimes M)$ and V are finite-dimensional, we have from Lemmas 3.5 and 3.9 that

$$\begin{aligned} H_0(V \otimes M) &\cong {}^*H^0((V \otimes M)^*) \cong {}^*H^0(M^* \otimes V^*) \\ &\cong {}^*(H^0(M^*) \otimes V^*) = V \otimes H_0(M). \end{aligned}$$

Everything else is purely formal; see [BK1, §8.4] for further discussion. \square

Since V^\vee is both a left dual and right dual to $V \in \text{Rep}_0(\mathfrak{g})$, it is automatic that $V^\vee \otimes -$ is both left and right adjoint to $V \otimes -$. Moreover, the monoidal isomorphism described in Theorem 3.10 intertwines the canonical adjunctions between $V \otimes -$ and $V^\vee \otimes -$ with the ones between $V \otimes -$ and $V^\vee \otimes -$.

3.5. Whittaker coinvariants of $M'(\mathbf{A})$. The following theorem will allow us to determine the effect of the Whittaker coinvariants functor on the Verma supermodule $M'(\mathbf{A})$.

Theorem 3.11. *The map $U(\mathfrak{p}) \rightarrow Q, u \mapsto 1_\chi u$ is an isomorphism of $(W, U(\mathfrak{p}))$ -superbimodules. Hence, for any left \mathfrak{p} -supermodule M , there is an isomorphism*

$$M \downarrow_W^{U(\mathfrak{p})} \xrightarrow{\sim} H_0(U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} M), \quad v \mapsto 1 \otimes v + \mathfrak{m}_\chi(U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} M)$$

Proof. The first assertion is immediate from the PBW theorem and the definition of Q . Hence,

$$H_0(U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} M) \cong Q \otimes_{U(\mathfrak{g})} U(\mathfrak{g}) \otimes_{U(\mathfrak{p})} M \cong Q \otimes_{U(\mathfrak{p})} M \cong U(\mathfrak{p}) \otimes_{U(\mathfrak{p})} M \cong M,$$

which translates into the given isomorphism. \square

Recall the definition of the W -supermodule $\overline{K}(\mathbf{A})$ from §3.2.

Corollary 3.12. *For any $\mathbf{A} \in \text{Tab}$, we have that $H_0(M'(\mathbf{A})) \cong \overline{K}(\mathbf{A})$.*

Proof. Apply Theorem 3.11 to the \mathfrak{p} -supermodule obtained by inflating $K(\mathbf{A})$ through $\mathfrak{p} \rightarrow \mathfrak{h}$ and use (3.9). \square

Corollary 3.13. *For any $\mathbf{A} \in \text{Tab}$, there exists a supermodule $M \in \mathcal{O}$ such that $H_0(M) \cong \overline{L}(\mathbf{A})$.*

Proof. Since $\overline{L}(\mathbf{A})$ only depends on the row equivalence class of \mathbf{A} , we may assume that $\text{atyp}(\mathbf{A}) = \text{def}(\mathbf{A})$. Then apply Theorem 3.11 to the \mathfrak{p} -supermodule obtained by inflating $V(\mathbf{A})$ through $\mathfrak{p} \rightarrow \mathfrak{h}$ and use Theorem 3.3. \square

3.6. Whittaker coinvariants of $M(\mathbf{A})$. We regard the following theorem as one of the central results of this article.

Theorem 3.14. *For any $\mathbf{A} \in \text{Tab}$, we have that $H_0(M(\mathbf{A})) \cong \overline{M}(\mathbf{A})$.*

Proof. See Appendix A. \square

Corollary 3.15. *For any $\mathbf{A} \in \text{Tab}$, we have that $[\overline{M}(\mathbf{A})] = [\overline{K}(\mathbf{A})]$ in the Grothendieck group $K_0(W\text{-smod}_{\text{fd}})$.*

Proof. By Lemmas 2.1 and 3.6 we have that $[H_0(M(\mathbf{A}))] = [H_0(M'(\mathbf{A}))]$. Now apply Theorem 3.14 and Corollary 3.12. \square

Corollary 3.16. *Suppose $\mathbf{A} \in \text{Tab}$ is chosen so that $\text{def}(\mathbf{A}) = \text{atyp}(\mathbf{A})$. Then*

$$[\overline{M}(\mathbf{A})] = \sum_{\mathbf{B} \cup \mathbf{A}} [\overline{L}(\mathbf{B})].$$

Proof. This is immediate from Corollary 3.15, Lemma 3.4 and Theorem 3.3. \square

3.7. Whittaker coinvariants of $L(\mathbf{A})$. Next we describe the effect of H_0 on the irreducible objects of \mathcal{O} .

Theorem 3.17. *Let $\mathbf{A} \in \text{Tab}$. Then*

$$H_0(L(\mathbf{A})) \cong \begin{cases} \overline{L}(\mathbf{A}) & \text{if } \mathbf{A} \text{ is anti-dominant,} \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We first show that $H_0(L(\mathbf{A})) = 0$ if \mathbf{A} is not anti-dominant. Let \mathfrak{p}' be the parabolic subalgebra of \mathfrak{g} spanned by $\{e_{i,j}\}_{\text{row}(i) \leq \text{row}(j)}$, i.e. $\mathfrak{p}' = \mathfrak{g}_{\overline{0}} + \mathfrak{b}$. Let $M_{ev}(\mathbf{A}) := U(\mathfrak{g}_{\overline{0}}) \otimes_{U(\mathfrak{b}_{\overline{0}})} \mathbb{C}_{\lambda_{\mathbf{A}}}$ be the Verma module for $\mathfrak{g}_{\overline{0}}$ of $\mathfrak{b}_{\overline{0}}$ -highest weight $\lambda_{\mathbf{A}}$. We view it also as a \mathfrak{p}' -supermodule concentrated in parity $\text{par}(\lambda_{\mathbf{A}})$ via the natural projection $\mathfrak{p}' \rightarrow \mathfrak{g}_{\overline{0}}$. Then we have obviously that

$$M(\mathbf{A}) \cong U(\mathfrak{g}) \otimes_{U(\mathfrak{p}')} M_{ev}(\mathbf{A}).$$

Assume that \mathbf{A} is not anti-dominant, and let \mathbf{B} be the unique anti-dominant tableau such that $\mathbf{A} \sim \mathbf{B} \approx \mathbf{A}$. By classical theory, the Verma module $M_{ev}(\mathbf{B})$ embeds into $M_{ev}(\mathbf{A})$. Hence, applying $U(\mathfrak{g}) \otimes_{U(\mathfrak{p}')} -$, we see that $M(\mathbf{B})$ embeds into $M(\mathbf{A})$ too. Now apply the exact functor H_0 to the resulting short exact sequence $0 \rightarrow M(\mathbf{B}) \rightarrow M(\mathbf{A}) \rightarrow C \rightarrow 0$ using Theorem 3.14, to obtain an exact sequence $0 \rightarrow \overline{M}(\mathbf{B}) \rightarrow \overline{M}(\mathbf{A}) \rightarrow H_0(C) \rightarrow 0$. But $\overline{M}(\mathbf{B}) \cong \overline{M}(\mathbf{A})$ as $\mathbf{B} \sim \mathbf{A}$, hence, we must have that $H_0(C) = 0$. Since $C \twoheadrightarrow L(\mathbf{A})$, this implies that $H_0(L(\mathbf{A})) = 0$.

We next show for anti-dominant \mathbf{A} that

$$[H_0(L(\mathbf{A}))] = [\overline{L}(\mathbf{A})] + (\text{a sum of } [\overline{L}(\mathbf{B})] \text{ for } \mathbf{B} \in \text{Tab with } a(\mathbf{B}) < a(\mathbf{A})). \quad (3.24)$$

To see this, note since $L(\mathbf{A})$ is a quotient of $M(\mathbf{A})$ that $H_0(L(\mathbf{A}))$ is a quotient of $H_0(M(\mathbf{A})) \cong \overline{M}(\mathbf{A})$. Applying Corollary 3.16, we deduce either that (3.24) holds or that $H_0(L(\mathbf{A})) = 0$. Also by Lemma 2.6 we know (as \mathbf{A} is anti-dominant) that

$$[M(\mathbf{A})] = [L(\mathbf{A})] + (\text{a sum of } [L(\mathbf{B})] \text{ for } \mathbf{B} \in \text{Tab with } a(\mathbf{B}) < a(\mathbf{A})).$$

Applying H_0 and using (3.24) whenever $H_0(L(\mathbf{B})) \neq 0$, we deduce that

$$[\overline{M}(\mathbf{A})] = [H_0(L(\mathbf{A}))] + (\text{a sum of } [\overline{L}(\mathbf{B})] \text{ for } \mathbf{B} \in \text{Tab with } a(\mathbf{B}) < a(\mathbf{A})).$$

Since this definitely involves $[\overline{L}(\mathbf{A})]$, we must have that $H_0(L(\mathbf{A})) \neq 0$, and we have established (3.24).

Now we claim for any $\mathbf{A} \in \text{Tab}$ that there exists some anti-dominant $\mathbf{B} \sim \mathbf{A}$ such that $H_0(L(\mathbf{B})) \cong \overline{L}(\mathbf{A})$. To see this, we know by Corollary 3.13 that there exists some $M \in \mathcal{O}$ with $H_0(M) \cong \overline{L}(\mathbf{A})$. Say we have that $[M] = [L(\mathbf{B}_1)] + \cdots + [L(\mathbf{B}_k)]$ in the Grothendieck group for some $\mathbf{B}_1, \dots, \mathbf{B}_k \in \text{Tab}$. In view of (3.24), only one of $\mathbf{B}_1, \dots, \mathbf{B}_k$ can be anti-dominant, and this \mathbf{B}_i must satisfy $H_0(L(\mathbf{B}_i)) \cong \overline{L}(\mathbf{B}_i) \cong \overline{L}(\mathbf{A})$. This proves the claim.

We have now shown in any row equivalence class of π -tableaux that there exists at least one anti-dominant \mathbf{A} with $H_0(L(\mathbf{A})) \cong \overline{L}(\mathbf{A})$. Suppose that \mathbf{B} is a different anti-dominant tableau in the same row equivalence class as \mathbf{A} . We need to show that $H_0(L(\mathbf{B})) \cong \overline{L}(\mathbf{B})$ too. To prove this we may assume that $\mathbf{B} = \sigma(\mathbf{A})$ for some simple transposition $\sigma \in S_m \times S_n$. Let ξ be the linkage class

containing A , so that $\sigma(\xi)$ is the linkage class containing B . By Lemma 2.9, there is an equivalence $T_\sigma : \mathcal{O}_\xi \rightarrow \mathcal{O}_{\sigma(\xi)}$ such that $T_\sigma(L(A)) \cong L(B)$ and $T_\sigma(M(C)) \cong M(\sigma(C))$ for each $C \approx A$. The \mathbb{Z} -linear maps $[H_0] : K_0(\mathcal{O}_\xi) \rightarrow K_0(\overline{\mathcal{O}})$ and $[H_0 \circ T_\sigma] : K_0(\mathcal{O}_\xi) \rightarrow K_0(\overline{\mathcal{O}})$ are equal; this follows because they are equal on $[M(C)]$ for each $C \approx A$ as $\overline{M}(C) \cong \overline{M}(\sigma(C))$. Hence, we get that

$$[H_0]([L(B)]) = [H_0 \circ T_\sigma]([L(A)]) = [H_0]([L(A)]) = [\overline{L}(A)] = [\overline{L}(B)].$$

This implies that $H_0(L(B)) \cong \overline{L}(B)$ as required. \square

Corollary 3.18. *The full subcategory of \mathcal{O} consisting of all objects annihilated by H_0 consists of all the supermodules in \mathcal{O} of strictly less than maximal Gelfand–Kirillov dimension.*

Proof. This follows from Theorem 3.17 on recalling that A is anti-dominant if and only if $L(A)$ is of maximal Gelfand–Kirillov dimension. \square

3.8. The center of W . In this subsection we determine the center of W . The argument here is similar in spirit to the proof of an analogous result in the purely even setting from [BK1, §6.4]; it depends crucially on Corollary 3.15. Let $\text{pr} : U(\mathfrak{g}) \rightarrow U(\mathfrak{p})$ be the projection along the direct sum decomposition $U(\mathfrak{g}) = U(\mathfrak{p}) \oplus \mathfrak{m}_\chi U(\mathfrak{g})$. It is easy to see from (3.2) that the restriction of pr defines an algebra homomorphism

$$\text{pr} : Z(\mathfrak{g}) \rightarrow Z(W). \quad (3.25)$$

The goal is to show that this map is actually an *isomorphism*.

Consider the Harish-Chandra homomorphism $\text{HC} : Z(\mathfrak{g}) \xrightarrow{\sim} I(\mathfrak{t})$ from Theorem 2.2. Recalling Theorem 2.3, we adopt the definition of HC that is adapted to the Borel subalgebra \mathfrak{b}' , i.e. we view HC as the restriction of the map

$$S_{-\rho'} \circ \phi' : U(\mathfrak{g})_0 \rightarrow S(\mathfrak{t}), \quad (3.26)$$

where $\phi' : U(\mathfrak{g})_0 \rightarrow S(\mathfrak{t})$ is projection along $U(\mathfrak{g})_0 = S(\mathfrak{t}) \oplus (U(\mathfrak{g})_0 \cap U(\mathfrak{g})\mathfrak{n}')$ and \mathfrak{n}' is the nilradical of \mathfrak{b}' . The restriction of (3.26) to $Z(\mathfrak{h})$ also gives a conveniently normalized Harish-Chandra homomorphism for the Lie superalgebra \mathfrak{h} , that is, an isomorphism $\text{hc} : Z(\mathfrak{h}) \xrightarrow{\sim} J(\mathfrak{t})$ where

$$J(\mathfrak{t}) := \left\{ f \in S(\mathfrak{t}) \mid \begin{array}{l} \frac{\partial f}{\partial x_i} + \frac{\partial f}{\partial y_j} \equiv 0 \pmod{x_i - y_j} \\ \text{for } 1 \leq i \leq m \text{ and } j = i + s_- \end{array} \right\}. \quad (3.27)$$

Also let $\pi : U(\mathfrak{p}) \rightarrow U(\mathfrak{h})$ be the usual projection, so that $\ker \pi = U(\mathfrak{p})\mathfrak{r}$ where \mathfrak{r} is the nilradical of \mathfrak{p} . We have now set up all of the notation to make sense of the following diagram:

$$\begin{array}{ccc} U(\mathfrak{g})_0 & \xrightarrow{S_{-\rho'} \circ \phi'} & S(\mathfrak{t}) \\ \text{pr} \downarrow & & \uparrow S_{-\rho'} \circ \phi' \\ U(\mathfrak{p})_0 & \xrightarrow{\pi} & U(\mathfrak{h})_0 \end{array} \quad (3.28)$$

Moreover, this diagram commutes. The final important point is that the restriction of π to W is injective; this is equivalent to the injectivity of the Miura transform in [BBG, Theorem 4.5].

Lemma 3.19. *The images of $d_1^{(r)}$ and $\tilde{d}_2^{(r)}$ under the map $S_{-\rho'} \circ \phi' \circ \pi$ are equal to $e_r(x_1, \dots, x_m)$ and $(-1)^r h_r(y_1, \dots, y_n)$, respectively.*

Proof. This depends on the explicit formulae for these elements of W from [BBG, Section 4]. For example, for $\tilde{d}_2^{(r)}$ remembering the typo pointed out in the footnote on p. 11, we have that

$$\pi(\tilde{d}_2^{(r)}) = S_{\rho'} \left(\sum (-1)^{r+|i_1|+\dots+|i_r|} e_{i_1, j_1} \cdots e_{i_r, j_r} \right),$$

summing over all $1 \leq i_1, \dots, i_r, j_1, \dots, j_r \leq m+n$ such that

- $\text{row}(i_1) = \text{row}(j_r) = 2$;
- $\text{col}(i_s) = \text{col}(j_s)$ for each s ;
- $\text{row}(i_{s+1}) = \text{row}(j_s)$ and $\text{col}(i_{s+1}) \leq \text{col}(j_s)$ for $s = 1, \dots, r-1$.

To apply ϕ' to this, note for one of the monomials $e_{i_1, j_1} \cdots e_{i_r, j_r}$ that ϕ' gives zero if $\text{row}(i_r) = 1$, hence, we may assume that $i_r = j_r$; then we get zero if $\text{row}(i_{r-1}) = 1$, hence, $i_{r-1} = j_{r-1}$; and so on. We deduce $S_{-\rho'}(\phi'(\pi(\tilde{d}_2^{(r)}))) = (-1)^r h_r(y_1, \dots, y_n)$ as claimed. \square

Lemma 3.20. *We have that $\pi(Z(W)) \subseteq Z(\mathfrak{h})$.*

Proof. We must show for $z \in Z(W)$ and $u \in U(\mathfrak{h})$ that $[\pi(z), u] = 0$. If $\mathbf{A} \in \text{Tab}$ is any typical tableau, i.e. $\text{atyp}(\mathbf{A}) = 0$, then $K(\mathbf{A})$ is an irreducible \mathfrak{h} -supermodule which remains irreducible (with one-dimensional endomorphism algebra) on restriction to W , as follows from Corollaries 3.15 and 3.16. Hence, $\pi(z)$ acts as a scalar on $K(\mathbf{A})$, implying that $[\pi(z), u] \in \text{Ann}_{U(\mathfrak{h})} K(\mathbf{A})$. It remains to observe that

$$\bigcap_{\substack{\mathbf{A} \in \text{Tab} \\ \text{atyp}(\mathbf{A})=0}} \text{Ann}_{U(\mathfrak{h})} K(\mathbf{A}) = 0.$$

This follows because $\{\mathbf{A} \in \text{Tab} \mid \text{atyp}(\mathbf{A}) = 0\}$ is Zariski dense in Tab (identified with \mathbb{A}^{m+n} in the obvious way). Now we can apply the standard fact that the annihilator of any dense set of Verma supermodules is zero, see for example the proof of [M, Lemma 13.1.4]³. \square

Theorem 3.21. *The homomorphism $\text{pr} : Z(\mathfrak{g}) \rightarrow Z(W)$ from (3.25) is an algebra isomorphism. Moreover, we have that $\text{pr}(z_r) = \tilde{c}^{(r)}$, where $z_r \in Z(\mathfrak{g})$ and $\tilde{c}^{(r)} \in Z(W)$ are defined by (2.8) and (3.5), respectively.*

Proof. We observe to start with that $Z(W) \subseteq U(\mathfrak{h})_0 \oplus U(\mathfrak{p})\mathfrak{r}$. To see this, note that $U(\mathfrak{p}) = U(\mathfrak{h}) \oplus U(\mathfrak{p})\mathfrak{r}$. Hence, we can write $z \in Z(W)$ as $z_0 + z_1$ with $z_0 \in U(\mathfrak{h})$ and $z_1 \in U(\mathfrak{p})\mathfrak{r}$. Applying π and using Lemma 3.20, we get that $z_0 = \pi(z) \in Z(\mathfrak{h}) \subseteq U(\mathfrak{h})_0$, as required. Hence, it makes sense to restrict all the maps in the commutative diagram (3.28) to obtain another commutative

³It is easy to supply a direct proof of this statement in the present situation since \mathfrak{h} is a direct sum of copies of $\mathfrak{gl}_1(\mathbb{C})$ and $\mathfrak{gl}_{1|1}(\mathbb{C})$.

diagram

$$\begin{array}{ccc}
 Z(\mathfrak{g}) & \xrightarrow{\text{HC}} & S(\mathfrak{t}) \\
 \text{pr} \downarrow & & \uparrow \text{hc} \\
 Z(W) & \xrightarrow{\pi} & Z(\mathfrak{h})
 \end{array} \tag{3.29}$$

Since HC is injective, so too is the map pr. Since $\tilde{c}^{(r)} = \sum_{s+t=r} d_1^{(s)} \tilde{d}_2^{(t)}$, we get from Lemma 3.19 and Theorem 2.4 that

$$\text{hc}(\pi(\tilde{c}^{(r)})) = e_r(x_1, \dots, x_m/y_1, \dots, y_n) = \text{HC}(z_r).$$

Hence, $\text{pr}(z_r) = \tilde{c}^{(r)}$.

To complete the proof of the theorem, we must show that pr is surjective. As $\text{hc} \circ \pi$ is injective and $\text{HC}(Z(\mathfrak{g})) = I(\mathfrak{t})$, this follows if we can show that $\text{hc}(\pi(Z(W))) \subseteq I(\mathfrak{t})$. Since $\overline{M}(A) = \overline{M}(B)$ for $A \sim B$, Corollary 3.15 implies that the generalized central character of the W -supermodule $\overline{K}(A)$ depends only on the row equivalence class of A . Hence, for $z \in Z(W)$ we deduce that $\pi(z)$ acts by the same scalar on the \mathfrak{h} -supermodules $K(A)$ for all A in the same row equivalence class. In other words, $\text{hc}(\pi(Z(W))) \subseteq S(\mathfrak{t})^{S_m \times S_n}$. We also have that $\text{hc}(\pi(Z(W))) \subseteq \text{hc}(Z(\mathfrak{h})) = J(\mathfrak{t})$. It remains to observe by the definitions (2.5) and (3.27) that $I(\mathfrak{t}) = S(\mathfrak{t})^{S_m \times S_n} \cap J(\mathfrak{t})$. \square

Corollary 3.22. *The center $Z(W)$ is generated by the elements $\{\tilde{c}^{(r)}\}_{r \geq 1}$; equivalently, it is generated by the elements $\{c^{(r)}\}_{r \geq 1}$.*

Proof. This follows from Theorems 3.21 and 2.4. \square

4. THE QUOTIENT CATEGORY $\overline{\mathcal{O}}_{\mathbb{Z}}$

For the remainder of the article, we restrict attention to integral central characters, denoting the corresponding subcategory of \mathcal{O} by $\mathcal{O}_{\mathbb{Z}}$. We introduce an Abelian subcategory $\overline{\mathcal{O}}_{\mathbb{Z}}$ of $W\text{-smod}_{\text{fd}}$ such that the Whittaker coinvariants functor restricts to a quotient functor

$$H_0 : \mathcal{O}_{\mathbb{Z}} \rightarrow \overline{\mathcal{O}}_{\mathbb{Z}}.$$

We show that this functor satisfies the double centralizer property, i.e. it is fully faithful on projectives. Then we discuss the locally unital (“idempotent”) algebras that are Morita equivalent to the blocks of $\overline{\mathcal{O}}_{\mathbb{Z}}$, and give some applications to the classification of blocks of $\mathcal{O}_{\mathbb{Z}}$.

4.1. Categorical actions. Let $\text{Tab}_{\mathbb{Z}}$ be the subset of Tab consisting of the tableaux all of whose entries are integers. Let $\mathcal{O}_{\mathbb{Z}}$ be the Serre subcategory of \mathcal{O} generated by the \mathfrak{g} -supermodules $\{L(A)\}_{A \in \text{Tab}_{\mathbb{Z}}}$. It is a sum of blocks of \mathcal{O} :

$$\mathcal{O}_{\mathbb{Z}} = \bigoplus_{\xi \in \text{Tab}_{\mathbb{Z}}/\approx} \mathcal{O}_{\xi}. \tag{4.1}$$

In particular, $\mathcal{O}_{\mathbb{Z}}$ is itself a highest weight category with weight poset $(\text{Tab}_{\mathbb{Z}}, \leq)$.

Adopting some standard Lie theoretic notation, let \mathfrak{sl}_{∞} be the Kac-Moody algebra of type A_{∞} (over \mathbb{C}), with Chevalley generators $\{E_i, F_i\}_{i \in \mathbb{Z}}$, weight lattice $P := \bigoplus_{i \in \mathbb{Z}} \mathbb{Z}\varepsilon_i$, simple roots $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$, etc.. We denote its natural module

by V^+ and the dual by V^- . These have standard bases $\{v_j^+\}_{j \in \mathbb{Z}}$ and $\{v_j^-\}_{j \in \mathbb{Z}}$, respectively. The vector v_j^\pm is of weight $\pm \varepsilon_j$, and the Chevalley generators act by

$$F_i v_j^+ = \begin{cases} v_{j+1}^+ & \text{if } j = i \\ 0 & \text{otherwise,} \end{cases} \quad E_i v_j^+ = \begin{cases} v_{j-1}^+ & \text{if } j = i + 1 \\ 0 & \text{otherwise,} \end{cases} \quad (4.2)$$

$$F_i v_j^- = \begin{cases} v_{j-1}^- & \text{if } j = i + 1 \\ 0 & \text{otherwise,} \end{cases} \quad E_i v_j^- = \begin{cases} v_{j+1}^- & \text{if } j = i \\ 0 & \text{otherwise.} \end{cases} \quad (4.3)$$

As goes back to [B1] (or [CR, §7.4] in the purely even case), there is a categorical action of \mathfrak{sl}_∞ on \mathcal{O} in the sense of Rouquier [R1, Definition 5.32]; see also [BLW, Definition 2.6] for our precise conventions. We just give a brief summary of the construction, referring to the proof of [BLW, Theorem 3.10] for details.

- The required biadjoint endofunctors F and E are the functors

$$F := U \otimes -, \quad E := U^\vee \otimes -, \quad (4.4)$$

where U is the natural \mathfrak{g} -supermodule of column vectors and U^\vee is its dual.

- The natural transformations $x : F \Rightarrow F$ and $s : F^2 \Rightarrow F^2$ are defined so that $x_M : U \otimes M \rightarrow U \otimes M$ is left multiplication by the Casimir tensor

$$\Omega := \sum_{i,j=1}^{m+n} (-1)^{|j|} e_{i,j} \otimes e_{j,i} \in \mathfrak{g} \otimes \mathfrak{g}, \quad (4.5)$$

and $s_M : U \otimes U \otimes M \rightarrow U \otimes U \otimes M$ is induced by the tensor flip $U \otimes U \rightarrow U \otimes U, u \otimes v \mapsto (-1)^{|u||v|} v \otimes u$.

- Let F_i be the summand of F defined by taking the generalized i -eigenspace of x , and E_i be the unique summand of E that is biadjoint to it. Let $\mathcal{O}_{\mathbb{Z}}^\Delta$ be the exact subcategory of $\mathcal{O}_{\mathbb{Z}}$ consisting of all supermodules admitting a Verma flag, and $K_0(\mathcal{O}_{\mathbb{Z}}^\Delta)_{\mathbb{C}}$ be its complexified Grothendieck group. Let $T^{m|n} := (V^+)^{\otimes m} \otimes (V^-)^{\otimes n}$, and set

$$v_{\mathbf{A}} := v_{a_1}^+ \otimes \cdots \otimes v_{a_m}^+ \otimes v_{b_1}^- \otimes \cdots \otimes v_{b_n}^- \in T^{m|n} \quad (4.6)$$

for each $\mathbf{A} = \begin{smallmatrix} a_1 & \cdots & a_m \\ b_1 & \cdots & b_n \end{smallmatrix} \in \text{Tab}_{\mathbb{Z}}$. Then, there is a vector space isomorphism

$$K_0(\mathcal{O}_{\mathbb{Z}}^\Delta)_{\mathbb{C}} \xrightarrow{\sim} T^{m|n}, \quad [M(\mathbf{A})] \mapsto v_{\mathbf{A}}. \quad (4.7)$$

Moreover, this map intertwines the operators induced by the endofunctors F_i and E_i on the left hand space with the actions of the Chevalley generators of \mathfrak{sl}_∞ on the right.

- Under the isomorphism from (4.7), the Grothendieck groups $K_0(\mathcal{O}_{\xi}^\Delta)_{\mathbb{C}}$ of the blocks correspond to the weight spaces of $T^{m|n}$.

In fact, $\mathcal{O}_{\mathbb{Z}}$ is a *tensor product categorification* of $T^{m|n}$ in the general sense of [BLW, Definition 2.10].

In the rest of the subsection, we are going to formulate an analogous categorification theorem at the level of W . Observe that a π -tableau $\mathbf{A} = \begin{smallmatrix} a_1 & \cdots & a_m \\ b_1 & \cdots & b_n \end{smallmatrix} \in \text{Tab}_{\mathbb{Z}}$ is anti-dominant if and only if $a_1 \leq \cdots \leq a_m$ and $b_1 \geq \cdots \geq b_n$. Let $\text{Tab}_{\mathbb{Z}}^\circ$ denote the set of all such tableaux. It gives a distinguished set of representatives

for $\text{Tab}_{\mathbb{Z}}/\sim$. For a linkage class $\xi \in \text{Tab}_{\mathbb{Z}}/\approx$, we let ξ° denote the set $\xi \cap \text{Tab}_{\mathbb{Z}}^\circ$ of anti-dominant tableaux that it contains.

Recall for $A \in \text{Tab}_{\mathbb{Z}}$ that $P(A)$ is the projective cover of $L(A)$ in $\mathcal{O}_{\mathbb{Z}}$. Let

$$\overline{P}(A) := H_0(P(A)). \quad (4.8)$$

Then we define $\overline{\mathcal{O}}_{\mathbb{Z}}$ to be the full subcategory of $W\text{-smod}_{\text{fd}}$ consisting of all W -supermodules that are isomorphic to subquotients of finite direct sums of the supermodules $\{\overline{P}(A)\}_{A \in \text{Tab}_{\mathbb{Z}}^\circ}$. This is obviously an Abelian subcategory of $W\text{-smod}_{\text{fd}}$. Similarly, given a linkage class $\xi \in \text{Tab}_{\mathbb{Z}}/\approx$, we let $\overline{\mathcal{O}}_{\xi}$ be the full subcategory consisting of subquotients of finite direct sums of the supermodules $\{\overline{P}(A)\}_{A \in \xi^\circ}$.

Lemma 4.1. *The Whittaker coinvariants functor restricts to an exact functor $H_0 : \mathcal{O}_{\mathbb{Z}} \rightarrow \overline{\mathcal{O}}_{\mathbb{Z}}$ sending each block \mathcal{O}_{ξ} to $\overline{\mathcal{O}}_{\xi}$. Each $\overline{\mathcal{O}}_{\xi}$ is itself a block (i.e. it is indecomposable), and $\overline{\mathcal{O}}_{\mathbb{Z}}$ decomposes as*

$$\overline{\mathcal{O}}_{\mathbb{Z}} = \bigoplus_{\xi \in \text{Tab}_{\mathbb{Z}}/\approx} \overline{\mathcal{O}}_{\xi}.$$

Moreover, the supermodules $\{\overline{L}(A)\}_{A \in \xi^\circ}$ give a complete set of inequivalent irreducible objects in each $\overline{\mathcal{O}}_{\xi}$.

Proof. We first show that the essential image of H_0 is contained in $\overline{\mathcal{O}}_{\mathbb{Z}}$. Any object $M \in \mathcal{O}_{\mathbb{Z}}$ is a quotient of a direct sum of the projective objects $P(A)$ for $A \in \text{Tab}_{\mathbb{Z}}$. Since H_0 is exact, we deduce that $H_0(M)$ is a quotient of a direct sum of the objects $\overline{P}(A)$ for $A \in \text{Tab}_{\mathbb{Z}}$. Since, by definition, $\overline{\mathcal{O}}_{\mathbb{Z}}$ is closed under taking quotients and direct sums, we are thus reduced to showing that each $\overline{P}(A)$ for $A \in \text{Tab}_{\mathbb{Z}}$ belongs to $\overline{\mathcal{O}}_{\mathbb{Z}}$. This is immediate by the definition of $\overline{\mathcal{O}}_{\mathbb{Z}}$ if A is anti-dominant. So suppose that A is not anti-dominant. Then $P(A)$ has a Verma flag, and the socle of any Verma is anti-dominant, hence, the injective hull of $P(A)$ is a direct sum of $P(B)$ for $B \in \text{Tab}_{\mathbb{Z}}^\circ$; see [BLW, Theorem 2.24]. Applying H_0 we deduce that $\overline{P}(A)$ embeds into a direct sum of $\overline{P}(B)$ for $B \in \text{Tab}_{\mathbb{Z}}^\circ$. Since $\overline{\mathcal{O}}_{\mathbb{Z}}$ is closed under taking submodules, this implies that $\overline{P}(A)$ belongs to $\overline{\mathcal{O}}_{\mathbb{Z}}$.

Thus, H_0 restricts to a well-defined exact functor $\mathcal{O}_{\mathbb{Z}} \rightarrow \overline{\mathcal{O}}_{\mathbb{Z}}$. The same argument at the level of blocks shows that H_0 maps \mathcal{O}_{ξ} to $\overline{\mathcal{O}}_{\xi}$, and clearly $\overline{\mathcal{O}}_{\mathbb{Z}}$ decomposes as the direct sum of the $\overline{\mathcal{O}}_{\xi}$'s. The irreducible objects in $\overline{\mathcal{O}}_{\xi}$ are just the irreducible objects of $W\text{-smod}$ that it contains, so they are represented by $\{\overline{L}(A) \mid A \in \xi^\circ\}$ thanks to Theorem 3.17.

It remains to show that each $\overline{\mathcal{O}}_{\xi}$ is indecomposable. Corollary 3.16 implies for any tableaux A, B with $B \uparrow A$ that the irreducible supermodules $\overline{L}(A)$ and $\overline{L}(B)$ are both composition factors of the indecomposable object $\overline{M}(A)$. Hence, $\overline{L}(A)$ and $\overline{L}(B)$ belong to the same block of $\overline{\mathcal{O}}$. Now observe that the equivalence relation \approx on $\text{Tab}_{\mathbb{Z}}$ is generated by the relations \sim and \uparrow . \square

Remark 4.2. By Lemma 3.8 and the definition of $\overline{\mathcal{O}}_{\mathbb{Z}}$, the elements $d_1^{(1)}$ and $d_2^{(1)}$ act semisimply on any object $M \in \overline{\mathcal{O}}_{\mathbb{Z}}$. Lemma 3.8 shows moreover that the z -eigenspace of $d_2^{(1)}$ is concentrated in parity $\text{par}(z)$, i.e. the $\mathbb{Z}/2$ -grading is determined by the eigenspace decomposition of $d_2^{(1)}$. This is a similar situation

to category \mathcal{O} itself, where the $\mathbb{Z}/2$ -grading was determined by the weight space decomposition.

Next we introduce endofunctors \bar{F} and \bar{E} of $\bar{\mathcal{O}}_{\mathbb{Z}}$. Consider the biadjoint endofunctors $\bar{F} := U \circledast -$ and $\bar{E} := U^\vee \circledast -$ of $W\text{-smod}_{\text{fd}}$ from §3.4 (where U is still the natural \mathfrak{g} -supermodule). By Theorem 3.10, we have canonical isomorphisms of functors

$$\bar{F} \circ H_0 \xrightarrow{\cong} H_0 \circ F, \quad \bar{E} \circ H_0 \xrightarrow{\cong} H_0 \circ E, \quad (4.9)$$

going from \mathcal{O} to $W\text{-smod}_{\text{fd}}$. It follows immediately that $\bar{F}(\bar{P}(\mathbf{A})) \cong H_0(FP(\mathbf{A}))$, hence, it is in the subcategory $\bar{\mathcal{O}}_{\mathbb{Z}}$. Since \bar{F} is exact, it follows that \bar{F} leaves the subcategory $\bar{\mathcal{O}}_{\mathbb{Z}}$ of $W\text{-smod}_{\text{fd}}$ invariant. Similarly, so does \bar{E} . Hence, we can restrict these endofunctors to obtain a biadjoint pair of endofunctors

$$\bar{F} : \bar{\mathcal{O}}_{\mathbb{Z}} \rightarrow \bar{\mathcal{O}}_{\mathbb{Z}}, \quad \bar{E} : \bar{\mathcal{O}}_{\mathbb{Z}} \rightarrow \bar{\mathcal{O}}_{\mathbb{Z}}. \quad (4.10)$$

Theorem 4.3. *There are natural transformations $\bar{x} : \bar{F} \Rightarrow \bar{F}$ and $\bar{s} : \bar{F}^2 \Rightarrow \bar{F}^2$ making $\bar{\mathcal{O}}_{\mathbb{Z}}$ into an integrable \mathfrak{sl}_{∞} -categorification. Moreover, the Whittaker coinvariants functor $H_0 : \mathcal{O}_{\mathbb{Z}} \rightarrow \bar{\mathcal{O}}_{\mathbb{Z}}$ is strongly equivariant in the usual sense of categorical actions (e.g. see [BLW, Definition 2.7]).*

Proof. First, we go through the construction of \bar{x} . For a left $U(\mathfrak{g})$ -supermodule M , we already have $x_M : U \otimes M \rightarrow U \otimes M$ defined by left multiplication by the tensor Ω from (4.5). Under the isomorphism (3.18), the dual map $(x_M)^* : (U \otimes M)^* \rightarrow (U \otimes M)^*$ is the map $x_{M^*} : M^* \otimes U^* \rightarrow M^* \otimes U^*$ defined by right multiplication by Ω . Now suppose that $M \in W\text{-smod}_{\text{fd}}$. Applying H^0 to $x_{M^* \otimes_W Q}$ gives us an endomorphism \bar{x}_{M^*} of $M^* \circledast U^* = H^0((M^* \otimes_W Q) \otimes U^*)$. Finally, taking the left dual gives us an endomorphism $\bar{x}_M := {}^*(\bar{x}_{M^*})$ of $U \circledast M$.

The definition of \bar{s} can be obtained in a very similar way, but it is easier to define this using the coherence isomorphism $U \circledast (U \circledast M) \cong (U \otimes U) \circledast M$ coming from the monoidal functor (3.21), starting from the endomorphism of $(U \otimes U) \circledast M$ obtained by applying $- \circledast \text{id}_M$ to the tensor flip $U \otimes U \mapsto U \otimes U$.

The fact that \bar{x} and \bar{s} satisfy the appropriate degenerate affine Hecke algebra relations is just a formal consequence of the fact that x and s do on $U(\mathfrak{g})\text{-smod}$. Also, we've already constructed \bar{F} and \bar{E} so that they are canonically biadjoint.

Next we show that H_0 is a strongly equivariant functor. We have already constructed the required data of an isomorphism $\zeta : \bar{F} \circ H_0 \xrightarrow{\cong} H_0 \circ F$ on the left hand side of (4.9). We next have to check that x and s are intertwined with \bar{x} and \bar{s} in the appropriate sense (we need the F -version of [CR, 5.2.1(5)] as recorded in [BLW, Definition 2.7(E2)–(E3)]). This is a formal exercise from the definitions (which were set up exactly for this purpose). Finally, we must check the F -version of [CR, 5.1.2(4)] (which is [BLW, Definition 2.7(E1)]). This asserts that a certain natural transformations $H_0 \circ E \Rightarrow \bar{E} \circ H_0$ constructed from ζ using the adjunction is an isomorphism. In fact, one shows that it is the inverse of the right hand side of (4.9). We omit the details here.

Then we decompose \bar{F} into its \bar{x} -generalized eigenspaces \bar{F}_i as before, and let \bar{E}_i be the adjoint summands of \bar{E} . Finally, we need to show that the induced actions of $[\bar{F}_i]$ and $[\bar{E}_i]$ make $K_0(\bar{\mathcal{O}}_{\mathbb{Z}})_{\mathbb{C}}$ into an integrable representation of \mathfrak{sl}_{∞} . This follows from the equivariance of H_0 : we already know that $K_0(\mathcal{O}_{\mathbb{Z}})_{\mathbb{C}}$ is integrable upstairs, and the \mathfrak{sl}_{∞} -equivariant map $[H_0] : K_0(\mathcal{O}_{\mathbb{Z}})_{\mathbb{C}} \rightarrow K_0(\bar{\mathcal{O}}_{\mathbb{Z}})_{\mathbb{C}}$ is

surjective according to Theorem 3.17 and the description of irreducible objects in Lemma 4.1. \square

The Grothendieck group $K_0(\overline{\mathcal{O}}_{\mathbb{Z}})_{\mathbb{C}}$ may be understood from the point of view of this categorification theorem as follows.

Lemma 4.4. *Let $S^{m|n} := S^m V^+ \otimes S^n V^-$ (tensor product of symmetric powers). Then, there is a unique injective linear map j making the following into a commutative diagram of \mathfrak{sl}_{∞} -module homomorphisms:*

$$\begin{array}{ccc} T^{m|n} & \xrightarrow{\text{can}} & S^{m|n} \\ \downarrow i & & \downarrow j \\ K_0(\mathcal{O}_{\mathbb{Z}})_{\mathbb{C}} & \xrightarrow{[H_0]} & K_0(\overline{\mathcal{O}}_{\mathbb{Z}})_{\mathbb{C}} \end{array}$$

Here, the top map is the canonical map from tensor powers to symmetric powers, and i is the composition of the inverse of (4.7) with the natural inclusion $K_0(\mathcal{O}_{\mathbb{Z}}^{\Delta})_{\mathbb{C}} \hookrightarrow K_0(\mathcal{O}_{\mathbb{Z}})_{\mathbb{C}}$.

Proof. To see this, one just has to observe that $H_0(M(\mathbf{A})) \cong H_0(M(\mathbf{B}))$ for all $\mathbf{A} \sim \mathbf{B}$ thanks to Theorem 3.14. Moreover, the classes of the Verma supermodules $\{[\overline{M}(\mathbf{A})]\}_{\mathbf{A} \in \text{Tab}^{\circ}}$ are linearly independent in $K_0(\overline{\mathcal{O}}_{\mathbb{Z}})_{\mathbb{C}}$ by the classification of irreducible objects. \square

4.2. Serre quotients and the double centralizer property. Throughout the subsection, we often appeal to Theorem 3.17 and the exactness of H_0 from Lemma 3.6. Although it is immediate from the definition that $\overline{\mathcal{O}}_{\mathbb{Z}}$ is an Abelian category, we do not yet know that it has enough projectives or injectives. We proceed to establish this, essentially mimicking the proof of [BK2, Lemma 5.7].

Lemma 4.5. *For each $\mathbf{A} \in \text{Tab}_{\mathbb{Z}}^{\circ}$, the supermodule $\overline{P}(\mathbf{A})$ is both the projective cover and the injective hull of $\overline{L}(\mathbf{A})$ in $\overline{\mathcal{O}}_{\mathbb{Z}}$.*

Proof. We need the following fact established in [BLW, Theorem 2.24]: for any $\mathbf{A} \in \text{Tab}_{\mathbb{Z}}^{\circ}$, the prinjective supermodule $P(\mathbf{A})$ is a direct summand of $F^d P(\mathbf{B})$ for some $d \geq 0$ and some $\mathbf{B} \in \text{Tab}_{\mathbb{Z}}^{\circ}$ of the special form $\mathbf{B} = \overset{a}{b} \dots \overset{a}{b}$ with $a \neq b$. Define $d(\mathbf{A})$ to be the smallest d such that this is the case.

We'll prove the lemma by induction on $d(\mathbf{A})$. For the base case $d(\mathbf{A}) = 0$, we have that \mathbf{A} is the only π -tableau in its linkage class, so that $P(\mathbf{A}) = L(\mathbf{A})$. Hence, $\overline{P}(\mathbf{A}) = H_0(P(\mathbf{A})) = H_0(L(\mathbf{A})) = \overline{L}(\mathbf{A})$. We deduce immediately from its definition that $\overline{\mathcal{O}}_{\xi}$ is simple (i.e. equivalent to the category of finite-dimensional vector spaces). Now the conclusion is trivial in this case.

For the induction step, take $\mathbf{A} \in \text{Tab}_{\mathbb{Z}}^{\circ}$ with $d(\mathbf{A}) > 0$. The functors F and \overline{F} both have biadjoints, hence, they send prinjectives to prinjectives. Using Lemma 2.8 and the definition of $d(\mathbf{A})$, we can find some $\mathbf{C} \in \text{Tab}_{\mathbb{Z}}^{\circ}$ with $d(\mathbf{C}) = d(\mathbf{A}) - 1$ such that $P(\mathbf{A})$ is a summand of $FP(\mathbf{C})$. By induction, $\overline{P}(\mathbf{C})$ is both the projective cover and the injective hull of $\overline{L}(\mathbf{C})$. So we have that

$$FP(\mathbf{C}) \cong \bigoplus_{\mathbf{B} \in \text{Tab}_{\mathbb{Z}}^{\circ}} P(\mathbf{B})^{\oplus m_{\mathbf{B}}}, \quad \overline{F}\overline{P}(\mathbf{C}) \cong \bigoplus_{\mathbf{B} \in \text{Tab}_{\mathbb{Z}}^{\circ}} \overline{P}(\mathbf{B})^{\oplus m_{\mathbf{B}}},$$

for some multiplicities m_B with $m_A > 0$, and deduce that $\overline{P}(A)$ is prinjective in $\overline{\mathcal{O}}_{\mathbb{Z}}$.

Let $B \in \text{Tab}_{\mathbb{Z}}^{\circ}$. Since $L(B)$ appears in the head of $P(B)$, we see that $\overline{L}(B)$ appears in the head of $\overline{P}(B)$. So for $D \in \text{Tab}_{\mathbb{Z}}^{\circ}$, we have $\dim \text{Hom}_W(\overline{P}(B), \overline{L}(D)) \geq \delta_{B,D}$ and

$$\dim \text{Hom}_W(\overline{F}\overline{P}(C), \overline{L}(D)) \geq \sum_{B \in \text{Tab}_{\mathbb{Z}}^{\circ}} m_B \delta_{B,D} = m_D.$$

Moreover, the equality holds here if and only if $\dim \text{Hom}_W(\overline{P}(B), \overline{L}(D)) = \delta_{B,D}$ for all B with $m_B > 0$. This is indeed the case thanks to the following calculation:

$$\begin{aligned} \dim \text{Hom}_W(\overline{F}\overline{P}(C), \overline{L}(D)) &= \dim \text{Hom}_W(\overline{P}(C), \overline{E}\overline{L}(D)) \\ &= [\overline{E}\overline{L}(D) : \overline{L}(C)] \\ &= [EL(D) : L(C)] \\ &= \dim \text{Hom}_{\mathfrak{g}}(P(C), EL(D)) \\ &= \dim \text{Hom}_{\mathfrak{g}}(FP(C), L(D)) = m_D. \end{aligned}$$

The previous paragraph establishes that $\dim \text{Hom}_W(\overline{P}(A), \overline{L}(B)) = \delta_{A,B}$ for all B , so $\overline{P}(A)$ has irreducible head $\overline{L}(A)$. Thus we have shown that $\overline{P}(A)$ is the projective cover of $\overline{L}(A)$ in $\overline{\mathcal{O}}_{\mathbb{Z}}$, as required. A similar calculation shows that $\dim \text{Hom}_W(\overline{L}(B), \overline{P}(A)) = \delta_{A,B}$, and $\overline{P}(A)$ is the injective hull of $\overline{L}(A)$ too. \square

Lemma 4.6. *For any $A \in \text{Tab}^{\circ}$ and $M \in \mathcal{O}_{\mathbb{Z}}$, the functor H_0 induces an isomorphism*

$$\text{Hom}_{\mathfrak{g}}(P(A), M) \xrightarrow{\sim} \text{Hom}_W(\overline{P}(A), H_0(M)).$$

Proof. We are trying to show that the natural transformation $\text{Hom}_{\mathfrak{g}}(P(A), -) \Rightarrow \text{Hom}_W(\overline{P}(A), H_0(-))$ induced by the functor H_0 is an isomorphism. Since H_0 is exact, it suffices to check this gives an isomorphism as in the statement for M an irreducible supermodule in $\mathcal{O}_{\mathbb{Z}}$. If $M = L(B)$ for $B \in \text{Tab}_{\mathbb{Z}}$, then both sides are zero unless $B = A$, thanks to Theorem 3.17 and Lemma 4.5. If $B = A$ then, by Lemma 4.5, both sides are one-dimensional. The left hand side is spanned by an epimorphism $P(A) \twoheadrightarrow L(A)$, so remains non-zero when we apply H_0 . Hence, H_0 does indeed give an isomorphism. \square

Lemma 4.7. *The functor H_0 is essentially surjective.*

Proof. Let $M \in \overline{\mathcal{O}}_{\mathbb{Z}}$. Applying Lemma 4.5, we can construct a two-step projective resolution

$$\overline{P}_1 \xrightarrow{\overline{f}} \overline{P}_0 \rightarrow M \rightarrow 0$$

in $\overline{\mathcal{O}}_{\mathbb{Z}}$. This means that $M \cong \text{coker } \overline{f}$ for projectives $\overline{P}_1, \overline{P}_0 \in \overline{\mathcal{O}}_{\mathbb{Z}}$ and $\overline{f} \in \text{Hom}_W(\overline{P}_1, \overline{P}_0)$. Let $P_1, P_0 \in \mathcal{O}_{\mathbb{Z}}$ be prinjectives such that $H_0(P_1) \cong \overline{P}_1$ and $H_0(P_0) \cong \overline{P}_0$. By Lemma 4.6, the functor H_0 defines an isomorphism

$$\text{Hom}_{\mathfrak{g}}(P_1, P_0) \xrightarrow{\sim} \text{Hom}_W(\overline{P}_1, \overline{P}_0).$$

Hence, there exists $f \in \text{Hom}_{\mathfrak{g}}(P_1, P_0)$ so that $H_0(f)$ identifies with \overline{f} . Then, using exactness, we get that $H_0(\text{coker } f) \cong \text{coker } \overline{f} \cong M$. \square

Theorem 4.8. *The functor $H_0 : \mathcal{O}_{\mathbb{Z}} \rightarrow \overline{\mathcal{O}}_{\mathbb{Z}}$ satisfies the universal property of the Serre quotient of $\mathcal{O}_{\mathbb{Z}}$ by the Serre subcategory $\mathcal{T}_{\mathbb{Z}}$ consisting of all supermodules of less than maximal Gelfand-Kirillov dimension.*

Proof. Recalling Lemma 2.8, $\mathcal{T}_{\mathbb{Z}}$ is generated by $\{L(\mathbf{A})\}_{\mathbf{A} \in \text{Tab}_{\mathbb{Z}} \setminus \text{Tab}_{\mathbb{Z}}^{\circ}}$. By Theorem 3.17, the exact functor H_0 annihilates all of these objects. Hence, by the universal property of the Serre quotient functor $Q : \mathcal{O}_{\mathbb{Z}} \rightarrow \mathcal{O}_{\mathbb{Z}}/\mathcal{T}_{\mathbb{Z}}$, we get an induced functor $G : \mathcal{O}_{\mathbb{Z}}/\mathcal{T}_{\mathbb{Z}} \rightarrow \overline{\mathcal{O}}_{\mathbb{Z}}$ such that $H_0 = G \circ Q$. By Lemma 4.7, G is essentially surjective. It just remains to show that it is fully faithful, i.e. for all $M, N \in \mathcal{O}_{\mathbb{Z}}$ we have that $G : \text{Hom}_{\mathcal{O}_{\mathbb{Z}}/\mathcal{T}_{\mathbb{Z}}}(QM, QN) \xrightarrow{\sim} \text{Hom}_W(H_0(M), H_0(N))$. This is clear from Lemma 4.6 in case M is prinjective, since Q satisfies an analogous property by the general theory of quotient functors. Take any $M' := QM$ and $N' := QN$ and a two-step projective resolution $P'_1 \rightarrow P'_0 \rightarrow M' \rightarrow 0$ in $\mathcal{O}_{\mathbb{Z}}/\mathcal{T}_{\mathbb{Z}}$. We get a commuting diagram

$$\begin{array}{ccccc} 0 \rightarrow \text{Hom}_{\mathcal{O}_{\mathbb{Z}}/\mathcal{T}_{\mathbb{Z}}}(M', N') & \longrightarrow & \text{Hom}_{\mathcal{O}_{\mathbb{Z}}/\mathcal{T}_{\mathbb{Z}}}(P'_0, N') & \longrightarrow & \text{Hom}_{\mathcal{O}_{\mathbb{Z}}/\mathcal{T}_{\mathbb{Z}}}(P'_1, N') \\ & & \downarrow G & & \downarrow G \\ 0 \rightarrow \text{Hom}_W(GM', GN') & \longrightarrow & \text{Hom}_W(GP'_0, GN') & \longrightarrow & \text{Hom}_W(GP'_1, GN') \end{array}$$

with exact rows. We've already established that the last two vertical maps are isomorphisms, hence, so is the first one. \square

Corollary 4.9. *The functor $H_0 : \mathcal{O}_{\mathbb{Z}} \rightarrow \overline{\mathcal{O}}_{\mathbb{Z}}$ is fully faithful on projectives.*

Proof. Given the above theorem, this follows from [BLW, Theorem 4.10]. \square

We stress that, although $\overline{\mathcal{O}}_{\mathbb{Z}}$ is a quotient of a highest weight category, it is not highest weight itself (except in the trivial case $m + n = 1$).

4.3. Parametrization of blocks by core and atypicality. At this point, it is convenient to switch from using anti-dominant π -tableaux as our preferred index set for the irreducible objects of $\overline{\mathcal{O}}_{\mathbb{Z}}$ to some equivalent but more suggestive formalism. By a *composition* $\lambda \vDash n$, we mean an infinite tuple $\lambda = (\lambda_i)_{i \in \mathbb{Z}}$ of non-negative integers whose sum is n . The sum of two compositions is obtained simply by adding their corresponding parts. The *strictification* λ^+ of λ is the strict composition $(\lambda_1^+, \dots, \lambda_\ell^+)$ of n obtained from λ by discarding all of its parts that equal zero. The *transpose* λ^T of λ is the partition $(\lambda_1^T, \lambda_2^T, \dots)$ of n defined from $\lambda_i^T := \#\{j \in \mathbb{Z} \mid 0 < \lambda_j \leq i\}$. For example, if $\lambda = (\dots, 0, 2, 4, 0, 0, 1, 0, \dots)$ then $\lambda^+ = (2, 4, 1)$ and $\lambda^T = (3, 2, 1, 1, 0, 0, \dots)$. Also, we say that two compositions $\mu, \nu \vDash n$ are *equal up to translation and duality* if there exists $s \in \mathbb{Z}$ such that either $\mu_i = \nu_{s+i}$ for all $i \in \mathbb{Z}$ or $\mu_i = \nu_{s-i}$ for all $i \in \mathbb{Z}$.

Compositions $\lambda \vDash n$ may be identified with special elements of the weight lattice P of \mathfrak{sl}_{∞} via the dictionary $\lambda \vDash n \leftrightarrow \sum_{i \in \mathbb{Z}} \lambda_i \varepsilon_i \in P$. For example, $t\varepsilon_i$ is the composition whose i th part is equal to t , with all other parts being zero. Then the usual dominance order $<$ on P determined by the simple roots $\alpha_i = \varepsilon_i - \varepsilon_{i+1}$ corresponds to the partial order on compositions given by $\lambda < \mu$ if $\sum_{j \leq i} \lambda_j < \sum_{j \leq i} \mu_j$ for all i . If $\lambda \vDash n$ then $\lambda + \alpha_i \in P$ is a well-defined composition of n if and only if $\lambda_{i+1} > 0$, in which case it is the composition

with $\lambda_i + 1$ as its i th part, $\lambda_{i+1} - 1$ as its $(i + 1)$ th part, and all other parts the same as λ .

The point of this is that the set

$$\Xi(m|n) := \left\{ (\mu, \nu; t) \mid \begin{array}{l} 0 \leq t \leq m, \mu \vDash m - t, \nu \vDash n - t \\ \text{such that } \mu_i \nu_i = 0 \text{ for all } i \in \mathbb{Z} \end{array} \right\} \quad (4.11)$$

is in bijection with the set of linkage classes $\xi \in \text{Tab}_{\mathbb{Z}} / \approx$. To understand how this goes, given $(\mu, \nu; t) \in \Xi(m|n)$ and $\lambda \vDash t$, we define $A(\mu, \nu; \lambda)$ to be the unique anti-dominant tableau that has $\lambda_i + \mu_i$ entries equal to i in its top row, and $\lambda_i + \nu_i$ entries equal to i in its bottom row. For example, if $m = 3, n = 3, t = 1$ and $\mu \vDash 2, \nu \vDash 2$ and $\lambda \vDash 1$ are the compositions with $\mu_5 = 2, \nu_3 = \nu_4 = 1$ and $\lambda_j = 1$ for some $j \in \mathbb{Z}$, then

$$A(\mu, \nu; \lambda) = \begin{cases} \begin{array}{c} 55j \\ j43 \end{array} & \text{if } j \geq 5, \\ \begin{array}{c} 455 \\ 443 \end{array} & \text{if } j = 4, \\ \begin{array}{c} j55 \\ 43j \end{array} & \text{if } j \leq 3. \end{cases} \quad (4.12)$$

In general, the set $\{A(\mu, \nu; \lambda)\}_{\lambda \vDash t}$ is equal to ξ° for a unique $\xi \in \text{Tab}_{\mathbb{Z}} / \approx$ of atypicality t , and all linkage classes arise in this way.

Henceforth, we *identify* elements $(\mu, \nu; t)$ of $\Xi(m|n)$ with linkage classes $\xi \in \text{Tab}_{\mathbb{Z}} / \approx$ via the bijection described in the previous paragraph, denoting the block decompositions of $\mathcal{O}_{\mathbb{Z}}$ and $\overline{\mathcal{O}}_{\mathbb{Z}}$ instead by

$$\mathcal{O}_{\mathbb{Z}} = \bigoplus_{\xi \in \Xi(m|n)} \mathcal{O}_{\xi}, \quad \overline{\mathcal{O}}_{\mathbb{Z}} = \bigoplus_{\xi \in \Xi(m|n)} \overline{\mathcal{O}}_{\xi},$$

respectively. Thus, blocks are parameterized by an *atypicality* t and a *core* (μ, ν) . As usual, the indecomposable projective, standard and irreducible objects of \mathcal{O}_{ξ} are represented by the supermodules $P(\mathbf{A}), M(\mathbf{A})$ and $L(\mathbf{A})$ for $\mathbf{A} \in \xi$. For $\xi = (\mu, \nu; t) \in \Xi(m|n)$ and $\lambda \vDash t$, we will usually write $\overline{P}_{\xi}(\lambda), \overline{M}_{\xi}(\lambda)$ and $\overline{L}_{\xi}(\lambda)$ in place of $\overline{P}(\mathbf{A}), \overline{M}(\mathbf{A})$ and $\overline{L}(\mathbf{A})$ for $\mathbf{A} := A(\mu, \nu; \lambda)$. In this way, these families of objects in $\overline{\mathcal{O}}_{\xi}$ are now parameterized by compositions $\lambda \vDash t$ rather than by anti-dominant tableaux. For example, the block ξ associated to the anti-dominant tableau $\mathbf{A} = \begin{array}{c} 1133 \\ 4211 \end{array}$ has atypicality 2 and core (μ, ν) where $\mu, \nu \vDash 2$ have $\mu_3 = 2$ and $\nu_2 = \nu_4 = 1$. Moreover, $\overline{L}(\mathbf{A}) = \overline{L}_{\xi}(\lambda)$ where $\lambda \vDash 2$ has $\lambda_1 = 2$.

4.4. Formal characters. Let $\{\chi_i\}_{i \in \mathbb{Z}}$ be indeterminates. Set $\chi^{\alpha_i} := \chi_i \chi_{i+1}^{-1}$ and $\chi^{\eta} := \prod_{i \in \mathbb{Z}} \chi_i^{\eta_i}$ for $\eta \vDash m$. Let $e_r(\eta)$ be the r th elementary symmetric function $e_r(a_1, \dots, a_m)$ where a_1, \dots, a_m are chosen so that η_i of them are equal to i for each $i \in \mathbb{Z}$. Then, for any finite-dimensional W -module M , we define its η -weight space

$$M_{\eta} := \{v \in M \mid (d_1^{(r)} - e_r(\eta))^N v = 0 \text{ for } N \gg 0\}. \quad (4.13)$$

The *formal character* of M is

$$\text{ch } M := \sum_{\eta \vDash m} (\dim M_{\eta}) \chi^{\eta} \in \mathbb{Z}[\chi_i \mid i \in \mathbb{Z}]. \quad (4.14)$$

Theorem 4.10. *For $\xi = (\mu, \nu; t) \in \Xi(m|n)$ and $\lambda \models t$, we have that*

$$\text{ch } \overline{M}_\xi(\lambda) = \chi^{\lambda+\mu} \prod_{i \in \mathbb{Z}} (1 + \chi^{\alpha_i})^{\lambda_{i+1} + \mu_{i+1}}, \quad (4.15)$$

$$\text{ch } \overline{L}_\xi(\lambda) = \chi^{\lambda+\mu} \prod_{i \in \mathbb{Z}} (1 + \chi^{\alpha_i})^{\mu_{i+1}}. \quad (4.16)$$

Moreover, for any $M \in \overline{\mathcal{O}}_\xi$, we have that $M = \bigoplus_{\eta \models m} M_\eta$.

Proof. We first prove (4.15). Let $\mathbf{A} := \mathbf{A}(\mu, \nu; \lambda)$ and a_1, \dots, a_m be the entries along the top row of \mathbf{A} . By Corollary 3.15, we have that $\text{ch } \overline{M}_\xi(\lambda) = \text{ch } \overline{K}(\mathbf{A})$, and will compute the latter. The advantage of this is that $\overline{K}(\mathbf{A})$ is the restriction of the \mathfrak{h} -supermodule $K(\mathbf{A})$. Recalling (3.8), $K(\mathbf{A})$ possesses a basis of \mathfrak{t} -weight vectors $\{v_\theta\}_{\theta=(\theta_1, \dots, \theta_m) \in \{0,1\}^m}$ such that $S_{\rho'}(x_i)v_\theta = (a_i - \theta_i)v_\theta$ for each $i = 1, \dots, m$ (where $x_i = e_{i,i}$ as in §2.4). Hence,

$$S_{\rho'}(e_r(x_1, \dots, x_m))v_\theta = e_r(a_1 - \theta_1, \dots, a_m - \theta_m)v_\theta.$$

For two tuples $\theta, \theta' \in \{0,1\}^m$, we write $\theta' >_{\text{lex}} \theta$ if $\theta'_j > \theta_j, \theta'_{j+1} = \theta_{j+1}, \dots, \theta'_m = \theta_m$ for some $1 \leq j \leq m$. Then we observe that

$$d_1^{(r)}v_\theta = e_r(a_1 - \theta_1, \dots, a_m - \theta_m)v_\theta + (\text{a linear combination of } v_{\theta'}\text{'s for } \theta' >_{\text{lex}} \theta).$$

This follows from the explicit formula for $\pi(d_1^{(r)})$ recorded in the proof of [BBG, Lemma 8.3]; see also Lemma 3.19. Hence, we see that v_θ contributes the monomial $\chi_{a_1 - \theta_1} \cdots \chi_{a_m - \theta_m}$ to the formal character of $\overline{K}(\mathbf{A})$. We have now shown that

$$\text{ch } \overline{M}_\xi(\lambda) = \chi_{a_1} \cdots \chi_{a_m} \sum_{\theta \in \{0,1\}^m} \left[\prod_{\substack{1 \leq i \leq m \\ \theta_i = 1}} \chi^{\alpha_{a_i - 1}} \right],$$

which simplifies to give (4.15) since $\chi_{a_1} \cdots \chi_{a_m} = \chi^{\lambda+\mu}$.

The proof of (4.16) is very similar, using instead that $\text{ch } \overline{L}_\xi(\lambda) = \text{ch } \overline{V}(\mathbf{B})$ according to Theorem 3.3, where $\mathbf{B} \sim \mathbf{A}(\mu, \nu; \lambda)$ is chosen so that the entries along its top row are $b_1, \dots, b_{m-t}, c_1, \dots, c_t$ and the entry immediately below each of the c_i 's is another c_i . Then $V(\mathbf{B})$ possesses a basis of \mathfrak{t} -weight vectors $\{v_\theta\}_{\theta=(\theta_1, \dots, \theta_{m-t}) \in \{0,1\}^{m-t}}$ such that $S_{\rho'}(x_i)v_\theta = (b_i - \theta_i)v_\theta$ for $i = 1, \dots, m-t$ and $S_{\rho'}(x_i)v_\theta = c_i v_\theta$ for $i = m-t+1, \dots, m$. So the same argument as in the previous paragraph gives that

$$\text{ch } \overline{L}_\xi(\lambda) = \chi_{b_1} \cdots \chi_{b_{m-t}} \chi_{c_1} \cdots \chi_{c_t} \sum_{\theta \in \{0,1\}^{m-t}} \left[\prod_{\substack{1 \leq i \leq m-t \\ \theta_i = 1}} \chi^{\alpha_{b_i - 1}} \right],$$

which simplifies to give (4.16).

Finally, to get the last sentence, we just exhibited a basis showing that it is true for $M = \overline{L}_\xi(\lambda)$, which is enough to establish it in general. \square

Corollary 4.11. *The map $K_0(\overline{\mathcal{O}}_\xi) \rightarrow \mathbb{Z}[\chi_i \mid i \in \mathbb{Z}]$ given by $[M] \mapsto \text{ch}(M)$ is injective.*

Proof. By Theorem 4.10, $\text{ch } \overline{L}_\xi(\lambda)$ is equal to $\chi^{\lambda+\mu}$ plus a sum of terms of the form χ^ν for $\nu > \lambda + \mu$. Hence, the formal characters of the irreducible objects in $\overline{\mathcal{O}}_\xi$ are linearly independent, which implies the corollary. \square

The following lemma describes what the $e^{(r)}$'s and $f^{(r)}$'s do to weight spaces.

Lemma 4.12. *For any finite-dimensional W -module M and $\eta \vDash m$, we have that*

$$f^{(s_--r_1)} \dots f^{(s_--r_k)} M_\eta \subseteq \bigoplus_{\substack{\theta \vDash k \\ \theta_i \leq \eta_{i+1}}} M_{\eta + \sum_i \theta_i \alpha_i}, \quad (4.17)$$

$$e^{(s_++r_1)} \dots e^{(s_++r_k)} M_\eta \subseteq \bigoplus_{\substack{\theta \vDash k \\ \theta_i \leq \eta_i}} M_{\eta - \sum_i \theta_i \alpha_i} \quad (4.18)$$

for all $k \geq 0$ and $r_1, \dots, r_k > 0$.

Proof. We will prove (4.17). Then (4.18) follows by twisting with the involution $\iota : W \xrightarrow{\sim} W$ given by $d_i^{(r)} \mapsto (-1)^r d_i^{(r)}$, $e^{(s_++r)} \mapsto (-1)^r f^{(s_--r)}$, and $f^{(s_--r)} \mapsto (-1)^r e^{(s_++r)}$; note for this that $\iota^*(M_\eta) = \iota^*(M)_{\eta'}$ where $\eta'_i = \eta_{-i}$.

To establish (4.17), let W_1^0 (resp. W_1^\flat) be the subalgebra of W generated by $d_1^{(1)}, \dots, d_1^{(m)}$ (resp. by $d_1^{(1)}, \dots, d_1^{(m)}, f^{(s_--1)}, \dots, f^{(s_--m)}$). For $\eta \vDash m$, we define the weight spaces M_η of a finite-dimensional W_1^\flat -module M by the same formula (4.13) as before. Let \mathbb{C}_η be a one-dimensional W_1^0 -module with basis $\bar{1}_\eta$ such that $d_1^{(r)} \bar{1}_\eta = e_r(\eta) \bar{1}_\eta$ for each r . Then form the induced module $\bar{M}(\eta) := W_1^\flat \otimes_{W_1^0} \mathbb{C}_\eta$, setting $\bar{m}_\eta := 1 \otimes \bar{1}_\eta$. We claim that

$$f^{(s_--r_1)} \dots f^{(s_--r_k)} \bar{m}_\eta \in \bigoplus_{\substack{\theta \vDash k \\ \theta_i \leq \eta_{i+1}}} \bar{M}(\eta)_{\eta + \sum_i \theta_i \alpha_i}. \quad (4.19)$$

To deduce (4.17) from this claim, it suffices to show for any $v \in M_\eta$ that is a simultaneous eigenvector for all $d_1^{(1)}, \dots, d_1^{(m)}$ that $f^{(s_--r_1)} \dots f^{(s_--r_k)} v$ belongs to the subspace on the right hand side of (4.17). This follows from (4.19) because there is a unique W_1^\flat -module homomorphism $\omega : \bar{M}(\eta) \rightarrow M$ such that $\bar{m}_\eta \mapsto v$, and $\omega\left(\bar{M}(\eta)_{\eta + \sum_i \theta_i \alpha_i}\right) \subseteq M_{\eta + \sum_i \theta_i \alpha_i}$.

Finally, to prove (4.19), we pick any block $\xi = (0, \nu; m) \in \Xi(m|n)$ of maximal atypicality. Applying the PBW theorem for W , we see that W_1^\flat -module $\bar{M}(\eta)$ may be identified with the restriction of the Verma supermodule $\bar{M}_\xi(\eta)$, so that \bar{m}_η is the highest weight vector in $\bar{M}_\xi(\eta)$. As \bar{m}_η is a $d_1^{(1)}$ -eigenvector of eigenvalue $\sum_{i \in \mathbb{Z}} i \eta_i$, the relations imply that $f^{(s_--r_1)} \dots f^{(s_--r_k)} \bar{m}_\eta$ is a $d_1^{(1)}$ -eigenvector of eigenvalue $\sum_{i \in \mathbb{Z}} i \eta_i - k$. Hence, this vector lies in the sum of the weight spaces $\bar{M}_\xi(\eta)_{\eta'}$ for $\eta' \vDash m$ with $\sum_i i \eta'_i = \sum_i i \eta_i - k$. Finally, we apply (4.15) to see that $\bar{M}_\xi(\eta)_{\eta'}$ is zero unless $\eta' = \eta + \sum_i \theta_i \alpha_i$ for $\theta \vDash k$ with $\theta_i \leq \eta_{i+1}$ for all i . \square

4.5. Cartan matrix of $\bar{\mathcal{O}}_\xi$. The next goal is to calculate the Cartan matrix of the block $\bar{\mathcal{O}}_\xi$. We will deduce this from the following lemma describing the composition multiplicities in the Verma supermodules $\bar{M}_\xi(\lambda)$. This is a reformulation of Corollary 3.16, but we will give an alternative proof here using the formal characters computed in Theorem 4.10.

Lemma 4.13. *For any $\lambda, \kappa \vDash t$, the Verma multiplicity $[\bar{M}_\xi(\lambda) : \bar{L}_\xi(\kappa)]$ is non-zero if and only if $\kappa = \lambda + \sum_i \theta_i \alpha_i$ for $\theta = (\theta_i)_{i \in \mathbb{Z}}$ satisfying $0 \leq \theta_i \leq \lambda_{i+1}$*

for all i ; equivalently, $\lambda = \kappa - \sum_i \theta_i \alpha_i$ for θ with $0 \leq \theta_i \leq \kappa_i$ for all i . When this holds, we have that

$$[\overline{M}_\xi(\lambda) : \overline{L}_\xi(\kappa)] = \prod_{i \in \mathbb{Z}} \binom{\lambda_{i+1}}{\theta_i}.$$

Proof. In view of Corollary 4.11, this follows from the following calculation:

$$\begin{aligned} \text{ch } \overline{M}_\xi(\lambda) &= \chi^{\lambda+\mu} \prod_{i \in \mathbb{Z}} (1 + \chi^{\alpha_i})^{\lambda_{i+1} + \mu_{i+1}} \\ &= \left(\chi^\lambda \prod_{i \in \mathbb{Z}} (1 + \chi^{\alpha_i})^{\lambda_{i+1}} \right) \left(\chi^\mu \prod_{i \in \mathbb{Z}} (1 + \chi^{\alpha_i})^{\mu_{i+1}} \right) \\ &= \left(\sum_{\substack{\theta = (\theta_i)_{i \in \mathbb{Z}} \\ 0 \leq \theta_i \leq \lambda_{i+1}}} \chi^\lambda \prod_{i \in \mathbb{Z}} \binom{\lambda_{i+1}}{\theta_i} (\chi^{\alpha_i})^{\theta_i} \right) \left(\chi^\mu \prod_{i \in \mathbb{Z}} (1 + \chi^{\alpha_i})^{\mu_{i+1}} \right) \\ &= \sum_{\substack{\theta = (\theta_i)_{i \in \mathbb{Z}} \\ 0 \leq \theta_i \leq \lambda_{i+1}}} \left(\prod_{i \in \mathbb{Z}} \binom{\lambda_{i+1}}{\theta_i} \right) \text{ch } \overline{L}_\xi \left(\lambda + \sum_{i \in \mathbb{Z}} \theta_i \alpha_i \right). \end{aligned}$$

Here we have used both of the formulae from Theorem 4.10. \square

Theorem 4.14. For $\xi = (\mu, \nu; t) \in \Xi(m|n)$ and any $\lambda, \kappa \models t$, the multiplicity $[\overline{P}_\xi(\lambda) : \overline{L}_\xi(\kappa)]$ is non-zero if and only if $\kappa = \lambda + \sum_i (\lambda_{i+1} - \rho_{i+1}) \alpha_i$ for $\rho = (\rho_i)_{i \in \mathbb{Z}}$ satisfying $0 \leq \rho_{i+1} \leq \lambda_{i+1} + \min(\lambda_i, \rho_i)$ for all i , in which case

$$[\overline{P}_\xi(\lambda) : \overline{L}_\xi(\kappa)] = m! n! \sum_{\substack{\tau = (\tau_i)_{i \in \mathbb{Z}} \\ \max(\lambda_{i+1}, \rho_{i+1}) \leq \tau_{i+1} \leq \lambda_{i+1} + \min(\lambda_i, \rho_i)}} \prod_{i \in \mathbb{Z}} \frac{\binom{\lambda_{i+1} + \tau_i - \tau_{i+1}}{\tau_i - \lambda_i} \binom{\lambda_{i+1} + \tau_i - \tau_{i+1}}{\tau_i - \rho_i}}{(\lambda_{i+1} + \tau_i - \tau_{i+1})! (\lambda_{i+1} + \tau_i - \tau_{i+1} + \gamma_i)!},$$

where $\gamma := \mu + \nu$. Moreover, $[\overline{P}_\xi(\lambda) : \overline{L}_\xi(\kappa)] = [\overline{P}_\xi(\kappa) : \overline{L}_\xi(\lambda)]$.

Proof. Let $\mathbf{A} := \mathbf{A}(\mu, \nu; \kappa)$ and $\mathbf{C} := \mathbf{A}(\mu, \nu; \lambda)$. Since these are anti-dominant, Theorem 3.17 and the exactness of H_0 imply that the multiplicity we are trying to compute is equal to $[P(\mathbf{C}) : L(\mathbf{A})]$. This can be computed by the usual BGG reciprocity formula in the highest weight category $\mathcal{O}_{\mathbb{Z}}$:

$$[P(\mathbf{C}) : L(\mathbf{A})] = \sum_{\mathbf{B} \in \xi} [M(\mathbf{B}) : L(\mathbf{A})][M(\mathbf{B}) : L(\mathbf{C})]. \quad (4.20)$$

In particular, this already establishes the symmetry property at the end of the statement of the theorem.

For any $\mathbf{B} \in \xi$, Theorem 3.14 shows that $H_0(M(\mathbf{B}))$ is isomorphic to $\overline{M}_\xi(\beta)$ for $\beta \models t$ determined uniquely from $\mathbf{A}(\mu, \nu; \beta) \sim \mathbf{B}$. Also, for a given β , the number of different \mathbf{B} satisfying $\mathbf{B} \sim \mathbf{A}(\mu, \nu; \beta)$ is

$$m! n! / \prod_i (\beta_i + \mu_i)! (\beta_i + \nu_i)! = m! n! / \prod_i \beta_i! (\beta_i + \gamma_i)!.$$

We deduce from (4.20) that

$$[\overline{P}_\xi(\lambda) : \overline{L}_\xi(\kappa)] = m! n! \sum_{\beta \models t} \left(\prod_i \frac{1}{\beta_i! (\beta_i + \gamma_i)!} \right) [\overline{M}_\xi(\beta) : \overline{L}_\xi(\lambda)] [\overline{M}_\xi(\beta) : \overline{L}_\xi(\kappa)].$$

are counting its lattice points. Theorem 4.14 tells us that $h(\lambda)$ is the number of $\kappa \models t$ such that $\overline{L}_\xi(\kappa)$ appears as a composition factor of $\overline{P}_\xi(\lambda)$.

If $\lambda_j = 0$ for some j , then we have by the definition that

$$h(\lambda) = h(\lambda^{\leq j}) h(\lambda^{\geq j}), \quad (4.22)$$

where $\lambda^{\leq j}$ (resp. $\lambda^{\geq j}$) is the composition obtained from λ by setting all parts in positions $> j$ (resp. $< j$) to zero. Also it is clear that $h(\lambda) = h(\mu)$ if μ is obtained from λ by a translation. This reduces the problem of computing $h(\lambda)$ to the case that λ is *connected*, i.e. it is of the form

$$h(\lambda_1, \dots, \lambda_r) := h((\dots, 0, \lambda_1, \dots, \lambda_r, 0, \dots))$$

for some r and $\lambda_1, \dots, \lambda_r > 0$.

Lemma 4.16. *For any $\lambda \models t$, we have that $h(\lambda) \geq \binom{t+2}{2}$, with equality if and only if $\lambda = t\varepsilon_i$ for some $i \in \mathbb{Z}$.*

Proof. It is easy to check that $h(t\varepsilon_i) = \binom{t+2}{2}$. Conversely, we must show that $h(\lambda) > \binom{t+2}{2}$ whenever it has at least two non-zero parts. If λ is not connected, the proof is easily computed by induction on t , using the elementary inequality $\binom{t'+2}{2} \binom{t''+2}{2} > \binom{t+2}{2}$ if $t', t'' \geq 1$ satisfy $t' + t'' = t$. If λ is connected, let λ_k, λ_{k+1} be its rightmost two non-zero parts, so $\lambda = (\dots, \lambda_{k-1}, \lambda_k, \lambda_{k+1})$. Then the conclusion follows easily from the claim that

$$h(\dots, \lambda_{k-1}, \lambda_k, \lambda_{k+1}) > h(\dots, \lambda_{k-1}, \lambda_k + \lambda_{k+1}).$$

This can be proved by showing for each choice of the entries of ρ up to ρ_{k-1} that there are more ways of extending this to a sequence satisfying (4.21) for $\lambda = (\dots, \lambda_{k-1}, \lambda_k, \lambda_{k+1}, \dots)$ than for $\lambda' = (\dots, \lambda_{k-1}, \lambda_k + \lambda_{k+1}, \dots)$. \square

Remark 4.17. We also expect for $\lambda \models t$ that $h(\lambda) \leq 3^t$, with equality if and only if λ is generic in the sense that it consists of isolated 1's. We won't need this observation here, but note that $h(\lambda) = 3^t$ for generic λ follows from $h(\varepsilon_i) = 3$ and (4.22).

Lemma 4.18. *The space $\text{Hom}_{\overline{\mathcal{O}}_\xi}(\overline{P}_\xi(t\varepsilon_j), \overline{P}_\xi(t\varepsilon_i))$ is non-zero if and only if $|i - j| \leq 1$.*

Proof. This is the multiplicity computed in Theorem 4.14 for $\lambda = t\varepsilon_i$ and $\kappa = t\varepsilon_j$. Since the Cartan matrix is symmetric, we may assume that $i < j$. From the equation $\lambda - \kappa = \sum_i (\rho_i - \lambda_i) \alpha_{i-1}$, we deduce that we must have $\rho_i = \rho_{i+1} = \dots = \rho_j = t$ and all other parts zero. But this contradicts the system of inequalities $\rho_{k+1} \leq \lambda_{k+1} + \min(\lambda_k, \rho_k)$ unless we actually have that $j = i + 1$. \square

Lemma 4.19. *Let $\gamma := \mu + \nu$. For each $i \in \mathbb{Z}$ we have that*

$$\dim \text{End}_{\overline{\mathcal{O}}_\xi}(\overline{P}_\xi(t\varepsilon_i)) = \frac{m!n!}{t! \prod_j \gamma_j!} \sum_{r=0}^t \binom{t}{r} \frac{\gamma_i! \gamma_{i+1}!}{(\gamma_i + t - r)! (\gamma_{i+1} + r)!}.$$

This is equal to $\frac{m!n!}{(t!)^2 \prod_j \gamma_j!} \binom{2t}{t}$ whenever $\gamma_i = \gamma_{i+1} = 0$.

Proof. The first formula is easily derived from Theorem 4.14; in the summation over τ there, one just has τ 's with $\tau_i = t, \tau_{i+1} = r$ for $0 \leq r \leq t$, and all other parts equal to zero. To deduce the second formula, use $\sum_{r=0}^t \binom{t}{r}^2 = \binom{2t}{t}$. \square

4.6. An idempotented form for W . In general, the blocks $\overline{\mathcal{O}}_\xi$ of the category $\overline{\mathcal{O}}_\mathbb{Z}$ have infinitely isomorphism classes of irreducible objects. In such situations, it is often appropriate to consider locally unital rather than unital algebras. Here, by a *locally unital algebra*, we mean an associative algebra A equipped with a distinguished system of mutually orthogonal idempotents $\{1_i\}_{i \in I}$ such that $A = \bigoplus_{i,j \in I} 1_i A 1_j$. By a (left) module M over a locally unital algebra A , we always mean a module as usual which is locally unital in the sense that $M = \bigoplus_{i \in I} 1_i M$. Let $A\text{-mod}_{\text{fd}}$ be the category of all such modules with $\dim M < \infty$. In this subsection, we construct a locally unital algebra W_ξ whose module category is equivalent to $\overline{\mathcal{O}}_\xi$. In the next subsection, we will show for maximally atypical blocks that W_ξ is isomorphic to the locally unital endomorphism algebra of a minimal projective generating family in $\overline{\mathcal{O}}_\xi$. For further discussion of our motivation, see [B5, §4].

To define W_ξ , we must first pass from W to an idempotented form \dot{W} . By definition, this is the locally unital algebra with distinguished idempotents $\{1_\eta\}_{\eta \models m}$, generators

$$\begin{aligned} c^{(r)} 1_\eta &\in 1_\eta \dot{W} 1_\eta && \text{for } \eta \models m \text{ and } r \geq 0, \\ d^{(r)} 1_\eta &\in 1_\eta \dot{W} 1_\eta && \text{for } \eta \models m \text{ and } r \geq 0, \\ f_i^{(r)} 1_\eta &\in 1_{\eta + \alpha_i} \dot{W} 1_\eta && \text{for } \eta \models m, r > s_- \text{ and } i \in \mathbb{Z} \text{ such that } \eta_{i+1} > 0, \\ e_i^{(r)} 1_\eta &\in 1_{\eta - \alpha_i} \dot{W} 1_\eta && \text{for } \eta \models m, r > s_+ \text{ and } i \in \mathbb{Z} \text{ such that } \eta_i > 0, \end{aligned}$$

and certain relations. In order to write these down, we need a couple of conventions. We interpret the following currently undefined expressions as zero: 1_η , $c^{(r)} 1_\eta$ and $d^{(r)} 1_\eta$ if $\eta \not\models m$; $e_i^{(r)} 1_\eta$ if either $\eta \not\models m$ or $\eta - \alpha_i \not\models m$; $f_i^{(r)} 1_\eta$ if either $\eta \not\models m$ or $\eta + \alpha_i \not\models m$. This means that for given $\eta \models m$, we have $e_i^{(r)} 1_\eta = f_i^{(r)} 1_\eta = 0$ for all but finitely many i . Also we will omit idempotents from the middles of monomials when they are clear from the context. Then, the relations are as follows:

$$c^{(0)} 1_\eta = d^{(0)} 1_\eta = 1_\eta, \quad d^{(1)} 1_\eta = \sum_i i \eta_i 1_\eta, \quad d^{(r)} 1_\eta = 0 \text{ for } r > m, \quad (4.23)$$

$$c^{(r)} 1_\eta \text{ is central,} \quad d^{(r)} d^{(s)} 1_\eta = d^{(s)} d^{(r)} 1_\eta, \quad (4.24)$$

$$d^{(r)} e_i^{(s)} 1_\eta - e_i^{(s)} d^{(r)} 1_\eta = \sum_{a=0}^{r-1} d^{(a)} e_i^{(r+s-1-a)} 1_\eta, \quad (4.25)$$

$$d^{(r)} f_i^{(s)} 1_\eta - f_i^{(s)} d^{(r)} 1_\eta = - \sum_{a=0}^{r-1} f_i^{(r+s-1-a)} d^{(a)} 1_\eta, \quad (4.26)$$

$$e_i^{(r)} e_j^{(s)} 1_\eta + e_j^{(r)} e_i^{(s)} 1_\eta + e_i^{(s)} e_j^{(r)} 1_\eta + e_j^{(s)} e_i^{(r)} 1_\eta = 0, \quad (4.27)$$


$$f_i^{(r)} f_j^{(s)} 1_\eta + f_j^{(r)} f_i^{(s)} 1_\eta + f_i^{(s)} f_j^{(r)} 1_\eta + f_j^{(s)} f_i^{(r)} 1_\eta = 0, \quad (4.28)$$

$$e_i^{(r)} f_j^{(s)} 1_\eta + f_j^{(s)} e_i^{(r)} 1_\eta = 0 \text{ for } i \neq j, \quad (4.29)$$

$$\sum_i (e_i^{(r)} f_i^{(s)} 1_\eta + f_i^{(s)} e_i^{(r)} 1_\eta) = c^{(r+s-1)} 1_\eta. \quad (4.30)$$

We leave it as an exercise for the reader to check in the degenerate case $m = 0$ that $\dot{W} = \mathbb{C}[c^{(1)}, \dots, c^{(n)}]$, i.e. it is a polynomial algebra in n variables. The case $m = 1$ is also particularly easy to understand; in this case, we let $1_i := 1_{\varepsilon_i}$, $e1_i := e_i^{(s+1)}1_{\varepsilon_i}$, $f1_i := f_{i-1}^{(s+1)}1_{\varepsilon_i}$ and $z1_i := c^{(n)}1_{\varepsilon_i}$ for short.

Lemma 4.20. *In the case $m = 1$, the algebra \dot{W} is $\mathbb{C}[c^{(1)}, \dots, c^{(n-1)}] \otimes T$ where T is the path algebra of the infinite quiver*



subject to the relations $e^21_i = f^21_i = 0$ and $ef1_i + fe1_i = z1_i$ for all $i \in \mathbb{Z}$.

Proof. It is easy to check from the defining relations for \dot{W} that there is a locally unital algebra homomorphism $f : \mathbb{C}[c^{(1)}, \dots, c^{(n-1)}] \otimes T \rightarrow \dot{W}$ sending $1_i, c^{(r)}1_i, e1_i, f1_i$ and $z1_i$ to the corresponding elements of \dot{W} . To show this is an isomorphism, we define $f^{-1} : \dot{W} \rightarrow \mathbb{C}[c^{(1)}, \dots, c^{(n-1)}] \otimes T$ by sending

$$\begin{aligned} 1_{\varepsilon_i} &\mapsto 1_i, \\ c^{(r)}1_{\varepsilon_i} &\mapsto \begin{cases} c^{(r)}1_i & \text{if } r < n, \\ (-1)^{r-n}((i+1)^{r-n+1} - i^{r-n+1})z1_i & \text{if } r \geq n, \end{cases} \\ d^{(r)}1_{\varepsilon_i} &\mapsto \delta_{r,1}i1_i, \\ e_j^{(s+r)}1_{\varepsilon_i} &\mapsto \delta_{i,j}(-i-1)^{r-1}e1_i, \\ f_j^{(s+r)}1_{\varepsilon_i} &\mapsto \delta_{i,j-1}(-i)^{r-1}f1_i, \end{aligned}$$

for each $r \geq 1$ and $i, j \in \mathbb{Z}$. This is well defined by another (longer) relation check. Finally, it is obvious that $f^{-1} \circ f = \text{id}$. To see that $f \circ f^{-1} = \text{id}$, note from the relations that the elements $1_i, e1_i, f1_i$ and $c^{(1)}1_i, \dots, c^{(n)}1_i$ for all $i \in \mathbb{Z}$ already suffice to generate \dot{W} . \square

Lemma 4.21. *The following two categories may be identified:*

- (1) *The full subcategory of $W\text{-mod}_{\text{fd}}$ consisting of all modules M such that M is equal to the direct sum $\bigoplus_{\eta \models m} M_{\eta}$ of its weight spaces from (4.13) and the endomorphism defined by the action of $d_1^{(1)}$ is diagonalizable;*
- (2) *The full subcategory of $\dot{W}\text{-mod}_{\text{fd}}$ consisting of all modules M such that $d^{(r)}1_{\eta} - e_r(\eta)1_{\eta}$ acts nilpotently for all $\eta \models m$ and $r > 1$.*

Proof. Take M belonging to the category (1). We make it into a \dot{W} -module by declaring that

- 1_{η} acts as the projection pr_{η} along the weight space decomposition;
- $c^{(r)}1_{\eta}$ acts as $c^{(r)} \circ \text{pr}_{\eta}$ and $d^{(r)}1_{\eta}$ acts as $d_1^{(r)} \circ \text{pr}_{\eta}$; and
- $e_i^{(r)}1_{\eta}$ acts as $\text{pr}_{\eta - \alpha_i} \circ e^{(r)} \circ \text{pr}_{\eta}$ and $f_i^{(r)}1_{\eta}$ acts as $\text{pr}_{\eta + \alpha_i} \circ f^{(r)} \circ \text{pr}_{\eta}$.

To see that this makes sense, we need to verify that the defining relations of \dot{W} are satisfied. This follows from the relations for W in Theorem 3.1. For example, to check (4.29)–(4.30), we have by the definition of the action of the

generators of \dot{W} on $v \in M_\eta$ that

$$\begin{aligned} & \sum_{i,j} (e_i^{(r)} f_j^{(s)} 1_\eta + f_j^{(s)} e_i^{(r)} 1_\eta) v \\ &= \sum_{i,j} (\text{pr}_{\eta+\alpha_j-\alpha_i} \circ e^{(r)} \circ \text{pr}_{\eta+\alpha_j} \circ f^{(s)} + \text{pr}_{\eta-\alpha_i+\alpha_j} \circ f^{(s)} \circ \text{pr}_{\eta-\alpha_i} \circ e^{(r)}) v \\ &= (e^{(r)} f^{(s)} + f^{(s)} e^{(r)}) v = \delta_{i,j} c^{(r+s-1)} v = c^{(r+s-1)} 1_\eta v, \end{aligned}$$

using Lemma 4.12 with $k = 1$ for the second equality. Applying $\text{pr}_{\eta+\alpha_j-\alpha_i}$ to both sides, this establishes (4.29) when $i \neq j$ and (4.30) when $i = j$.

Conversely, take M belonging to the category (2). Then we make it into a W -module by declaring that

- $c^{(r)}, d_1^{(r)}, e^{(r)}$ and $f^{(r)}$ act on $1_\eta M$ as $c^{(r)} 1_\eta, d^{(r)} 1_\eta, \sum_i e_i^{(r)} 1_\eta$ and $\sum_i f_i^{(r)} 1_\eta$;
- $d_2^{(r)}$ acts via the formula $d_2^{(r)} = \sum_{s=0}^r d_1^{(s)} c^{(r-s)}$ from (3.5).

Then one needs to verify that the relations for W from Theorem 3.1 are satisfied. Moreover, we have that $M_\eta = 1_\eta M$.

Finally, the two constructions just explained are inverses of each other. Again, this uses the $k = 1$ case of Lemma 4.12. \square

Theorem 4.22. *For $\xi = (\mu, \nu; t) \in \Xi(m|n)$, the category $\overline{\mathcal{O}}_\xi$ is isomorphic to the category $W_\xi\text{-mod}_{\text{fd}}$, where*

$$W_\xi := \dot{W} / \bigcap_{\lambda \vDash t} \text{Ann}_{\dot{W}}(\overline{P}_\xi(\lambda)).$$

This is a locally unital algebra with distinguished idempotents $\{1_\eta\}_{\eta \vDash m}$ that are the images of the ones in \dot{W} . Moreover:

- (1) *All of the left ideals $W_\xi 1_\eta$ and right ideals $1_\nu W_\xi$ are finite-dimensional.*
- (2) *We have that $c^{(1)} 1_\eta = \sum_i i(\nu_i - \mu_i) 1_\eta$ in W_ξ .*

Proof. Set $V := \bigoplus_{\lambda \vDash t} \overline{P}_\xi(\lambda)$. For each $\eta \vDash m$, there are only finitely many $\kappa \vDash t$ such that $1_\eta \overline{L}_\xi(\kappa) \neq 0$. This follows from Theorem 4.10: we must have that $\kappa = \eta - \mu - \sum_i \theta_i \alpha_i$ for $\theta = (\theta_i)_{i \in \mathbb{Z}}$ with $0 \leq \theta_i \leq \mu_{i+1}$, and there are only finitely many such θ 's. Moreover, for each $\kappa \vDash t$, there are only finitely many $\lambda \vDash t$ such that $[\overline{P}_\xi(\lambda) : \overline{L}_\xi(\kappa)] \neq 0$, as is clear from Theorem 4.14. Hence, there are only finitely many $\lambda \vDash t$ such that $1_\eta \overline{P}_\xi(\lambda) \neq 0$. Thus, we have shown that all of the weight spaces $1_\eta V$ of the \dot{W} -module V are finite-dimensional.

Consider the locally unital algebra

$$E := \bigoplus_{\eta, \nu \vDash m} \text{Hom}_{\mathbb{C}}(1_\eta V, 1_\nu V)$$

with multiplication coming from the usual composition. This is a simple locally unital matrix algebra with unique (up to isomorphism) irreducible module V . The representation of \dot{W} on V defines a locally unital algebra homomorphism $\rho : \dot{W} \rightarrow E$ sending $a \in 1_\nu \dot{W} 1_\eta$ to the linear map $1_\eta V \rightarrow 1_\nu V$ defined by left multiplication by a . We have that $\ker \rho = \bigcap_{\lambda \vDash t} \text{Ann}_{\dot{W}} \overline{P}_\xi(\lambda)$, hence, ρ induces an embedding $W_\xi \hookrightarrow E$. Since each $1_\nu E 1_\eta = \text{Hom}_{\mathbb{C}}(1_\eta V, 1_\nu V)$ is finite-dimensional, we deduce that each $1_\nu W_\xi 1_\eta$ is finite-dimensional. Also $1_\nu W_\xi 1_\eta$ is non-zero only if there exists $\lambda \vDash t$ such that $1_\nu \overline{P}_\xi(\lambda) \neq 0 \neq 1_\eta \overline{P}_\xi(\lambda)$. For

fixed η , there are only finitely many such λ , hence, only finitely many such ν . This shows that $W_\xi 1_\eta$ is finite-dimensional. Similarly, so is $1_\nu W_\xi$, and we have established (1).

Now we explain how to identify $\overline{\mathcal{O}}_\xi$ with $W_\xi\text{-mod}_{\text{fd}}$. Given any $M \in \overline{\mathcal{O}}_\xi$, we can forget the $\mathbb{Z}/2$ -grading then apply Lemma 4.21 to view it as a \dot{W} -module, using also Lemma 3.8 and the last part of Theorem 4.10. This defines a functor from $\overline{\mathcal{O}}_\xi$ to the full subcategory of $\dot{W}\text{-mod}_{\text{fd}}$ consisting of subquotients of finite direct sums of the modules $\{\overline{P}_\xi(\lambda)\}_{\lambda \models t}$. Each $\overline{P}_\xi(\lambda)$ factors through the quotient W_ξ , so the latter category may also be described as the full subcategory of $W_\xi\text{-mod}_{\text{fd}}$ consisting of subquotients of finite direct sums of the modules $\{\overline{P}_\xi(\lambda)\}_{\lambda \models t}$. In fact, this defines an isomorphism of categories, since the $\mathbb{Z}/2$ -grading on M can be recovered uniquely thanks to Remark 4.2.

It just remains to observe that *every* finite-dimensional W_ξ -module belongs to the full subcategory of $W_\xi\text{-mod}_{\text{fd}}$ consisting of subquotients of finite direct sums of the modules $\{\overline{P}_\xi(\lambda)\}_{\lambda \models t}$. To see this, note that any $M \in W_\xi\text{-mod}_{\text{fd}}$ is a quotient of a finite direct sum of the projective modules $W_\xi 1_\eta$ for $\eta \models m$. Moreover, $W_\xi 1_\eta \hookrightarrow E 1_\eta$, which is a direct sum of copies of V , so that $W_\xi 1_\eta$ embeds into a (possibly infinite) direct sum of the modules $\overline{P}_\xi(\lambda)$. In fact, it embeds into a finite direct sum of these modules since it is finite-dimensional, as is each $\overline{P}_\xi(\lambda)$.

To establish (2), note that $c^{(1)} = d_2^{(1)} - d_1^{(1)}$, so it acts diagonalizably on any object of $\overline{\mathcal{O}}_\xi$ thanks to Lemma 3.8. Moreover, $d_2^{(1)} - d_1^{(1)}$ acts on $\overline{L}(\mathbf{A})$ as $b(\mathbf{A}) - a(\mathbf{A}) = \sum_i i(\nu_i - \mu_i)$ for any $\mathbf{A} \in \xi^\circ$. Thus, $c^{(1)} - \sum_i i(\nu_i - \mu_i)$ annihilates all $\overline{P}_\xi(\lambda)$, and (2) follows. \square

Remark 4.23. Here is a slightly different construction of the locally unital algebra W_ξ . Given any finite subset $X \subset \xi$, the W -module $\bigoplus_{\lambda \in X} \overline{P}_\xi(\lambda)$ is finite-dimensional. Hence,

$$W_X := W / \bigcap_{\lambda \in X} \text{Ann}_W(\overline{P}_\xi(\lambda))$$

is a finite-dimensional algebra. Each W_X possesses a distinguished family of idempotents $\{1_\eta\}_{\eta \models m}$ such that $W_X = \bigoplus_{\nu, \eta \models m} 1_\nu W_X 1_\eta$, namely, 1_η is the primitive idempotent in the finite-dimensional commutative subalgebra of W_X generated by $d_1^{(1)}, \dots, d_m^{(1)}$ that projects any module onto its η -weight space. Then W_ξ is the inverse limit $\varprojlim_X W_X$ over all finite subsets X of ξ , taking the inverse limit in the category of locally unital algebras with idempotents indexed by compositions of m .

Remark 4.24. The special case $m = n = 1$ is particularly trivial. If $m = n = t = 1$ then W_ξ is the algebra T from Lemma 4.20 subject to the additional relations $z 1_i = 0$ for all $i \in \mathbb{Z}$. If $m = n = 1$ and $t = 0$, then we have that $\mu = \varepsilon_i$ and $\nu = \varepsilon_j$ for $i \neq j$, and W_ξ is the algebra T with the additional relations $z 1_i = (j - i) 1_i, z 1_{i-1} = (j - i) 1_{i-1}$, and $1_k = 0$ for $k \neq i, i - 1$. (In this case, $W_\xi \cong M_2(\mathbb{C})$.)

4.7. Graded lifts and the Soergel algebra of a block. Throughout the subsection, we fix $\xi = (\mu, \nu; t) \in \Xi(m|n)$ and set $\gamma := \mu + \nu$ for short. Let

$$A_\xi := \bigoplus_{A, B \in \xi} \text{Hom}_{\mathfrak{g}}(P(A), P(B)), \quad (4.31)$$

viewed as an algebra with multiplication that is the opposite of composition. Note A_ξ is a locally unital in the sense introduced at the start of §4.6, with distinguished idempotents $\{1_A\}_{A \in \xi}$ coming from the identity endomorphisms of each $P(A)$. By standard theory, the functor

$$\bigoplus_{A \in \xi} \text{Hom}_{\mathfrak{g}}(P(A), -) : \mathcal{O}_\xi \rightarrow A_\xi\text{-mod}_{\text{fd}} \quad (4.32)$$

is an equivalence of categories. Since we have that $1_A A_\xi 1_B = \text{Hom}_{\mathfrak{g}}(P(A), P(B))$, this functor sends $P(B)$ to the left ideal $A_\xi 1_B$.

By [BLW, Theorem 5.26] (plus [BLW, Theorem 3.10]), the basic algebra A_ξ admits a positive grading $A_\xi = \bigoplus_{d \geq 0} (A_\xi)_d$ making it into a Koszul algebra. In view of (4.32), this means that we can introduce a *graded lift* of the block \mathcal{O}_ξ , namely, the category $A_\xi\text{-grmod}_{\text{fd}}$ of finite-dimensional graded A_ξ -modules. This is entirely analogous to the situation in [S], where Soergel introduced a graded lift of the category \mathcal{O} for a semisimple Lie algebra. For further discussion, see [BLW, §§5.2–5.5]. In particular, [BLW, Theorem 5.11] shows that categorical action on $\mathcal{O}_{\mathbb{Z}}$ discussed in §4.1 also lifts to the graded setting. Note finally by [BGS, Corollary 2.5.2] (which extends obviously to the present locally unital setting) that the Koszul grading on A_ξ is unique up to automorphism. This means that the grading is canonical, so it can be used to refine the invariants of blocks computed already in Theorem 4.14; see Theorem 4.27 below.

Recalling next that the indecomposable projective objects in the quotient category $\overline{\mathcal{O}}_\xi$ are the W -supermodules $\{\overline{P}_\xi(\lambda)\}_{\lambda \vDash t}$, we can also consider the locally unital algebra

$$B_\xi := \bigoplus_{\kappa, \lambda \vDash t} \text{Hom}_W(\overline{P}_\xi(\kappa), \overline{P}_\xi(\lambda)), \quad (4.33)$$

again with multiplication that is opposite of composition. Its distinguished idempotents are denoted $\{1_\lambda\}_{\lambda \vDash t}$, so that $1_\kappa B_\xi 1_\lambda = \text{Hom}_W(\overline{P}_\xi(\kappa), \overline{P}_\xi(\lambda))$. In view of the double centralizer property of Corollary 4.9, we can identify B_ξ with the subalgebra $\bigoplus_{\kappa, \lambda \vDash t} 1_{A(\mu, \nu; \kappa)} A_\xi 1_{A(\mu, \nu; \lambda)}$ of A_ξ . In particular, B_ξ inherits a positive grading $B_\xi = \bigoplus_{d \geq 0} (B_\xi)_d$ from the Koszul grading on A_ξ . We call the graded algebra B_ξ the *Soergel algebra* of the block \mathcal{O}_ξ . Just like in (4.32), there is an equivalence of categories

$$\bigoplus_{\kappa \vDash t} \text{Hom}_W(\overline{P}_\xi(\kappa), -) : \overline{\mathcal{O}}_\xi \rightarrow B_\xi\text{-mod}_{\text{fd}}, \quad (4.34)$$

so that the category $B_\xi\text{-grmod}_{\text{fd}}$ is a graded lift of $\overline{\mathcal{O}}_\xi$. The algebra B_ξ is Morita equivalent to the algebra W_ξ from Theorem 4.22. The following theorem shows that these two algebras coincide for maximally atypical blocks. It gives us hope that the algebra B_ξ can be described in these cases as the path algebra of an infinite quiver with explicit relations.

Theorem 4.25. *For any block $\xi = (0, \nu; m) \in \Xi(m|n)$ of maximal atypicality, there is an isomorphism $W_\xi \xrightarrow{\sim} B_\xi$ such that $1_\lambda \mapsto 1_\lambda$ for each $\lambda \vDash m$.*

Proof. Maximal atypicality implies that all of the irreducible W -modules in $\overline{\mathcal{O}}_\xi$ are one-dimensional. By Theorem 4.22, these are the irreducible W_ξ -modules. Hence, W_ξ is a locally unital basic algebra such that $W_\xi\text{-mod}_{\text{fd}}$ is isomorphic to $\overline{\mathcal{O}}_\xi$. Since the idempotent $1_\lambda \in W_\xi$ acts as the identity on $\overline{L}_\xi(\lambda)$ and as zero on all other $\overline{L}_\xi(\kappa)$'s, it is a primitive idempotent, and the left ideal $W_\xi 1_\lambda$ is isomorphic to the projective cover $\overline{P}_\xi(\lambda)$ of $\overline{L}_\xi(\lambda)$. Then comparing with (4.33), we get that

$$B_\xi \cong \bigoplus_{\kappa, \lambda \models m} \text{Hom}_{W_\xi}(W_\xi 1_\kappa, W_\xi 1_\lambda) \cong \bigoplus_{\kappa, \lambda \models m} 1_\kappa W_\xi 1_\lambda = W_\xi.$$

This isomorphism sends $1_\lambda \mapsto 1_\lambda$ for each $\lambda \models m$. \square

Remark 4.26. Let ξ be as in Theorem 4.25, so that $B_\xi \cong W_\xi$. If $m = n = 1$, we described this algebra already in Remark 4.24. If $m = 1 < n$, using also Theorem 4.22, we see that B_ξ is a quotient of $\mathbb{C}[c^{(2)}, \dots, c^{(n-1)}] \otimes T$, where T is as in Lemma 4.20. Setting $x_i := ef1_i$ and $y_i := fe1_i$, we have that $1_i T 1_i \cong \mathbb{C}[x_i, y_i]$. It follows that $1_{\varepsilon_i} B_\xi 1_{\varepsilon_i}$ is a quotient of the polynomial algebra $\mathbb{C}[c^{(2)}, \dots, c^{(n-1)}, x_i, y_i]$.

Our next theorem computes the graded dimensions of the spaces $1_\kappa B_\xi 1_\lambda$. It is a graded analog of Theorem 4.14. To formulate it, for a positively graded vector space $V = \bigoplus_{n \geq 0} V_n$, we let $\dim_q V := \sum_{n \geq 0} (\dim V_n) q^n$ where q is an indeterminate. Let $[n]$ be the quantum integer $(q^n - q^{-n}) / (q - q^{-1})$, let $[n]!$ be the corresponding quantum factorial, and let $\begin{bmatrix} n \\ r \end{bmatrix}$ be the quantum binomial coefficient.

Theorem 4.27. *For any $\lambda, \kappa \models t$, the space $1_\kappa B_\xi 1_\lambda$ is non-zero if and only if $\kappa = \lambda + \sum_i (\lambda_{i+1} - \rho_{i+1}) \alpha_i$ for a composition ρ with $0 \leq \rho_{i+1} \leq \lambda_{i+1} + \min(\lambda_i, \rho_i)$ for all i , in which case*

$$\dim_q 1_\kappa B_\xi 1_\lambda = [m]! [n]! \sum_{\substack{\tau = (\tau_i)_{i \in \mathbb{Z}} \\ \max(\lambda_{i+1}, \rho_{i+1}) \leq \tau_{i+1} \leq \lambda_{i+1} + \min(\lambda_i, \rho_i)}} q^{s(\tau)} \prod_i \frac{\begin{bmatrix} \lambda_{i+1} + \tau_i - \tau_{i+1} \\ \tau_i - \lambda_i \end{bmatrix} \begin{bmatrix} \lambda_{i+1} + \tau_i - \tau_{i+1} \\ \tau_i - \rho_i \end{bmatrix}}{[\lambda_{i+1} + \tau_i - \tau_{i+1}]! [\lambda_{i+1} + \tau_i - \tau_{i+1} + \gamma_i]!},$$

where

$$s(\tau) := \binom{m}{2} + \binom{n}{2} + \sum_i (2\tau_i - \lambda_i - \rho_i)(\lambda_{i+1} + \tau_i - \tau_{i+1} + \gamma_i) - \sum_i \binom{\lambda_{i+1} + \tau_i - \tau_{i+1}}{2} - \sum_i \binom{\lambda_{i+1} + \tau_i - \tau_{i+1} + \gamma_i}{2}.$$

Proof. See Appendix B. \square

Corollary 4.28. *For all compositions $\lambda \models t$ which are generic in the sense that $\lambda_i \neq 0 \Rightarrow \lambda_i + \lambda_{i+1} + \gamma_i + \gamma_{i+1} = 1$, we have that*

$$\dim_q 1_\lambda B_\xi 1_\lambda = q^{\binom{m}{2} + \binom{n}{2} - \sum_i \binom{\gamma_i}{2}} (1 + q^2)^t [m]! [n]! / \prod_i [\gamma_i]!.$$

Proof. We apply Theorem 4.27 with $\kappa = \lambda$, hence, $\rho_i = 0$ for all i . Letting $I := \{i \in \mathbb{Z} \mid \lambda_{i-1} = 1\}$, the summation is over the compositions τ^J for $J \subseteq I$ defined from $\tau_i^J := \lambda_i + 1$ if $i \in J$, and $\tau_i^J := \lambda_i$ otherwise. We have that

$s(\tau^J) = \binom{m}{2} + \binom{n}{2} - \sum_i \binom{\gamma_i}{2} + 2|J|$, hence, $\sum_{J \subseteq I} q^{s(\tau^J)} = q^{\binom{m}{2} + \binom{n}{2} - \sum_i \binom{\gamma_i}{2}} (1+q^2)^t$, while the big product is always equal to $1/\prod_i [\gamma_i]!$. \square

By Lemma 4.5, the basic algebra B_ξ is a Frobenius algebra. It is also graded and indecomposable. It is a general fact about such algebras that the top degrees of the endomorphism algebras of the indecomposable projective objects are all the same (and these endomorphism algebras are one-dimensional in this degree); see e.g. [R3, Proposition 5.18]. The following describes this top degree explicitly.

Corollary 4.29. *Let $d := m^2 + n^2 - \sum_i \gamma_i^2$. For every $\lambda \models t$, we have that $\dim(1_\lambda B_\xi 1_\lambda)_d = 1$ and $(1_\lambda B_\xi 1_\lambda)_{d'} = 0$ for all $d' > d$.*

Proof. In view of the remarks just made, it suffices to compute the top degree of $1_\lambda B_\xi 1_\lambda$ for generic λ , which is easily done using Corollary 4.28. (One can also check this directly for arbitrary λ ; in general, the monomial of top degree in the polynomial $\dim_q 1_\lambda B_\xi 1_\lambda$ as computed by Theorem 4.27 comes from the summand with $\tau_{i+1} = \lambda_{i+1} + \lambda_i$ for each i .) \square

Conjecture 4.30. The algebra B_ξ is a graded symmetric Frobenius algebra.

The last important algebra that we associate to the block \mathcal{O}_ξ is its *center* C_ξ , i.e. the endomorphism algebra of the identity functor $\text{Id} : \mathcal{O}_\xi \rightarrow \mathcal{O}_\xi$. The double centralizer property implies that C_ξ is also the center of the quotient category $\overline{\mathcal{O}}_\xi$. It can be recovered from the Soergel algebra B_ξ . To explain this, we view elements of B_ξ as infinite matrices of the form $x = (x_{\kappa,\lambda})_{\kappa,\lambda \models t}$ for $x_{\kappa,\lambda} \in 1_\kappa B_\xi 1_\lambda$, all but finitely many of which are zero. If we drop this finiteness condition, we obtain a completion \widehat{B}_ξ of this algebra. In fact, we have simply that

$$\widehat{B}_\xi = \text{End}_W \left(\bigoplus_{\lambda \models t} \overline{P}_\xi(\lambda) \right)^{\text{op}}. \quad (4.35)$$

Moreover, finite-dimensional B_ξ -modules are the same as finite-dimensional modules over the completion. This all depends on the fact that B_ξ is *bounded* in the sense that all of the ideals $1_\kappa B_\xi$ and $B_\xi 1_\lambda$ are finite-dimensional. Finally, we may identify

$$C_\xi = Z(\widehat{B}_\xi). \quad (4.36)$$

The grading on B_ξ induces a positive grading $C_\xi = \bigoplus_{d \geq 0} (C_\xi)_d$.

Lemma 4.31. *The top graded component of C_ξ is $\bigoplus_{\lambda \models t} (1_\lambda C_\xi 1_\lambda)_d$ where $d := m^2 + n^2 - \sum_i \gamma_i^2$, with each summand $(1_\lambda C_\xi 1_\lambda)_d$ being one-dimensional. Also, the Jacobson radical of C_ξ is nilpotent of codimension 1.*

Proof. The first assertion follows from Corollary 4.29. The second assertion then follows because \mathcal{O}_ξ , hence, $\overline{\mathcal{O}}_\xi$, is indecomposable. \square

For the following conjecture, we observe that $B_\xi \cap C_\xi$ is an ideal of C_ξ .

Conjecture 4.32. The image of the canonical map $Z(W) \rightarrow C_\xi$ is isomorphic to $C_\xi / B_\xi \cap C_\xi$.

4.8. Morita and derived equivalences between blocks. In the final subsection, we make some remarks about the problem of classifying blocks of \mathcal{O} up to Morita and/or derived equivalence. Actually, we just look at the integral blocks, which is justified thanks to [CMW, Theorem 3.10]. Recall the definitions of λ^+ and λ^T from the beginning of §4.3.

We begin by discussing derived equivalences. Following [CM, Definition 4.2], we say that \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$ are *gradable derived equivalent* if there is a \mathbb{C} -linear equivalence $F : D^b(\mathcal{O}_\xi) \rightarrow D^b(\mathcal{O}_{\xi'})$ of triangulated categories with inverse G , such that both F and G admit graded lifts.

The following theorem gives many examples of gradable derived equivalences. It comes for free from the theory of braid group actions from [CR].

Theorem 4.33. *Suppose that $\xi = (\mu, \nu; t) \in \Xi(m|n)$ and $i \in \mathbb{Z}$. Let $s_i(\xi) := (s_i(\mu), s_i(\nu); t)$, where $s_i(\mu)$ (resp. $s_i(\nu)$) is obtained by interchanging the i th and $(i+1)$ th parts of μ (resp. ν). Then, there is a gradable derived equivalence*

$$\Theta_i : D^b(\mathcal{O}_\xi) \rightarrow D^b(\mathcal{O}_{s_i(\xi)})$$

inducing a map of the form $[M(\mathbf{A})] \mapsto \pm[M(s_i(\mathbf{A}))]$ at the level of Grothendieck groups, where $s_i(\mathbf{A})$ is obtained by replacing all entries i of $\mathbf{A} \in \xi$ by $(i+1)$ and vice versa. If $t = \mu_i \mu_{i+1} = \nu_i \nu_{i+1} = 0$, this functor is induced by a \mathbb{C} -linear equivalence $\Theta_i : \mathcal{O}_\xi \rightarrow \mathcal{O}_{s_i(\xi)}$ such that $\Theta_i M(\mathbf{A}) \cong M(s_i(\mathbf{A}))$ for all $\mathbf{A} \in \xi$.

Proof. Since we have a categorical action of \mathfrak{sl}_∞ on $\mathcal{O}_\mathbb{Z}$ as described in §4.1, and this categorical action admits a graded lift by [BLW, Theorem 5.26], the existence of Θ_i follows from [CR, Theorem 6.4]. The functor Θ_i is defined there by tensoring with the ‘‘Rickard complex’’, which admits a graded lift by [R1, §5.3.2]. When $t = \mu_i \mu_{i+1} = \nu_i \nu_{i+1} = 0$, the Rickard complex collapses to a single term, hence, it is a ‘‘Scopes equivalence’’. \square

Theorem 4.33 motivates the following conjecture.

Conjecture 4.34. Take blocks $\xi = (\mu, \nu; t) \in \Xi(m|n)$ for $0 \leq m \leq n$ and $\xi' = (\mu', \nu'; t') \in \Xi(m'|n')$ for $0 \leq m' \leq n'$, such that \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$ are non-trivial, i.e. they have more than one isomorphism class of irreducible object. Then \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$ are gradably derived equivalent if and only if $t = t'$, $m = m'$, $n = n'$ and $(\mu + \nu)^T = (\mu' + \nu')^T$.

Part of the ‘‘if’’ implication of this conjecture is implied by Theorem 4.33: \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$ are gradably derived equivalent if $t = t'$, $m = m'$, $n = n'$, $\mu^T = \mu'^T$ and $\nu^T = \nu'^T$. Our hope is that there should be some additional gradable derived equivalences allowing these existing ones to be upgraded to include the case that $(\mu + \nu)^T = (\mu' + \nu')^T$.

The graded algebra C_ξ from (4.36) is an invariant of gradable derived equivalence thanks to [CM, Lemma 4.6]. So, to prove the ‘‘only if’’ direction of Conjecture 4.34, one should look for more information about the structure of C_ξ along the lines of Lemma 4.31. At present, we do not even know how to show that gradably derived equivalent blocks have the same atypicality. We expect that the atypicality of a block should be related to the dimension of its derived category in the sense of [R2].

The remainder of the subsection is concerned with Morita equivalences between blocks. We first point out some obvious ones which arise by twisting with automorphisms of $U(\mathfrak{g})$. For $\xi = (\mu, \nu; t)$ and $\xi' = (\mu', \nu'; t)$, the blocks \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$ are equivalent as \mathbb{C} -linear categories if any of the following hold:

- (“Translation”) There exists $s \in \mathbb{Z}$ such that $\mu_i = \mu'_{i+s}$ and $\nu_i = \nu'_{i+s}$ for all i ; use the automorphism $e_{i,j} \mapsto e_{i,j} + (-1)^{|i|} s \delta_{i,j}$.
- (“Duality”) We have that $\mu_i = \mu'_{-i}$ and $\nu_i = \nu'_{-i}$ for all i ; use the automorphism $e_{i,j} \mapsto -(-1)^{|i||j|} e_{w_0(j), w_0(i)}$ where w_0 is the longest element of $S_m \times S_n$.
- We have that $m = n$, $\mu = \nu'$ and $\nu = \mu'$; use the automorphism that switches the top left and bottom right blocks and the top right and bottom left blocks in the standard matrix realization of \mathfrak{g} .

We also get some more interesting Morita equivalences between typical blocks from the last part of Theorem 4.33: if $t = t' = 0$, $\mu^+ = (\mu')^+$ and $\nu^+ = (\nu')^+$ then \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$ are equivalent.

The following theorem shows that there are very few equivalences between atypical blocks. This was pointed out already in low rank by Coulembier and Serganova in [CS, §6.3]; see also [CS, Remark 6.6] which predicts the importance of the invariants used in the proof of Theorem 4.35.

Theorem 4.35. *Let $\xi = (\mu, \nu; t) \in \Xi(m|n)$ for $0 \leq m \leq n$, and $\xi' = (\mu', \nu'; t') \in \Xi(m'|n')$ for $0 \leq m' \leq n'$. Suppose that \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$ are equivalent as \mathbb{C} -linear categories. Then:*

$$(1) \quad t = t'.$$

Suppose in addition that there is more than one isomorphism class of irreducible object in the blocks \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$. Then:

$$(2) \quad m = m' \text{ and } n = n'.$$

Finally, assume that there are infinitely many isomorphism classes of irreducible objects, so that $t, t' > 0$. Then:

$$(3) \quad \mu + \nu \text{ and } \mu' + \nu' \text{ are equal up to translation and duality.}$$

Proof. In view of Theorem 4.8, the assumption that \mathcal{O}_ξ is equivalent to $\mathcal{O}_{\xi'}$ implies that $\overline{\mathcal{O}}_\xi$ is equivalent to $\overline{\mathcal{O}}_{\xi'}$. If either of the blocks $\overline{\mathcal{O}}_\xi$ or $\overline{\mathcal{O}}_{\xi'}$ has a unique irreducible object up to isomorphism, then so does the other, and we must have that $t = t' = 0$. Otherwise, we have that $t, t' > 0$ and these blocks have infinitely many classes of irreducible objects. To prove (1) and (3) in these cases, we will show that t and $\gamma := \mu + \nu$ can be recovered uniquely (up to translation and duality) from the category $\overline{\mathcal{O}}_\xi$ without using any information about its structure that is external to the abstract \mathbb{C} -linear category.

Starting from $\overline{\mathcal{O}}_\xi$, we can choose a complete set of pairwise inequivalent irreducible objects $\{L(x) \mid x \in X\}$ indexed by some set X . Since we assumed $t > 0$, the set X is infinite. Let $P(x)$ be a projective cover of $L(x)$ and $h(x) := \#\{y \in X \mid [P(x) : L(y)] \neq 0\}$. By Lemma 4.16, we know that the minimal possible value for $h(x)$ as x ranges over all of X is equal to $\binom{t+2}{2}$ for some $t \geq 1$. Thus, we have recovered the atypicality t of the block $\overline{\mathcal{O}}_\xi$ from the underlying abstract category.

Now that we know t , we can define $X_{\min} := \{x \in X \mid h(x) = \binom{t+2}{2}\}$. By Lemma 4.16 again, we know that X_{\min} is in bijection with \mathbb{Z} . To fix a choice of such a bijection, we arbitrarily pick some $x_0 \in X$. Now we appeal to Lemma 4.18. It tells us that the set $\{x \in X_0 \setminus \{x_0\} \mid [P(x_0) : L(x)] \neq 0\}$ contains exactly two elements. We arbitrarily call one of these elements x_1 and the other x_{-1} . Then the set $\{x \in X_0 \setminus \{x_0, x_1\} \mid [P(x_1) : L(x)] \neq 0\}$ is a singleton $\{x_2\}$, the set $\{x \in X_0 \setminus \{x_1, x_2\} \mid [P(x_2) : L(x)] \neq 0\}$ is a singleton $\{x_3\}$, and so on. Similarly, $\{x \in X_0 \setminus \{x_0, x_{-1}\} \mid [P(x_{-1}) : L(x)] \neq 0\}$ is a singleton $\{x_{-2}\}$, and so on. In this way, we have enumerated the elements of X_{\min} as $\{x_i \mid i \in \mathbb{Z}\}$. We have done this in a way that ensures that it agrees with the canonical labelling $\{\bar{L}_\xi(t\varepsilon_i) \mid i \in \mathbb{Z}\}$, at least up to some duality and translation which we can simply ignore due to the symmetry of the invariants that we are about to use.

Next, we explain how to recover the composition γ from the dimensions $\dim \text{End}_{\mathcal{O}_\xi}(P(x_i))$. By the formula in Lemma 4.19, these all take the same value $N := \frac{m!n!}{(t!)^2 \prod_j \gamma_j!} \binom{2t}{t}$ for all but finitely many $i \in \mathbb{Z}$. Thus, we have recovered the number N . Rescaling, we get the numbers

$$d(i) := \binom{2t}{t} \dim \text{End}_{\mathcal{O}_\xi}(P(x_i)) / N = \sum_{r=0}^t \binom{t}{r} \frac{t! \gamma_i! \gamma_{i+1}!}{(\gamma_i + t - r)! (\gamma_{i+1} + r)!}. \quad (4.37)$$

for each $i \in \mathbb{Z}$, and will explain how to recover the γ_i 's uniquely from this sequence. We have that $d(i) = \binom{2t}{t}$ with equality if and only if $\gamma_i = 0 = \gamma_{i+1}$. This already determines all but finitely many of the γ_i 's. Observe moreover that the expression on the right hand side of (4.37) is monotonic in γ_i : it gets strictly smaller if we make the non-negative integer γ_i bigger. So we can use this equation to compute each γ_i uniquely, assuming γ_{i+1} has already been determined inductively (starting from the biggest i such that $\gamma_i \neq 0$).

At this point, we have established (1) and (3). Our proof of (2) requires considerably more force as we need to exploit the existence of the Koszul grading on the basic algebra A_ξ discussed in the previous section. If \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$ are Morita equivalent, then the algebras A_ξ and $A_{\xi'}$ are isomorphic as locally unital graded algebras thanks to the unicity of Koszul gradings. Hence, so too are the algebras B_ξ and $B_{\xi'}$. Since we already know that $t = t'$, we can then invoke Corollary 4.28 to deduce that

$$q^{\binom{m}{2} + \binom{n}{2} - \sum_i \binom{\gamma_i}{2}} [m]! [n]! / \prod_i [\gamma_i]! = q^{\binom{m'}{2} + \binom{n'}{2} - \sum_i \binom{\gamma'_i}{2}} [m']! [n']! / \prod_i [\gamma'_i]!,$$

where $\gamma' := \mu' + \nu'$ of course. When $t, t' > 0$, we already know that $(\gamma)^+ = (\gamma')^+$, so get easily from this that $m = m'$ and $n = n'$. If $t = t' = 0$, we need to use also that the blocks are not trivial (and deduce in addition that $\gamma^T = (\gamma')^T$); one also finds this argument in the proof of [CM, Lemma 8.2]. \square

Corollary 4.36. *For $\xi = (0, \nu; m) \in \Xi(m|n)$ and $\xi' = (0, \nu'; m') \in \Xi(m'|n')$ with $m, m' > 0$, the blocks \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$ are equivalent if and only if $m = m'$ and ν equals ν' up to translation and duality.*

Evidence for the following conjecture comes from Theorem 4.27: it shows that the Soergel algebras B_ξ and $B_{\xi'}$ in the statement have the same graded Cartan matrices.

Conjecture 4.37. Assume that $\xi = (\mu, \nu; t)$ and $\xi' = (\mu', \nu'; t)$ for $\mu, \mu' \models m - t$ and $\nu, \nu' \models n - t$ such that $\mu + \nu$ equals $\mu' + \nu'$ up to translation and duality. Then $B_\xi \cong B_{\xi'}$ as locally unital graded algebras, so that the blocks $\overline{\mathcal{O}}_\xi$ and $\overline{\mathcal{O}}_{\xi'}$ are equivalent.

Believing this conjecture, we also thought initially that the blocks \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$ themselves should also be equivalent (under the same hypotheses as in the conjecture). However, this is too optimistic, due to the following counterexample communicated to us by Coulembier: for $\mathfrak{g} = \mathfrak{gl}_{3|4}(\mathbb{C})$, $t = 1$, and ξ, ξ' defined so that $\mu_1 = 2$, $\mu'_2 = 1$, $\mu'_3 = 1$, $\nu_2 = 1$, $\nu_3 = 1$, $\nu_4 = 1$, $\nu'_1 = 2$ and $\nu'_4 = 1$, the blocks \mathcal{O}_ξ and $\mathcal{O}_{\xi'}$ are not equivalent. To establish this, Coulembier shows that they have different finitistic global dimensions by an application of [CS, Theorem 6.4].

APPENDIX A. PROOF OF THEOREM 3.14

Let notation be as in the statement of the theorem. We note that the case $m = 0$ follows from Corollary 3.12 along with the trivial observations that $M(\mathbf{A}) = M'(\mathbf{A})$ and $\overline{K}(\mathbf{A}) = \overline{M}(\mathbf{A})$ when $m = 0$. So we assume henceforth that $m > 0$. Set $M := M(\mathbf{A})$ for short.

Lemma A.1. $H_0(M)$ is spanned by a subset \mathcal{S} of size 2^m .

Proof. We define

$$\begin{aligned} I^- &:= \{(i, j) \mid i > j, i, j = 1, \dots, m + n\}, \\ I_{\geq}^- &:= \{(i, j) \in I^- \mid \text{col}(i) \leq \text{col}(j)\}, \\ I_{<}^- &:= \{(i, j) \in I^- \mid \text{col}(i) > \text{col}(j)\}. \end{aligned}$$

Let \mathfrak{n}^- be the subalgebra of \mathfrak{g} of strictly lower triangular matrices, so that $\{e_{i,j}\}_{(i,j) \in I^-}$ is a basis of \mathfrak{n}^- . Also $\{e_{i,j}\}_{(i,j) \in I_{\geq}^-}$ is a basis of $\mathfrak{n}^- \cap \mathfrak{p}$ and $\{e_{i,j}\}_{(i,j) \in I_{<}^-}$ is a basis of $\mathfrak{n}^- \cap \mathfrak{m}$. We note that $\mathfrak{n}^- \cap \mathfrak{p} \subseteq \mathfrak{gl}_1$, so the elements of $\{e_{i,j}\}_{(i,j) \in I_{\geq}^-}$ actually all supercommute.

Fix a total order on I^- in such a way that $I_{<}^-$ precedes I_{\geq}^- . The set of ordered monomials of the form $\left(\prod_{(i,j) \in I^-} e_{i,j}^{d_{i,j}}\right) m_{\mathbf{A}}$, where $d_{i,j} \in \mathbb{Z}$ if $\text{row}(i) = \text{row}(j)$ and $d_{i,j} \in \{0, 1\}$ if $\text{row}(i) > \text{row}(j)$, forms a basis of M . For $(i, j) \in I_{<}^-$, we have $e_{ij} \in \mathfrak{m}$, so $e_{ij} - \chi(e_{ij}) \in \mathfrak{m}_\chi$, and $\chi(e_{ij}) \in \{0, \pm 1\}$. Hence, the following ordered monomials span $M/\mathfrak{m}_\chi M$:

$$\left\{ \left(\prod_{(i,j) \in I_{\geq}^-} e_{i,j}^{d_{i,j}} \right) m_{\mathbf{A}} + \mathfrak{m}_\chi M \mid d_{i,j} \in \{0, 1\} \right\}.$$

We are going to cut this spanning set down to one of the required size 2^m .

For $K \subseteq I_{\geq}^-$, we use the notation

$$u(K) := \prod_{(i,j) \in K} e_{i,j} \in U(\mathfrak{n}^- \cap \mathfrak{p}), \quad (\text{A.1})$$

and define $\text{wt}(K) := \sum_{(i,j) \in K} (\delta_i - \delta_j) \in \mathfrak{t}_{\mathbb{Z}}^*$ to be the \mathfrak{t} -weight of $u(K)$. In this paragraph, we are going to focus on the monomials

$$u(K)e_{k,l}u(L)m_{\mathbf{A}} + \mathfrak{m}_\chi M \in M/\mathfrak{m}_\chi M,$$

where $K, L \subseteq I_{\geq}^-$ and $e_{k,l} \in \mathfrak{m}$ or $e_{k,l} \in \mathfrak{b}$. The goal is to describe a *straightening process* in order to prove that this monomial can be expressed as a linear combination of monomials of the form

$$u(J)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M, \quad (\text{A.2})$$

for $J \subseteq I_{\geq}^-$ such that one of the conditions (S1)–(S3) holds:

- (S1) $|J| = |K| + |L|$ and $\text{wt}(J) = \text{wt}(K) + \text{wt}(L) + \delta_k - \delta_l$;
- (S2) $|J| = |K| + |L|$ and $\text{wt}(J) = \text{wt}(K) + \text{wt}(L)$, this is only possible if $k \neq m + 1$ and $l = k - 1$, or if $l = k$; or
- (S3) $|J| < |K| + |L|$.

We proceed by induction on $|K| + |L|$. Our objective is clear when $|K| + |L| = 0$, as $e_{k,l}m_{\mathbf{A}} + \mathfrak{m}_{\chi}M = \chi(e_{k,l}) + \mathfrak{m}_{\chi}M$ if $e_{k,l} \in \mathfrak{m}$, and $e_{k,l}m_{\mathbf{A}} + \mathfrak{m}_{\chi}M = \lambda_{\mathbf{A}}(e_{k,l}) + \mathfrak{m}_{\chi}M$ if $e_{k,l} \in \mathfrak{b}$, where $\lambda_{\mathbf{A}}$ is viewed as an element of \mathfrak{b}^* . Now assume that $|K| + |L| > 0$.

For the case $e_{k,l} \in \mathfrak{m}$, we have

$$u(K)e_{k,l}u(L)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M = \pm e_{k,l}u(K)u(L)m_{\mathbf{A}} + [u(K), e_{k,l}]u(L)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M. \quad (\text{A.3})$$

We do not need to know the sign in the equation above, so we won't specify these explicitly; this is also the case in some other equations below. The first term in (A.3) is (up to sign)

$$e_{k,l}u(K)u(L)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M = \chi(e_{k,l})u(K)u(L) + \mathfrak{m}_{\chi}M,$$

which is zero or (up to sign) a monomial as in (A.2) satisfying (S2).

For the case $e_{k,l} \in \mathfrak{b}$, we have

$$u(K)e_{k,l}u(L)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M = \pm u(K)u(L)e_{k,l}m_{\mathbf{A}} + u(K)[e_{k,l}, u(L)]m_{\mathbf{A}} + \mathfrak{m}_{\chi}M. \quad (\text{A.4})$$

The first term in (A.4) is (up to sign)

$$u(K)u(L)e_{k,l}m_{\mathbf{A}} + \mathfrak{m}_{\chi}M = \lambda_{\mathbf{A}}(e_{k,l})u(K)u(L) + \mathfrak{m}_{\chi}M,$$

where $\lambda_{\mathbf{A}}$ is viewed as an element of \mathfrak{b}^* . This is zero or (up to sign) a monomial as in (A.2) satisfying (S2).

Now we consider the case $e_{k,l} \in \mathfrak{m}$ further. Observe that the term $[u(K), e_{k,l}]$ occurring in (A.3) is a sum of terms of the form $\pm u(K_{i,j})[e_{i,j}, e_{k,l}]u(K^{i,j})$, summed over $(i, j) \in K$ where $K_{i,j}$ is the set of elements of K before (i, j) in our fixed order of I^- and $K^{i,j}$ is the set of those after (i, j) . Either $[e_{i,j}, e_{k,l}]$ is zero, or an element of one of $\mathfrak{n}^- \cap \mathfrak{p}$, \mathfrak{m} or \mathfrak{b} ; note that $\mathfrak{m} \cap \mathfrak{b} \neq \{0\}$ in general, so $[e_{i,j}, e_{k,l}]$ can be an element of both \mathfrak{m} and \mathfrak{b} but this does not matter. If $[e_{i,j}, e_{k,l}] \in \mathfrak{n}^- \cap \mathfrak{p}$, then $u(K_{i,j})[e_{i,j}, e_{k,l}]u(K^{i,j})u(L)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M$ is (up to sign) a monomial of the form (A.2) for which (S1) holds. Whereas if $[e_{i,j}, e_{k,l}] \in \mathfrak{m}$ or $[e_{i,j}, e_{k,l}] \in \mathfrak{b}$, then we can apply induction to deduce that $u(K_{i,j})[e_{i,j}, e_{k,l}]u(K^{i,j})u(L)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M$ is a sum of monomials as in (A.2) that satisfy (S3).

For the case $e_{k,l} \in \mathfrak{b}$ we can argue entirely similarly, but working with the term $u(K)[e_{k,l}, u(L)]$, which occurs in (A.4).

We have now established that our straightening process works.

Now we let $H := \{(i, j) \in I_{\geq}^- \mid i > m + 1\}$, then define

$$\mathcal{H} := \{L \subseteq I_{\geq}^- \mid H \subseteq L\}, \quad \mathcal{S} := \{u(L) + \mathfrak{m}_\chi M \mid L \in \mathcal{H}\}. \quad (\text{A.5})$$

Let $<_{\text{lex}}$ be the order on $\mathfrak{t}_{\mathbb{Z}}^*$ defined by $\sum_{i=1}^{m+n} r_i \delta_i <_{\text{lex}} \sum_{i=1}^{m+n} s_i \delta_i$ if $r_j < s_j$ where j is maximal such that $r_j \neq s_j$. Take $K \subseteq I_{\geq}^-$. Proceeding by induction on $|K|$ and reverse induction on $\text{wt}(K)$ with respect to $<_{\text{lex}}$, we are going to prove that $u(K)$ is in the space spanned by \mathcal{S} . Before proceeding, we note that the case $m = 1$ and π is left justified is trivial, because $H = \emptyset$. So we assume that this is not the case.

For the base step we consider the case $|K| = 0$, so that $u(K) = m_{\mathbb{A}}$. In this case we let $k = 2m + s_-$. Our assumption above implies that k is maximal such that the column of k in the pyramid π has two boxes, and that this is not the leftmost column in π . In particular, $k - 1$ also lies in the second row of π .

Now consider $e_{m, k-1} e_{k, m} m_{\mathbb{A}} + \mathfrak{m}_\chi M$. We have that $e_{m, k-1} \in \mathfrak{m}$ and that $\chi(e_{m, k-1}) = 0$, so that $e_{m, k-1} e_{k, m} m_{\mathbb{A}} + \mathfrak{m}_\chi M = 0 + \mathfrak{m}_\chi M$. Moreover,

$$e_{m, k-1} e_{k, m} m_{\mathbb{A}} + \mathfrak{m}_\chi M = -e_{k, m} e_{m, k-1} m_{\mathbb{A}} + e_{k, k-1} m_{\mathbb{A}} + \mathfrak{m}_\chi M = -m_{\mathbb{A}} + \mathfrak{m}_\chi M,$$

where we use that $e_{m, k-1} m_{\mathbb{A}} = 0$, because $e_{m, k-1} \in \mathfrak{n}$, and that $e_{k, k-1} m_{\mathbb{A}} + \mathfrak{m}_\chi M = -m_{\mathbb{A}} + \mathfrak{m}_\chi M$ because $e_{k, k-1} + 1 \in \mathfrak{m}_\chi$. It follows that $m_{\mathbb{A}} \in \mathfrak{m}_\chi M$, so $m_{\mathbb{A}} + \mathfrak{m}_\chi M$ is certainly in the span of \mathcal{S} .

Next let $K \subseteq I_{\geq}^-$ and assume inductively that $u(L) m_{\mathbb{A}} + \mathfrak{m}_\chi M$ is in the span of \mathcal{S} whenever $|L| < |K|$ or when $|L| = |K|$ and $\text{wt}(L) >_{\text{lex}} \text{wt}(K)$. If $H \subseteq K$, then $u(K) \in \mathcal{S}$, so we may assume this is not the case. We choose $(k, l) \in H \setminus K$ such that k is maximal and l minimal given k . Consider $e_{l, k-1} e_{k, l} u(K) m_{\mathbb{A}}$. Since $e_{l, k-1} \in \mathfrak{m}$ and $\chi(e_{l, k-1}) = 0$, we have $e_{l, k-1} e_{k, l} u(K) m_{\mathbb{A}} + \mathfrak{m}_\chi M = 0 + \mathfrak{m}_\chi M$. Moreover,

$$\begin{aligned} e_{l, k-1} e_{k, l} u(K) m_{\mathbb{A}} + \mathfrak{m}_\chi M &= -e_{k, l} e_{l, k-1} u(K) m_{\mathbb{A}} + e_{k, k-1} u(K) m_{\mathbb{A}} + \mathfrak{m}_\chi M \\ &= -e_{k, l} e_{l, k-1} u(K) m_{\mathbb{A}} - u(K) m_{\mathbb{A}} + \mathfrak{m}_\chi M, \end{aligned}$$

where we use that $e_{k, k-1} u(K) m_{\mathbb{A}} + \mathfrak{m}_\chi M = -u(K) m_{\mathbb{A}} + \mathfrak{m}_\chi M$, because $e_{k, k-1} + 1 \in \mathfrak{m}_\chi$. Thus we see that it suffices to show that $e_{k, l} e_{l, k-1} u(K) m_{\mathbb{A}} + \mathfrak{m}_\chi M$ is in the span of \mathcal{S} .

To see this, we note that the \mathfrak{t} -weight of $e_{k, l} e_{l, k-1} u(K)$ is $\text{wt}(K) + \delta_k - \delta_{k-1} >_{\text{lex}} \text{wt}(K)$. Next we calculate

$$\begin{aligned} e_{k, l} e_{l, k-1} u(K) m_{\mathbb{A}} &= (-1)^{|K|} e_{k, l} u(K) e_{l, k-1} m_{\mathbb{A}} + e_{k, l} [e_{l, k-1}, u(K)] m_{\mathbb{A}} \\ &= e_{k, l} [e_{l, k-1}, u(K)] m_{\mathbb{A}}, \end{aligned}$$

because $e_{l, k-1} \in \mathfrak{n}$ so that $e_{l, k-1} m_{\mathbb{A}} = 0$. Then we see that $[e_{l, k-1}, u(K)]$ is a sum of terms of the form $(-1)^{|K_{i,j}|} u(K_{i,j}) [e_{l, k-1}, e_{i,j}] u(K^{i,j})$ over $(i, j) \in K$. The nonzero possibilities for $[e_{l, k-1}, e_{i,j}]$ are $e_{l, l} + e_{k-1, k-1}$, $e_{l, j}$ or $e_{i, k-1}$ all of which lie in either \mathfrak{m} or \mathfrak{b} . Moreover, we note that $e_{k, k-1}$ is not possible, because $e_{k, l} \notin K$, though $e_{l, l-1}$ can occur. Therefore, using the straightening process, we obtain that $e_{k, l} u(K_{i,j}) [e_{l, k-1}, e_{i,j}] u(K^{i,j}) + \mathfrak{m}_\chi M$ is a sum of monomials of the form $u(J) m_{\mathbb{A}} + \mathfrak{m}_\chi M$ for $J \subseteq I_{\geq}^-$. Moreover, in the present situation the conditions (S1)–(S3) translate to saying that

$$\begin{aligned} (\text{S1}') \quad |J| &= |K| \text{ and } \text{wt}(J) \text{ is either } \text{wt}(K) + \delta_k - \delta_{k-1} \text{ or } \text{wt}(K) + \delta_k - \delta_{k-1} + \\ &\quad \delta_l - \delta_{l-1}; \text{ or} \end{aligned}$$

$$(S2') \quad |J| < |K|.$$

In the first case the possibilities for $\text{wt}(M)$ satisfy $\text{wt}(K) <_{\text{lex}} \text{wt}(M)$. Hence, we can apply induction to deduce that $u(K)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M$ is in the span of \mathcal{S} . Since $|\mathcal{S}| = 2^m$, this completes the proof of the lemma. \square

Lemma A.2. *$H_0(M)$ is a highest weight supermodule for W .*

Proof. We recall that the grading on \mathfrak{g} determined by the pyramid π given by $\deg(e_{i,j}) = \text{col}(j) - \text{col}(i)$ induces a grading on $U(\mathfrak{p})$. We refer to this as grading as the *Lie grading* as in [BBG]. Also we require the explicit formulas for the elements $f^{(r)} \in W \subseteq U(\mathfrak{p})$ for $r = s_- + 1, \dots, s_- + m$ given in [BBG, §4]. We let $\bar{e}_{i,j} := (-1)^{|\text{row}(i)|} e_{i,j}$ and recall that

$$f^{(r)} = S_{\rho'} \left(\sum_{s=1}^r (-1)^{r-s} \sum_{\substack{i_1, \dots, i_s \\ j_1, \dots, j_s}} (-1)^{\#\{a=1, \dots, s-1 \mid \text{row}(j_a)=1\}} \bar{e}_{i_1, j_1} \cdots \bar{e}_{i_s, j_s} \right),$$

where the sum is over all $1 \leq i_1, \dots, i_s, j_1, \dots, j_s \leq m+n$ such that

- (R1) $\text{row}(i_1) = 2$ and $\text{row}(j_s) = 1$;
- (R2) $\text{col}(i_a) \leq \text{col}(j_a)$ ($a = 1, \dots, s$);
- (R3) $\text{row}(i_{a+1}) = \text{row}(j_a)$ ($a = 1, \dots, s-1$);
- (R4) if $\text{row}(j_a) = 2$, then $\text{col}(i_{a+1}) > \text{col}(j_a)$ ($a = 1, \dots, s-1$);
- (R5) if $\text{row}(j_a) = 1$, then $\text{col}(i_{a+1}) \leq \text{col}(j_a)$ ($a = 1, \dots, s-1$);
- (R6) $\deg(e_{i_1, j_1}) + \cdots + \deg(e_{i_s, j_s}) = r - s$.

To spell out the key properties we need from this formula, write

$$f^{(r)} = \pm \sum_{i=1}^{m-r+1} e_{m+i, i+r-(s_-+1)} + g^{(r)}$$

where $g^{(r)} \in U(\mathfrak{p})$ is a linear combination of terms of the form $e_{i_1, j_1} \cdots e_{i_s, j_s} \in U(\mathfrak{p})$ which satisfy the following conditions.

- (F1) The degree of $e_{i_1, j_1} \cdots e_{i_s, j_s}$ in the Lie grading is strictly less than $r-1$.
- (F2) There exists s' such that $\text{row}(i_t), \text{row}(j_t) = 1$ for all $t > s'$ and $\text{row}(i_{s'}) = 2, \text{row}(j_{s'}) = 1$; moreover, if $i_{s'} = m+1$, then $s' = 1$.

Now let \mathcal{H} and \mathcal{S} be as in (A.5). Let $K \in \mathcal{H}$, so that $u(K)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M \in \mathcal{S}$, and let $r \in \{s_- + 1, \dots, s_- + m\}$. We will prove that $g^{(r)}u(K)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M$ is a linear combination of terms of the form $u(L)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M$, where $L \in \mathcal{H}$ such that $u(L)$ has Lie degree strictly less than $u(K) + r - 1$.

Let $i, j \in \{1, \dots, m+n\}$ such that $\text{row}(i) = \text{row}(j) = 1, \text{col}(i) \leq \text{col}(j)$ and let $L \in \mathcal{H}$. Consider $e_{i,j}u(L)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M$. We have that

$$e_{i,j}u(L)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M = [e_{i,j}, u(L)]m_{\mathbf{A}} + u(L)\lambda_{\mathbf{A}}(e_{i,j})m_{\mathbf{A}} + \mathfrak{m}_{\chi}M.$$

Now $[e_{i,j}, u(L)]$ is a sum of terms of the form $\pm u(L_{k,l})[e_{i,j}, e_{k,l}]u(L^{k,l})$ over $(k,l) \in L$, where $L_{k,l}$ is the set of elements of K before (k,l) in our fixed order of I^- and $L^{k,l}$ is the set of those after (k,l) . We have that $[e_{i,j}, e_{k,l}]$ is either zero or equal to $-e_{k,j}$ if $i = l$, and in this case we have $(k,j) \in I_{\geq 0}^-$. It follows that $u(L_{i,j})[e_{i,j}, e_{k,l}]u(L^{i,j})$ is either zero or equal to $\pm u(L')$ for some $L' \in \mathcal{H}$ with the same Lie degree as $e_{i,j}u(L)$.

Next let $i, j \in \{1, \dots, m+n\}$ such that $\text{row}(i) = 2, \text{row}(j) = 1, \text{col}(i) \leq \text{col}(j)$ and let $L \in \mathcal{H}$. We observe that $e_{i,j}u(L) = 0$ if $\text{col}(i) > 1$. Also if $\text{col}(i) = 1$

(so $i = m + 1$), then we have that $e_{i,j}u(L)$ is either zero or equal to $\pm u(L')$ for some $L' \in \mathcal{H}$ with the same Lie degree as $e_{i,j}u(L)$.

Combining the discussion in the previous two paragraphs with the fact that $g^{(r)}u(K)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M$ is a sum of terms of the form $e_{i_1,j_1} \cdots e_{i_s,j_s}u(K)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M$ subject to conditions (F1) and (F2) we deduce that $g^{(r)}u(K)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M$ is a linear combination of terms of the form $u(L)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M$, where $L \in \mathcal{H}$ such that $u(L)$ has Lie degree strictly less than $u(K) + r - 1$.

Now suppose that $(m + 1, r - s_-) \notin K$. Then we have that $e_{m+1,r-s_-}u(K) = \pm u(K \cup \{(m + 1, r - s_-)\})$, and that $e_{m+i,i+r-(s_-+1)}u(K) = 0$ for all $i > 1$. Therefore,

$$f^{(r)}(u(K)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M) = u(K \cup \{(m + 1, r - s_-)\}) + g^{(r)}u(K) + \mathfrak{m}_{\chi}M.$$

We deduce that

$$\left\{ \prod_{r=s_-+1}^{s_-+m} (f^{(r)})^{a_r} u(H)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M \mid a_r \in \{0, 1\} \right\}$$

spans $H_0(M)$, because $(f^{(r)})^{a_r} u(H)m_{\mathbf{A}} + \mathfrak{m}_{\chi}M$ is equal to

$$u(H \cup \{(m + 1, r - s_-) \mid a_r = 1\})m_{\mathbf{A}} + (\text{terms of lower Lie degree}) + \mathfrak{m}_{\chi}M.$$

Hence, $H_0(M)$ is a highest weight supermodule as required. \square

Proof of Theorem 3.14. By Lemma A.2 and the universal property of Verma supermodules, there is a surjective homomorphism $\theta : \Pi^p \overline{M}(\mathbf{B}) \rightarrow H_0(M)$ for some $\mathbf{B} \in \text{Tab}$ and some parity $p \in \mathbb{Z}/2$. By Lemmas 2.1 and 3.6, we have that $[H_0(M)] = [H_0(M'(\mathbf{A}))]$ in the Grothendieck group $K_0(W\text{-smod}_{\text{fd}})$. By Corollary 3.12, $[H_0(M'(\mathbf{A}))] = [\overline{K}(\mathbf{A})]$. These facts imply that $\dim H_0(M) = \dim \overline{K}(\mathbf{A}) = 2^m$, so that the spanning set from Lemma A.1 is actually a basis. Since $\dim \overline{M}(\mathbf{B}) = 2^m$ too, this shows that θ is in fact an isomorphism, and moreover we have established that $[\Pi^p \overline{M}(\mathbf{B})] = [\overline{K}(\mathbf{A})]$.

It remains to show that $p = \bar{0}$ and $\mathbf{B} = \mathbf{A}$. By their definitions (3.10) and (3.7), the W -supermodules $\overline{K}(\mathbf{A})$ and $\overline{M}(\mathbf{B})$ are both diagonalizable with respect to $d_2^{(1)}$, the vectors $\bar{k}_{\mathbf{A}}$ and $\bar{m}_{\mathbf{B}}$ are eigenvectors of $d_2^{(1)}$ -eigenvalues $b(\mathbf{A})$ and $b(\mathbf{B})$, and have parities $\text{par}(b(\mathbf{A}))$ and $\text{par}(b(\mathbf{B}))$, respectively. Moreover, all other $d_2^{(1)}$ -eigenspaces in these supermodules correspond to strictly smaller eigenvalues. As $e^{(r)}$ raises $d_2^{(1)}$ -eigenvalues by one, $\bar{k}_{\mathbf{A}}$ must be a highest weight vector. Hence, there is a non-zero (but not necessarily surjective) homomorphism $\overline{M}(\mathbf{A}) \rightarrow \overline{K}(\mathbf{A}), \bar{m}_{\mathbf{A}} \mapsto \bar{k}_{\mathbf{A}}$. This discussion implies that

$$\begin{aligned} [\overline{K}(\mathbf{A})] &= [\overline{L}(\mathbf{A})] + ([\overline{L}(\mathbf{C})]'s \text{ and } [\Pi \overline{L}(\mathbf{C})]'s \text{ with } b(\mathbf{C}) < b(\mathbf{A})), \\ [\Pi^p \overline{M}(\mathbf{B})] &= [\Pi^p \overline{L}(\mathbf{B})] + ([\overline{L}(\mathbf{C})]'s \text{ and } [\Pi \overline{L}(\mathbf{C})]'s \text{ with } b(\mathbf{C}) < b(\mathbf{A})). \end{aligned}$$

In the previous paragraph, we established already that $[\Pi^p \overline{M}(\mathbf{B})] = [\overline{K}(\mathbf{A})]$. So we must have that $\mathbf{A} = \mathbf{B}$ and $p = 0$, and the proof is complete. \square

APPENDIX B. PROOF OF THEOREM 4.27

Fix $N \geq 2$ and let $U_q \mathfrak{sl}_N$ be the usual quantized enveloping algebra over the field $\mathbb{Q}(q)$ (q an indeterminate) that is associated to the simple Lie algebra $\mathfrak{sl}_N(\mathbb{C})$. We denote its standard generators by $\{F_i, E_i, K_i^{\pm}\}_{1 \leq i < N}$. Let $P :=$

$\bigoplus_{i=1}^N \mathbb{Z}\varepsilon_i$ be its weight lattice, with simple roots $\{\alpha_i := \varepsilon_i - \varepsilon_{i+1}\}_{1 \leq i < N}$ and symmetric form $(-, -)$ defined from $(\varepsilon_i, \varepsilon_j) := \delta_{i,j}$. We have the natural $U_q \mathfrak{sl}_N$ -module V^+ on basis $\{v_i^+\}_{1 \leq i \leq N}$ and the dual natural module V^- on basis $\{v_i^-\}_{1 \leq i \leq N}$. The actions of the generators on these bases are given by the following formulae:

$$\begin{aligned} F_i v_j^+ &= \delta_{i,j} v_{i+1}^+, & E_i v_j^+ &= \delta_{i+1,j} v_i^+, & K_i v_j^+ &= q^{(\alpha_i, \varepsilon_j)} v_j^+, \\ F_i v_j^- &= \delta_{i+1,j} v_i^-, & E_i v_j^- &= \delta_{i,j} v_{i+1}^-, & K_i v_j^- &= q^{(\alpha_i, -\varepsilon_j)} v_j^-. \end{aligned}$$

We'll work with the comultiplication $\Delta : U_q \mathfrak{sl}_N \rightarrow U_q \mathfrak{sl}_N \otimes U_q \mathfrak{sl}_N$ defined from $\Delta(F_i) = 1 \otimes F_i + F_i \otimes K_i$, $\Delta(E_i) = K_i^{-1} \otimes E_i + E_i \otimes 1$, $\Delta(K_i) = K_i \otimes K_i$. This is not quite the same as the comultiplication in Lusztig's book [Lu, §3.1.3]: our q and K_i are Lusztig's v^{-1} and K_i^{-1} . All definitions from [Lu] cited below should be modified accordingly.

Let Θ be the quasi- R -matrix from [Lu, §4.1.1] (with v replaced by q^{-1}), and $R = R_{V,W} : V \otimes W \xrightarrow{\sim} W \otimes V$ be the R -matrix from [Lu, §32.1.4] for any integrable modules V, W . Thus, for vectors $v \in V, w \in W$ of weights $\lambda, \mu \in P$, we have that $R(v \otimes w) = q^{(\lambda, \mu)} \Theta(w \otimes v)$. The following explicit formulae for the action of the inverse of the R -matrix on V^\pm are derived in [BSW, §5]:

$$\begin{aligned} R^{-1}(v_i^+ \otimes v_j^+) &= \begin{cases} v_j^+ \otimes v_i^+ & \text{if } i > j, \\ q^{-1} v_j^+ \otimes v_i^+ & \text{if } i = j, \\ v_j^+ \otimes v_i^+ - (q - q^{-1}) v_i^+ \otimes v_j^+ & \text{if } i < j; \end{cases} \\ R^{-1}(v_i^- \otimes v_j^-) &= \begin{cases} v_j^- \otimes v_i^- & \text{if } i < j, \\ q^{-1} v_j^- \otimes v_i^- & \text{if } i = j, \\ v_j^- \otimes v_i^- - (q - q^{-1}) v_i^- \otimes v_j^- & \text{if } i > j; \end{cases} \\ R^{-1}(v_i^+ \otimes v_j^-) &= \begin{cases} v_j^- \otimes v_i^+ & \text{if } i \neq j, \\ q v_j^- \otimes v_i^+ + (q - q^{-1}) \sum_{r=1}^{N-i} (-q)^r v_{j+r}^- \otimes v_{i+r}^+ & \text{if } i = j; \end{cases} \\ R^{-1}(v_i^- \otimes v_j^+) &= \begin{cases} v_j^+ \otimes v_i^- & \text{if } i \neq j, \\ q v_j^+ \otimes v_i^- + (q - q^{-1}) \sum_{r=1}^{i-1} (-q)^r v_{j-r}^+ \otimes v_{i-r}^- & \text{if } i = j. \end{cases} \end{aligned}$$

For a sign sequence $\sigma = (\sigma_1, \dots, \sigma_k) \in \{\pm\}^k$, we have the tensor space $V^{\otimes \sigma} := V^{\sigma_1} \otimes \dots \otimes V^{\sigma_k}$, with basis $\{v_{i_1}^{\sigma_1} \otimes \dots \otimes v_{i_k}^{\sigma_k}\}_{1 \leq i_1, \dots, i_k \leq N}$. Let $(-, -)$ be the symmetric bilinear form on $V^{\otimes \sigma}$ defined by declaring that this basis is orthonormal. There is an anti-linear algebra automorphism $\psi : U_q \mathfrak{sl}_N \rightarrow U_q \mathfrak{sl}_N$ defined by

$$\psi(F_i) := F_i, \quad \psi(E_i) := E_i, \quad \psi(K_i) := K_i^{-1}.$$

The modules V^\pm possess anti-linear bar-involutions ψ compatible with this in the sense that $\psi(uv) = \psi(u)\psi(v)$ for all $u \in \dot{I}, v \in V^\pm$; these are defined simply so that $\psi(v_i^\pm) = v_i^\pm$ for all $1 \leq i \leq N$. Applying Lusztig's general construction from [Lu, §27.3.1], we get also a (highly non-trivial) compatible bar involution $\psi : V^{\otimes \sigma} \rightarrow V^{\otimes \sigma}$. Finally, let $\psi^* : V^{\otimes \sigma} \rightarrow V^{\otimes \sigma}$ be the adjoint anti-linear

involution to ψ with respect to the form $(-, -)$, i.e. $\overline{(\psi(v), w)} = (v, \psi^*(w))$ for all $v, w \in V^{\otimes \sigma}$.

Lemma B.1. *Let w_0 be the longest element of the symmetric group S_k , so that $w_0(\sigma) = (\sigma_k, \dots, \sigma_1)$. For $s_i = (i \ i+1) \in S_k$, let R_i be the R -matrix $1^{\otimes(i-1)} \otimes R \otimes 1^{\otimes(k-i-1)}$. Then let $R_{w_0} : V^{\otimes w_0(\sigma)} \xrightarrow{\sim} V^{\otimes \sigma}$ be the isomorphism $R_{i_1} \circ \dots \circ R_{i_{k(k-1)/2}}$ obtained from any reduced expression $w_0 = s_{i_1} \dots s_{i_{k(k-1)/2}}$. Define $R_{w_0}^{-1} : V^{\otimes w_0(\sigma)} \xrightarrow{\sim} V^{\otimes \sigma}$ similarly using the inverse R -matrices throughout. Then, we have that*

$$\psi(v_{i_1}^{\sigma_1} \otimes \dots \otimes v_{i_k}^{\sigma_k}) = q^{-\sum_{1 \leq r < s \leq k} (\sigma_{i_r} \varepsilon_{i_r}, \sigma_{i_s} \varepsilon_{i_s})} R_{w_0}(v_{i_k}^{\sigma_k} \otimes \dots \otimes v_{i_1}^{\sigma_1}), \quad (\text{B.1})$$

$$\psi^*(v_{i_1}^{\sigma_1} \otimes \dots \otimes v_{i_k}^{\sigma_k}) = q^{\sum_{1 \leq r < s \leq k} (\sigma_{i_r} \varepsilon_{i_r}, \sigma_{i_s} \varepsilon_{i_s})} R_{w_0}^{-1}(v_{i_k}^{\sigma_k} \otimes \dots \otimes v_{i_1}^{\sigma_1}), \quad (\text{B.2})$$

for any $1 \leq i_1, \dots, i_k \leq N$.

Proof. The formula for ψ follows immediately from Lusztig's construction in [Lu, §27.3.1], plus the formula expressing the R -matrix in terms of the quasi- R -matrix from [Lu, §32.1.4]. To deduce the formula for the adjoint map ψ^* , one reduces to the case that $k = 2$, which may then be checked directly using the formulae for R and R^{-1} displayed above. \square

Henceforth, we will be interested just in the spaces $T^{m|n} := (V^+)^{\otimes m} \otimes (V^-)^{\otimes n}$ for $m, n \geq 0$. Set

$$T := \bigoplus_{m, n \geq 0} T^{m|n},$$

with bar involutions $\psi, \psi^* : T \rightarrow T$ obtained from the ones on each $T^{m|n}$. Like in (4.6), we denote the monomial basis of $T^{m|n}$ by $\{v_{\mathbf{A}}\}_{\mathbf{A} \in \text{Tab}_{m|n}}$, where $\text{Tab}_{m|n}$ denotes the set of all tableaux $\mathbf{A} = \begin{smallmatrix} a_1 \dots a_m \\ b_1 \dots b_n \end{smallmatrix}$ with entries satisfying $1 \leq a_1, \dots, a_m, b_1, \dots, b_n \leq N$. Also let $\text{Tab}_{m|n}^{\circ}$ be the set of all the anti-dominant tableaux in $\text{Tab}_{m|n}$, i.e. the tableaux $\mathbf{A} = \begin{smallmatrix} a_1 \dots a_m \\ b_1 \dots b_n \end{smallmatrix}$ satisfying $1 \leq a_1 \leq \dots \leq a_m \leq N \geq b_1 \geq \dots \geq b_n \geq 1$.

As in [Lu, §27.3.1], the bar involutions ψ and ψ^* have the properties

$$\psi(v_{\mathbf{A}}) = v_{\mathbf{A}} + (\text{a } \mathbb{Z}[q, q^{-1}]\text{-linear combination of } v_{\mathbf{B}} \text{'s for } \mathbf{B} > \mathbf{A}), \quad (\text{B.3})$$

$$\psi^*(v_{\mathbf{A}}) = v_{\mathbf{A}} + (\text{a } \mathbb{Z}[q, q^{-1}]\text{-linear combination of } v_{\mathbf{B}} \text{'s for } \mathbf{B} < \mathbf{A}). \quad (\text{B.4})$$

So we can apply Lusztig's Lemma as in the proof of [Lu, Theorem 27.3.2] to introduce the *canonical basis* $\{b_{\mathbf{A}}\}_{\mathbf{A} \in \text{Tab}_{m|n}}$ and *dual canonical basis* $\{b_{\mathbf{A}}^*\}_{\mathbf{A} \in \text{Tab}_{m|n}}$ of $T^{m|n}$, which are the unique bases determined by the following properties:

$$\psi(b_{\mathbf{A}}) = b_{\mathbf{A}}, \quad b_{\mathbf{A}} \in v_{\mathbf{A}} + \bigoplus_{\mathbf{B} \in \text{Tab}_{m|n}} q\mathbb{Z}[q]v_{\mathbf{B}}, \quad (\text{B.5})$$

$$\psi^*(b_{\mathbf{A}}^*) = b_{\mathbf{A}}^*, \quad b_{\mathbf{A}}^* \in v_{\mathbf{A}} + \bigoplus_{\mathbf{B} \in \text{Tab}_{m|n}} q\mathbb{Z}[q]v_{\mathbf{B}}. \quad (\text{B.6})$$

Since ψ and ψ^* are adjoint, the canonical and dual canonical bases are dual bases with respect to the form $(-, -)$.

Let S be the $\mathbb{Q}(q)$ -algebra defined by generators $\{x_i, y_i\}_{1 \leq i \leq N}$ subject to the following relations:

$$x_i x_j = q x_j x_i \quad \text{if } i > j, \quad (\text{B.7})$$

$$y_i y_j = q y_j y_i \quad \text{if } i < j, \quad (\text{B.8})$$

$$y_i x_j = x_j y_i \quad \text{if } i \neq j, \quad (\text{B.9})$$

$$y_i x_i = q x_i y_i + (q - q^{-1}) \sum_{r=1}^{i-1} (-q)^r x_{i-r} y_{i-r}. \quad (\text{B.10})$$

The algebra S admits compatible gradings $S = \bigoplus_{\gamma \in P} S_\gamma$ and $S = \bigoplus_{m, n \geq 0} S^{m|n}$, the first of which is defined by declaring that $\deg(x_i) := \varepsilon_i$ and $\deg(y_i) := -\varepsilon_i$ for each i , and the second by declaring that $S^{m|n}$ is the span of the monomials

$$u_{\mathbf{A}} := x_{a_1} \cdots x_{a_m} y_{b_1} \cdots y_{b_n} \quad (\text{B.11})$$

for all $\mathbf{A} = \begin{smallmatrix} a_1 \cdots a_m \\ b_1 \cdots b_n \end{smallmatrix} \in \text{Tab}_{m|n}$. The following theorem shows that each $S^{m|n}$ has two distinguished bases: the *monomial basis* $\{u_{\mathbf{A}}\}_{\mathbf{A} \in \text{Tab}_{m|n}^\circ}$ and the *dual canonical basis* $\{d_{\mathbf{A}}\}_{\mathbf{A} \in \text{Tab}_{m|n}^\circ}$, which is explicitly computed. The proof is analogous to that of [B3, Theorem 20].

Theorem B.2. *The vectors $\{u_{\mathbf{A}}\}_{\mathbf{A} \in \text{Tab}_{m|n}^\circ}$ give a basis for $S^{m|n}$. Moreover:*

- (1) *The $\mathbb{Q}(q)$ -linear map $\pi : T \rightarrow S$, $v_{\mathbf{A}} \mapsto u_{\mathbf{A}}$ intertwines the dual bar involution ψ^* on T with the unique anti-linear involution $\psi^* : S \rightarrow S$ such that $\psi^*(x_i) = x_i$, $\psi^*(y_i) = y_i$ and*

$$\psi^*(uu') = q^{(\gamma, \gamma') - mm' - nn'} \psi^*(u') \psi^*(u) \quad (\text{B.12})$$

for all $u \in S^{m|n} \cap S_\gamma$ and $u' \in S^{m'|n'} \cap S_{\gamma'}$.

- (2) *For $\mathbf{A} \in \text{Tab}_{m|n}$, we have that $\pi(b_{\mathbf{A}}^*) = 0$ unless \mathbf{A} is anti-dominant, in which case the vector $d_{\mathbf{A}} := \pi(b_{\mathbf{A}}^*)$ is characterized uniquely by the following properties: $\psi^*(d_{\mathbf{A}}) = d_{\mathbf{A}}$, $d_{\mathbf{A}} \in u_{\mathbf{A}} + \sum_{\mathbf{B} \in \text{Tab}_{m|n}^\circ} q\mathbb{Z}[q]u_{\mathbf{B}}$.*
- (3) *Let $z_0 := 0$ and $z_i := x_i y_i - q x_{i-1} y_{i-1} + \cdots + (-q)^{i-1} x_1 y_1$ for $1 \leq i \leq N$. Given $\mathbf{A} \in \text{Tab}_{m|n}^\circ$ of atypicality t , choose $\begin{smallmatrix} c_1 \cdots c_t a_1 \cdots a_{m-t} \\ c_1 \cdots c_t b_1 \cdots b_{n-t} \end{smallmatrix} \sim \mathbf{A}$ such that $a_1 \leq \cdots \leq a_{m-t}$ and $b_1 \geq \cdots \geq b_{n-t}$. Then:*

$$d_{\mathbf{A}} = q^{-t(t-1)/2 - \#\{(i,j) \mid a_i > c_j\} - \#\{(i,j) \mid b_i > c_j\}} x_{a_1} \cdots x_{a_{m-t}} z_{c_1} \cdots z_{c_t} y_{b_1} \cdots y_{b_{n-t}}.$$

The vectors $\{d_{\mathbf{A}}\}_{\mathbf{A} \in \text{Tab}_{m|n}^\circ}$ give another basis for $S^{m|n}$.

Proof. In this proof, we will cite some results from [B3]. The conventions followed there are consistent with those of [Lu], so that one needs to replace q by q^{-1} and K_i by K_i^{-1} when translating from [B3] to the present setting. The R -matrix $\mathcal{R}_{V,W}$ in [B3] is the same as our inverse R -matrix $R_{V,W}^{-1} := (R_{W,V})^{-1}$ (with q replaced by q^{-1}); hence, in view also of (B.2), the bar involution defined in [B3, (3.2)] corresponds to our ψ^* .

We begin by recalling the standard definitions of the quantum symmetric algebras $S(V^+) = \bigoplus_{m \geq 0} S^m(V^+)$ and $S(V^-) = \bigoplus_{n \geq 0} S^n(V^-)$. As discussed in detail in [B3, §5], the former is the quotient of the tensor algebra $T(V^+)$ by the two-sided ideal

$$I^+ := \langle v_i^+ \otimes v_j^+ - q v_j^+ \otimes v_i^+ \mid i > j \rangle.$$

It has a basis consisting of the images $v_{i_1}^+ \cdots v_{i_m}^+$ of the tensors $v_{i_1}^+ \otimes \cdots \otimes v_{i_m}^+$ for $m \geq 0$ and $i_1 \leq \cdots \leq i_m$. Similarly, $S(V^-)$ is the quotient of $T(V^-)$ by

$$I^- := \langle v_i^- \otimes v_j^- - qv_j^- \otimes v_i^- \mid i < j \rangle,$$

and it has basis $v_{j_1}^- \cdots v_{j_n}^-$ for $n \geq 0$ and $j_1 \geq \cdots \geq j_n$.

Let $\mu^\pm : S(V^\pm) \otimes S(V^\pm) \rightarrow S(V^\pm)$ be the multiplications on these two algebras. Then define a multiplication μ on the vector space $S(V^+) \otimes S(V^-)$ by the composition $(\mu^+ \otimes \mu^-) \circ (\text{id}_{S(V^+)} \otimes R_{S(V^-), S(V^+)}^{-1} \otimes \text{id}_{S(V^-)})$. Since the R -matrix is a braiding, this makes $S(V^+) \otimes S(V^-)$ into an associative algebra. Using the formula for $R^{-1}(v_i^- \otimes v_j^+)$ displayed above, it is easy to check the relations (B.7)–(B.10) to show that there is an algebra homomorphism

$$f : S \rightarrow S(V^+) \otimes S(V^-), \quad x_i \mapsto v_i^+ \otimes 1, y_j \mapsto 1 \otimes v_j^-.$$

Also the relations easily give that the anti-dominant monomials $\{u_A \mid A \in \text{Tab}_{m|n}^\circ\}$ span $S^{m|n}$. Moreover, their images under f are a basis for $S^m(V^+) \otimes S^n(V^-)$. This shows that f is an isomorphism, thereby establishing the first statement of the theorem about the monomial basis.

To prove (1), we consider the diagram

$$\begin{array}{ccc} & T & \\ \pi \swarrow & & \searrow \pi' := f \circ \pi \\ S & \xrightarrow[\sim]{f} & S(V^+) \otimes S(V^-). \end{array}$$

It suffices to define an anti-linear map $\psi^* : S(V^+) \otimes S(V^-) \rightarrow S(V^+) \otimes S(V^-)$ such that $\psi^* \circ \pi' = \pi \circ \psi^*$, and then show that

$$\psi^*((x \otimes y)(x' \otimes y')) = q^{(\alpha+\beta, \alpha'+\beta')} q^{-mm' - nn'} \psi^*(x' \otimes y') \psi^*(x \otimes y) \quad (\text{B.13})$$

for $x \in S^m(V^+)_\alpha, x' \in S^{m'}(V^+)_{\alpha'}, y \in S^n(V^-)_\beta$ and $y' \in S^{n'}(V^-)_{\beta'}$. The generators of the ideal I^+ belong to the dual canonical basis of $V^+ \otimes V^+$, hence, they are fixed by ψ^* . This implies that I^+ is ψ^* -invariant, hence, $\psi^* : T(V^+) \rightarrow T(V^+)$ factors through the quotient $S(V^+)$ to induce $\psi^* : S(V^+) \rightarrow S(V^+)$. The latter map may be defined directly: it is the unique anti-linear involution that fixes all the monomials $v_{i_1}^+ \cdots v_{i_m}^+$ for $m \geq 0$ and $i_1 \leq \cdots \leq i_m$. Similarly, $\psi^* : T(V^-) \rightarrow T(V^-)$ induces $\psi^* : S(V^-) \rightarrow S(V^-)$, which fixes $v_{j_1}^- \cdots v_{j_n}^-$ for $n \geq 0$ and $j_1 \geq \cdots \geq j_n$. Then we let $\psi^* : S(V^+) \otimes S(V^-) \rightarrow S(V^+) \otimes S(V^-)$ be defined from

$$\psi^*(x \otimes y) = q^{(\alpha, \beta)} R_{S(V^-), S(V^+)}^{-1}(\psi^*(y) \otimes \psi^*(x))$$

for $x \in S(V^+)$ of weight α and $y \in S(V^-)$ of weight β . It is immediate from this definition and (B.2) that $\psi^* \circ \pi' = \pi \circ \psi^*$. It remains to establish (B.13). Let $\tilde{\mu}^+ : S(V^+) \otimes S(V^+) \rightarrow S(V^+)$ be the twisted multiplication $m^+ \circ R_{S(V^+), S(V^+)}^{-1}$. Define $\tilde{\mu}^- : S(V^-) \otimes S(V^-) \rightarrow S(V^-)$ similarly. Then let $\tilde{\mu} := (\tilde{\mu}^+ \otimes \tilde{\mu}^-) \circ (\text{id}_{S(V^+)} \otimes R_{S(V^-), S(V^+)}^{-1} \otimes \text{id}_{S(V^-)})$. This gives a twisted multiplication on $S(V^+) \otimes S(V^-)$. Now let x, y, x' and y' be as in (B.13). We apply [B3, Lemma 2] to deduce immediately that

$$\psi^*((x \otimes y)(x' \otimes y')) = q^{(\alpha+\beta, \alpha'+\beta')} \tilde{\mu}(\psi^*(x' \otimes y') \otimes \psi^*(x \otimes y)).$$

We are thus reduced to checking that

$$\tilde{\mu}(\psi^*(x' \otimes y') \otimes \psi^*(x \otimes y)) = q^{-mm' - nn'} \mu(\psi^*(x' \otimes y') \otimes \psi^*(x \otimes y)),$$

which follows as $\tilde{\mu}^+(x \otimes x') = q^{-mm'} \mu^+(x \otimes x')$ and $\tilde{\mu}^-(y \otimes y') = q^{-nn'} \mu^-(y \otimes y')$, as is pointed out at the beginning of the proof of [B3, Theorem 16].

We turn our attention to (2). If $A \in \text{Tab}_{m|n}$ is *not* anti-dominant, then the defining relations for S imply that $u_A = \pi(v_A) = q^k u_B$ for $k > 0$ and $A < B \in \text{Tab}_{m|n}^\circ$. Combined with (B.4), we deduce for $A \in \text{Tab}_{m|n}^\circ$ that

$$\psi^*(u_A) = u_A + (\text{a } \mathbb{Z}[q, q^{-1}]\text{-linear combination of } u_B\text{'s for } A < B \in \text{Tab}_{m|n}^\circ).$$

Hence, we can apply Lusztig's Lemma once again to deduce that $S^{m|n}$ has another basis $\{d_A \mid A \in \text{Tab}_{m|n}^\circ\}$, with d_A being determined uniquely by the properties that $\psi^*(d_A) = d_A$ and $d_A \in u_A + \sum_{B \in \text{Tab}_{m|n}^\circ} q\mathbb{Z}[q]u_B$. This is the basis appearing in the final statement of the theorem. In view of (1), for $A \in \text{Tab}_{m|n}^\circ$, the vector $\pi(b_A^*)$ satisfies the defining properties of d_A , hence, $\pi(b_A^*) = d_A$. To complete the proof of (2), we need to show that $\pi(b_A^*) = 0$ for $A \in \text{Tab}_{m|n} \setminus \text{Tab}_{m|n}^\circ$. This follows because in that case $\pi(b_A^*)$ lies in $\bigoplus_{B \in \text{Tab}_{m|n}^\circ} q\mathbb{Z}[q]u_B$, which contains no non-zero ψ^* -invariant vectors.

Finally, we must establish (3). For this, we first prove the following commutation formulae involving the z_i 's:

$$x_j z_i = \begin{cases} q z_i x_j & \text{if } j > i, \\ q^{-1} z_i x_j & \text{if } j \leq i; \end{cases} \quad (\text{B.14})$$

$$y_j z_i = \begin{cases} q^{-1} z_i y_j & \text{if } j > i, \\ q z_i y_j & \text{if } j \leq i; \end{cases} \quad (\text{B.15})$$

$$z_j z_i = z_i z_j. \quad (\text{B.16})$$

Actually, we just prove (B.14); then the proof of (B.15) is similar, and together they obviously imply (B.16). It is obvious that $x_j z_i = q z_i x_j$ for $j > i$. Also from the definitions we have that

$$x_i y_i = z_i + q z_{i-1}, \quad (\text{B.17})$$

$$y_i x_i = q z_i + z_{i-1}. \quad (\text{B.18})$$

Hence, $z_i x_i = (x_i y_i - q z_{i-1}) x_i = x_i (y_i x_i - z_{i-1}) = q x_i z_i$. Finally, to show that $z_i x_j = q x_j z_i$ for $i > j$, we proceed by induction on i : $z_i x_j = (x_i y_i - q z_{i-1}) x_j = q x_j (x_i y_i - q z_{i-1}) = q x_j z_i$.

Next we derive the formula for d_A under the assumption that $m = n = t$. We need to show simply that $d_A = q^{-t(t-1)/2} z_{c_1} \cdots z_{c_t}$. Since the z 's commute, we may assume that $c_1 \leq \cdots \leq c_t$. We proceed by induction on t , leaving the base case $t = 1$ to the reader as an exercise. For the induction step, we have by induction that $d_{\bar{A}} = q^{-(t-1)(t-2)/2} z_{c_2} \cdots z_{c_t}$ where $\bar{A} := \begin{smallmatrix} c_2 c_3 \cdots c_t \\ c_t \cdots c_3 c_2 \end{smallmatrix}$, and must show that $d_A = q^{-(t-1)} z_{c_1} d_{\bar{A}}$. Expanding the definition of z_{c_1} then commuting y 's past $d_{\bar{A}}$, we get that

$$q^{-(t-1)} z_{c_1} d_{\bar{A}} = x_{c_1} d_{\bar{A}} y_{c_1} - q x_{c_1-1} d_{\bar{A}} y_{c_1-1} + \cdots + (-q)^{c_1-1} x_1 d_{\bar{A}} y_1.$$

It follows easily that this vector lies in $u_A + \sum_B q\mathbb{Z}[q]u_B$, and it just remains to show that it is ψ^* -invariant. Using (B.12), we have that

$$\psi^*(q^{-(t-1)}z_{c_1}d_{\bar{A}}) = q^{t-1}q^{-2(t-1)}d_{\bar{A}}z_{c_1} = q^{-(t-1)}z_{c_1}d_{\bar{A}}.$$

To complete the proof of (3), assume first that $m = t < n$. Let \bar{A} be obtained from A by removing the entry b_{n-t} from its bottom row. By induction on n , we may assume that

$$d_{\bar{A}} = q^{-t(t-1)/2 - \#\{(i,j) \mid b_i > c_j\} + \#\{j \mid b_{n-t} > c_j\}} z_{c_1} \cdots z_{c_t} y_{b_1} \cdots y_{b_{n-t-1}}.$$

We need to show that $d_A = q^{-\#\{j \mid b_{n-t} > c_j\}} d_{\bar{A}} y_{b_{n-t}}$. It is easy to see that it equals u_A plus a $q\mathbb{Z}[q]$ -linear combination of other u_B 's. Then one checks that it is ψ^* -invariant by a calculation using the commutation formulae and (B.12). Finally, one treats the case $m > t$ in a very similar way: let \bar{A} be A with the entry a_1 removed from its top row; by induction we have a formula for $d_{\bar{A}}$; then one deduces that $d_A = q^{-\#\{j \mid a_1 > c_j\}} x_{a_1} d_{\bar{A}}$. \square

Now we switch to the combinatorial framework of §4.3, modified slightly since we are working with \mathfrak{sl}_N rather than \mathfrak{sl}_∞ . We re-use the notation $\lambda \vDash n$ now to indicate that λ is an N -part composition of n , i.e. a sequence $\lambda = (\lambda_1, \dots, \lambda_N)$ of non-negative integers summing to n . Fix for the remainder of the appendix integers $m, n \geq 0$ and a triple $(\mu, \nu; t)$ such that $0 \leq t \leq \min(m, n)$, $\mu \vDash m - t$, $\nu \vDash n - t$, and $\mu_i \nu_i = 0$ for all $i = 1, \dots, N$. For each $\lambda \vDash t$, let $A(\mu, \nu; \lambda)$ be the unique anti-dominant tableau with $\lambda_i + \mu_i$ entries equal to i on its top row and $\lambda_i + \nu_i$ entries equal to i on its bottom row, for all $i = 1, \dots, N$. We denote b_A, v_A, b_A^*, u_A and d_A for $A := A(\mu, \nu; \lambda)$ simply by $b_\lambda, v_\lambda, b_\lambda^*, u_\lambda$ and d_λ , respectively. Set $\gamma := \mu + \nu$.

Lemma B.3. *For $\lambda, \kappa \vDash t$, the d_κ -coefficient of u_λ when expanded in terms of the dual canonical basis for $S^{m|n}$ is non-zero if and only if $\lambda = \kappa - \sum_{i=1}^{N-1} \theta_i \alpha_i$ for $(\theta_1, \dots, \theta_{N-1})$ with $0 \leq \theta_i \leq \lambda_i$ for all i , in which case the coefficient equals*

$$\prod_{i=1}^{N-1} q^{\theta_i(\lambda_{i+1} + \gamma_{i+1})} \begin{bmatrix} \lambda_{i+1} \\ \theta_i \end{bmatrix}.$$

Proof. We first observe by induction on $r \geq 0$ that

$$x_i^r y_i^r = \sum_{s=0}^r q^{sr - r(r-1)/2} \begin{bmatrix} r \\ s \end{bmatrix} z_i^{r-s} z_{i-1}^s. \quad (\text{B.19})$$

The base case is trivial, while the induction step follows using (B.14), (B.16), (B.17) and the usual identity $\begin{bmatrix} r+1 \\ s \end{bmatrix} = q^s \begin{bmatrix} r \\ s \end{bmatrix} + q^{s-r-1} \begin{bmatrix} r \\ s-1 \end{bmatrix}$. Combining (B.19) with (B.15), we get also that

$$(x_i^r y_i^r) y_j^s = q^{-sr} y_j^s (x_i^r y_i^r) \quad (\text{B.20})$$

whenever $j < i$.

Now take any $\lambda \vDash t$ and set $x^\lambda := x_1^{\lambda_1} \cdots x_N^{\lambda_N}$, $y^\lambda := y_N^{\lambda_N} \cdots y_1^{\lambda_1}$ and $z^\lambda := z_1^{\lambda_1} \cdots z_N^{\lambda_N}$. By (B.7)–(B.8) then (B.20), we have that

$$u_\lambda = q^{-\sum_{i < j} \lambda_i (\mu_j + \nu_j)} x^\mu x^\lambda y^\lambda y^\nu = q^{-\sum_{i < j} \lambda_i (\lambda_j + \gamma_j)} x^\mu (x_1^{\lambda_1} y_1^{\lambda_1}) \cdots (x_N^{\lambda_N} y_N^{\lambda_N}) y^\nu.$$

Expanding each $x_i^{\lambda_i} y_i^{\lambda_i}$ here using (B.19), we obtain

$$u_\lambda = q^{-\sum_{i<j} \lambda_i(\lambda_j+\gamma_j)} \sum_{\substack{(\theta_0, \theta_1, \dots, \theta_{N-1}) \\ 0 \leq \theta_i \leq \lambda_{i+1}}} \left(\prod_{i=1}^N q^{\theta_{i-1} \lambda_i - \lambda_i(\lambda_i-1)/2} \begin{bmatrix} \lambda_i \\ \theta_{i-1} \end{bmatrix} \right) x^\mu z^{(\lambda, \theta)} y^\nu$$

where $z^{(\lambda, \theta)} := z_0^{\theta_0} z_1^{\lambda_1+\theta_1-\theta_0} \dots z_{N-1}^{\lambda_{N-1}+\theta_{N-1}-\theta_{N-2}} z_N^{\lambda_N-\theta_{N-1}}$. Since $z_0^{\theta_0} = 0$ unless $\theta_0 = 0$, and $\sum_{i<j} \lambda_i \lambda_j + \sum_i \lambda_i(\lambda_i-1)/2 = t(t-1)/2$, this simplifies to

$$u_\lambda = q^{-t(t-1)/2 - \sum_{i<j} \lambda_i \gamma_j} \sum_{\substack{(\theta_0, \theta_1, \dots, \theta_{N-1}) \\ 0 \leq \theta_i \leq \lambda_{i+1} \\ \theta_0 = 0}} \left(\prod_{i=1}^{N-1} q^{\theta_i \lambda_{i+1}} \begin{bmatrix} \lambda_{i+1} \\ \theta_i \end{bmatrix} \right) x^\mu z^{(\lambda, \theta)} y^\nu.$$

Now pick some $(\theta_0, \theta_1, \dots, \theta_{N-1})$ appearing in this summation, and set $\kappa := \lambda + \sum_{i=1}^{N-1} \theta_i \alpha_i \models t$. By the formula from Theorem B.2(3), we get that

$$d_\kappa = q^{-t(t-1)/2 - \sum_{i<j} \kappa_i \gamma_j} x^\mu z^\kappa y^\nu.$$

Moreover,

$$\sum_{1 \leq i < j \leq N} (\kappa_i - \lambda_i) \gamma_j = \sum_{1 \leq i < j \leq N} (\theta_i - \theta_{i-1}) \gamma_j = \sum_{i=1}^{N-1} \theta_i \gamma_{i+1}.$$

The last three identities displayed combine to show that

$$u_\lambda = \sum_{\substack{(\theta_1, \dots, \theta_{N-1}) \\ 0 \leq \theta_i \leq \lambda_{i+1}}} \left(\prod_{i=1}^{N-1} q^{\theta_i(\lambda_{i+1}+\gamma_{i+1})} \begin{bmatrix} \lambda_{i+1} \\ \theta_i \end{bmatrix} \right) d_{\lambda + \sum_{i=1}^{N-1} \theta_i \alpha_i},$$

and the lemma follows. \square

Lemma B.4. *For any $\lambda, \kappa \models t$, the inner product (b_κ, b_λ) is non-zero if and only if $\kappa = \lambda + \sum_{i=1}^{N-1} (\lambda_{i+1} - \rho_{i+1}) \alpha_i$ for $\rho = (\rho_1, \dots, \rho_N)$ with $\rho_1 = \lambda_1$ and $0 \leq \rho_{i+1} \leq \lambda_{i+1} + \min(\lambda_i, \rho_i)$ for all $i = 1, \dots, N-1$. In that case*

$$(b_\kappa, b_\lambda) = [m]! [n]! \sum_{\tau} q^{s(\tau)} \frac{\prod_{i=2}^N \begin{bmatrix} \lambda_{i+1} + \tau_i - \tau_{i+1} \\ \tau_i - \lambda_i \end{bmatrix} \begin{bmatrix} \lambda_{i+1} + \tau_i - \tau_{i+1} \\ \tau_i - \rho_i \end{bmatrix}}{\prod_{i=1}^N [\lambda_{i+1} + \tau_i - \tau_{i+1}]! [\lambda_{i+1} + \tau_i - \tau_{i+1} + \gamma_i]!},$$

where we interpret λ_{N+1} as zero, the summation is over $\tau = (\tau_1, \dots, \tau_{N+1})$ with $\tau_1 = \lambda_1, \tau_{N+1} = 0$ and $\max(\lambda_{i+1}, \rho_{i+1}) \leq \tau_{i+1} \leq \lambda_{i+1} + \min(\lambda_i, \rho_i)$ for $i = 1, \dots, N-1$, and

$$s(\tau) := \binom{m}{2} + \binom{n}{2} + \sum_{i=2}^N (2\tau_i - \lambda_i - \rho_i)(\lambda_{i+1} + \tau_i - \tau_{i+1} + \gamma_i) - \sum_{i=1}^N \binom{\lambda_{i+1} + \tau_i - \tau_{i+1}}{2} - \sum_{i=1}^N \binom{\lambda_{i+1} + \tau_i - \tau_{i+1} + \gamma_i}{2}.$$

Proof. We have that $b_\lambda = \sum_{\mathbf{B} \in \text{Tab}_{m|n}} (b_\lambda, v_{\mathbf{B}}) v_{\mathbf{B}}$. Hence,

$$(b_\kappa, b_\lambda) = \sum_{\mathbf{B} \in \text{Tab}_{m|n}} (b_\kappa, v_{\mathbf{B}}) (b_\lambda, v_{\mathbf{B}}) = \sum_{\beta \models t} \left[\sum_{\mathbf{B} \sim \mathbf{A}(\mu, \nu; \beta)} (b_\kappa, v_{\mathbf{B}}) (b_\lambda, v_{\mathbf{B}}) \right]. \quad (\text{B.21})$$

To compute the number $(b_\kappa, v_{\mathbf{B}})$ appearing on the right hand side of this formula, we have that $v_{\mathbf{B}} = \sum_{\mathbf{A} \in \text{Tab}_{m|n}} (b_{\mathbf{A}}, v_{\mathbf{B}}) b_{\mathbf{A}}^*$. Hence, in view of Theorem B.2(2), we can compute $(b_\kappa, v_{\mathbf{B}})$ by applying π : it is the d_κ -coefficient of $u_{\mathbf{B}} = \pi(v_{\mathbf{B}})$ when expanded in terms of the dual canonical basis of $\mathcal{S}^{m|n}$. For $\mathbf{B} = \begin{smallmatrix} a_1 & \dots & a_m \\ b_1 & \dots & b_n \end{smallmatrix}$, we let $\ell(\mathbf{B}) := \#\{i < j \mid a_i > a_j\} + \#\{i < j \mid b_i < b_j\}$ so that $u_{\mathbf{B}} = q^{\ell(\mathbf{B})} u_\beta$. Then Lemma B.3 shows that $(b_\kappa, v_{\mathbf{B}})$ is non-zero only if $\beta = \kappa - \sum_{i=1}^{N-1} \theta_i \alpha_i$ for $(\theta_1, \dots, \theta_{N-1})$ with $0 \leq \theta_i \leq \kappa_i$ for each i , in which case

$$(b_\kappa, v_{\mathbf{B}}) = q^{\ell(\mathbf{B})} \prod_{i=1}^{N-1} q^{\theta_i(\beta_{i+1} + \gamma_{i+1})} \begin{bmatrix} \beta_{i+1} \\ \theta_i \end{bmatrix}.$$

Similarly, $(b_\lambda, v_{\mathbf{B}})$ is non-zero only if $\beta = \lambda - \sum_{i=1}^{N-1} \phi_i \alpha_i$ for $(\phi_1, \dots, \phi_{N-1})$ with $0 \leq \phi_i \leq \lambda_i$ for each i , in which case

$$(b_\lambda, v_{\mathbf{B}}) = q^{\ell(\mathbf{B})} \prod_{i=1}^{N-1} q^{\phi_i(\beta_{i+1} + \gamma_{i+1})} \begin{bmatrix} \beta_{i+1} \\ \phi_i \end{bmatrix}.$$

Observe also that

$$\begin{aligned} \sum_{\mathbf{B} \sim \mathbf{A}(\mu, \nu; \beta)} q^{2\ell(\mathbf{B})} &= \frac{q^{\binom{m}{2}} [m]! q^{\binom{n}{2}} [n]!}{\prod_{i=1}^N q^{\binom{\beta_i + \mu_i}{2}} [\beta_i + \mu_i]! q^{\binom{\beta_i + \nu_i}{2}} [\beta_i + \nu_i]!} \\ &= \frac{q^{\binom{m}{2}} [m]! q^{\binom{n}{2}} [n]!}{\prod_{i=1}^N q^{\binom{\beta_i}{2}} [\beta_i]! q^{\binom{\beta_i + \gamma_i}{2}} [\beta_i + \gamma_i]!}. \end{aligned}$$

Putting these observations together, we deduce that the β th summand on the right hand side of (B.21) is non-zero only if $\beta = \kappa - \sum_{i=1}^{N-1} \theta_i \alpha_i = \lambda - \sum_{i=1}^{N-1} \phi_i \alpha_i$ for $(\theta_1, \dots, \theta_{N-1})$ and $(\phi_1, \dots, \phi_{N-1})$ satisfying $0 \leq \theta_i \leq \kappa_i, 0 \leq \phi_i \leq \lambda_i$ for all $i = 1, \dots, N-1$, in which case it equals

$$q^{\binom{m}{2}} [m]! q^{\binom{n}{2}} [n]! \frac{\prod_{i=1}^{N-1} q^{(\phi_i + \theta_i)(\beta_{i+1} + \gamma_{i+1})} \begin{bmatrix} \beta_{i+1} \\ \phi_i \end{bmatrix} \begin{bmatrix} \beta_{i+1} \\ \theta_i \end{bmatrix}}{\prod_{i=1}^N q^{\binom{\beta_i}{2}} [\beta_i]! q^{\binom{\beta_i + \gamma_i}{2}} [\beta_i + \gamma_i]!}. \quad (\text{B.22})$$

Now we complete the proof by repeating the last part of the proof of Theorem 4.14: in the formula (B.22), we replace ϕ_i by $\tau_{i+1} - \lambda_{i+1}$ and θ_i by $\tau_{i+1} - \rho_{i+1}$, to deduce that the β th summand of (B.21) gives a non-zero contribution only if there exist (ρ_2, \dots, ρ_N) and (τ_2, \dots, τ_N) such that $\beta = \lambda + \sum_{i=1}^{N-1} (\lambda_{i+1} - \tau_{i+1}) \alpha_i$, $\kappa = \lambda + \sum_{i=1}^{N-1} (\lambda_{i+1} - \rho_{i+1}) \alpha_i$, and $\max(\lambda_{i+1}, \rho_{i+1}) \leq \tau_{i+1} \leq \lambda_{i+1} + \min(\lambda_i, \rho_i)$ for all $i = 1, \dots, N-1$, interpreting ρ_1 as λ_1 . Then we simplify as before. \square

Proof of Theorem 4.27. This follows from Lemma B.4 together with the discussion in [BLW, §5.9], on passing to the limit as $N \rightarrow \infty$. The main point is that the canonical basis $\{b_{\mathbf{A}} \mid \mathbf{A} \in \text{Tab}_{m|n}\}$ corresponds to the indecomposable graded projectives in the graded lift of $\mathcal{O}_{\mathbb{Z}}$ constructed in *loc. cit.* thanks to [BLW, Corollary 5.30] (and [BLW, Theorem 3.10]). The bilinear form $(-, -)$ is the same as the pairing from [BLW, (5.30)]. \square

REFERENCES

- [Ba] E. Backelin, Representation of the category \mathcal{O} in Whittaker categories, *Int. Math. Res. Notices* 4 (1997), 153–172.

- [BGS] A. Beilinson, V. Ginzburg and W. Soergel, Koszul duality patterns in representation theory, *J. Amer. Math. Soc.* **9** (1996), 473–527.
- [BBG] J. Brown, J. Brundan and S. M. Goodwin, Principal W -algebras for $GL(m|n)$, *Alg. Numb. Theory* **7** (2013), 1849–1882.
- [B1] J. Brundan, Kazhdan–Lusztig polynomials and character formulae for the Lie superalgebra $\mathfrak{gl}(m|n)$, *J. Amer. Math. Soc.* **16** (2003), 185–231.
- [B2] ———, Tilting modules for Lie superalgebras, *Comm. Algebra* **32** (2004), 2251–2268.
- [B3] ———, Dual canonical bases and Kazhdan–Lusztig polynomials, *J. Algebra* **306** (2006), 17–46.
- [B4] ———, Moeglin’s theorem and Goldie rank polynomials in Cartan type A, *Compos. Math.* **147** (2011), 1741–1771.
- [B5] ———, Representations of the general linear Lie superalgebra in the BGG category \mathcal{O} , in: ‘Developments and Retrospectives in Lie Theory: Algebraic Methods,’ eds. G. Mason et al., *Developments in Mathematics* **38**, Springer, 2014, pp. 71–98.
- [BK1] J. Brundan and A. Kleshchev, Representations of shifted Yangians and finite W -algebras, *Mem. Amer. Math. Soc.* **196** (2008).
- [BK2] ———, Schur–Weyl duality for higher levels, *Selecta Math.* **14** (2008), 1–57.
- [BLW] J. Brundan, I. Losev and B. Webster, Tensor product categorifications and the super Kazhdan–Lusztig conjecture, *Int. Math. Res. Notices* **20** (2017), 6329–6410.
- [BSW] J. Brundan, A. Savage and B. Webster, On the definition of quantum Heisenberg category; [arXiv:1812.04779](https://arxiv.org/abs/1812.04779).
- [CLW] S.-J. Cheng, N. Lam and W. Wang, Brundan–Kazhdan–Lusztig conjecture for general linear Lie superalgebras, *Duke Math. J.* **164** (2015), 617–695.
- [CMW] S.-J. Cheng, V. Mazorchuk and W. Wang, Equivalence of blocks for the general linear Lie superalgebra, *Lett. Math. Phys.* **103** (2013), 1313–1327.
- [CR] J. Chuang and R. Rouquier, Derived equivalences for symmetric groups and \mathfrak{sl}_2 -categorification, *Ann. Math.* **167** (2008), 245–298.
- [CM] K. Coulembier and V. Mazorchuk, Dualities and derived equivalences for category \mathcal{O} , *Israel J. Math.* **219** (2017), 661–706.
- [CS] K. Coulembier and V. Serganova, Homological invariants in category \mathcal{O} for the general linear superalgebra, *Trans. Amer. Math. Soc.* **369** (2017), 7961–7997.
- [EGNO] P. Etingof, S. Gelaki, D. Nikshych and V. Ostrik, *Tensor Categories*, Amer. Math. Soc., 2015.
- [GG] W. L. Gan and V. Ginzburg, Quantization of Slodowy slices, *Int. Math. Res. Notices* **5** (2002), 243–255.
- [GKLLRT] I. Gelfand, D. Krob, A. Lascoux, B. Leclerc, V. Retakh and J.-Y. Thibon, Non-commutative symmetric functions, *Adv. Math.* **112** (1995), 218–348.
- [Gk] M. Gorelik, The Kac construction of the center of $U(\mathfrak{g})$ for Lie superalgebras, *J. Nonlinear Math. Phys.* **11** (2004), 325–349.
- [Gw] L. Gow, On the Yangian $Y(\mathfrak{gl}_{m|n})$ and its quantum Berezinian, *Czech. J. Physics* **55** (2005), 1415–1420.
- [K] V. Kac, Characters of typical representations of classical Lie superalgebras, *Comm. Algebra* **5** (1977), 889–897.
- [Ko] B. Kostant, On Whittaker modules and representation theory, *Invent. Math.* **48** (1978), 101–184.
- [L1] I. Losev, Finite W -algebras, *Proceedings of the International Congress of Mathematicians*, vol. III, pp. 1281–1307, Hindustan Book Agency, New Delhi, 2010.
- [L2] ———, Dimensions of irreducible modules over W -algebras and Goldie ranks, *Invent. Math.* **200** (2015), 849–923.
- [Lu] G. Lusztig, *Introduction to Quantum Groups*, Birkhäuser, 1993.
- [M] I. Musson, *Lie Superalgebras and Enveloping Algebras*, Graduate Studies in Mathematics **131**, AMS, 2012.
- [N] M. Nazarov, Quantum Berezinian and the classical Capelli identity, *Lett. Math. Phys.* **21** (1991), 123–131.

- [P] A. Premet, Special transverse slices and their enveloping algebras, *Adv. Math.* **170** (2002), 1–55.
- [R1] R. Rouquier, 2-Kac-Moody algebras; [arXiv:0812.5023](https://arxiv.org/abs/0812.5023).
- [R2] ———, Dimensions of triangulated categories, *J. K-Theory* **1** (2008), 193–256.
- [R3] ———, Automorphismes, graduations et catégories triangulées, *J. Inst. Math. Jussieu* **10** (2011), 713–751.
- [Se] A. Sergeev, The invariant polynomials on simple Lie superalgebras, *Representation Theory* **3** (1999), 250–280.
- [Sk] S. Skryabin, A category equivalence, appendix to [P].
- [S] W. Soergel, Kategorie \mathcal{O} , perverse Garben und Moduln über den Koinvarianten zur Weylgruppe, *J. Amer. Math. Soc.* **3** (1990), 421–445.
- [Z] L. Zhao, Finite W -superalgebras for queer Lie superalgebras, *J. Pure Appl. Algebra* **218** (2014), 1184–1194.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR 97403, USA
E-mail address: brundan@uoregon.edu

SCHOOL OF MATHEMATICS, UNIVERSITY OF BIRMINGHAM, BIRMINGHAM, B15 2TT, UK
E-mail address: s.m.goodwin@bham.ac.uk