

NONDESARGUESIAN PLANES, PARTIAL GEOMETRIES, STRONGLY REGULAR GRAPHS AND CODES
ARISING FROM HYPERBOLIC QUADRICS

William M. Kantor

ABSTRACT

Spreads of hyperbolic quadrics are used to construct translation planes, partial geometries, strongly regular graphs and codes, all having a rich geometric structure.

1. INTRODUCTION

A hyperbolic quadric Q^+ in $PG(2n-1, q)$ has $(q^n-1)(q^{n-1}+1)/(q-1)$ points, and contains $n-1$ -spaces each having $(q^n-1)/(q-1)$ points. A *spread* of Q^+ is a family Σ of $q^{n-1}+1$ subspaces of Q^+ of dimension $n-1$ partitioning the points of Q^+ . A spread can only exist if n is even (when n is odd, Q^+ does not even contain 3 pairwise disjoint $n-1$ -spaces). The only known examples occur when $n=2$, when q is even, or when $n=4$ and $q \equiv 0$ or $2 \pmod{3}$.

In this paper, we will summarize some of the ways spreads of hyperbolic quadrics have been used recently.

2. TRANSLATION PLANES AND THE CONSTRUCTION OF SPREADS

If X is any subspace of $PG(2n-1, q)$, then X^\perp will denote its polar with respect to Q^+ . Consider a spread Σ of Q^+ , and assume that q is even. Fix any point $x \notin Q^+$, and form the family $x^\perp \cap \Sigma = \{x^\perp \cap W \mid W \in \Sigma\}$; clearly, this consists of $n-2$ -spaces partitioning the quadric $x^\perp \cap Q^+$. Since q is even, $x \in x^\perp$ and x^\perp/x is a $PG(2n-3, q)$ equipped with a symplectic (or null) polarity. If $\Sigma(x)$ denotes the projection of $x^\perp \cap \Sigma$ into x^\perp/x , then $\Sigma(x)$ is a spread in the more usual sense: $q^{n-1}+1$ subspaces of dimension $n-2$ which partition the points of x^\perp/x . Consequently Σ and x determine a translation plane $A(\Sigma(x))$ arising from a symplectic spread.

Conversely, suppose that n and q are even, and that Σ' is a symplectic spread of $PG(2n-3, q)$. Then Σ' arises from some Q^+ , Σ and x , as follows (Dillon [4], Dye [5]). We can regard $PG(2n-3, q)$ as our former x^\perp/x , related to Q^+ as be-

fore. Each of the $q^{n-1}+1$ members of Σ' is the projection of a unique $n-2$ -space of $x^\perp \cap Q^+$, and the resulting family Σ'' of $n-2$ -spaces partitions $x^\perp \cap Q^+$. Fix one of the two families of $n-1$ -spaces of Q^+ ; each $n-2$ -space of Q^+ is in a unique member of this family. Thus, Σ'' lifts to a family Σ of $q^{n-1}+1$ of these $n-1$ -spaces, any two having at most a point in common. Since no two of these $n-1$ -spaces can have exactly one point in common (as n is even), Σ consists of pairwise disjoint subspaces. Consequently, Σ is a spread of Q^+ .

The preceding construction produced an essentially unique spread of Q^+ from a symplectic spread of x^\perp/x . In particular, $\Sigma(x)$ essentially determines Σ (where "essentially" means that the symplectic geometry on x^\perp/x can arise from several quadrics in $PG(2n-1, q)$, and that we singled out one of the two families of $n-1$ -spaces of Q^+). However, more is true. Let Σ and Σ_1 be spreads of Q^+ , and let x and x_1 be points off Q^+ . Then any isomorphism from $A(\Sigma(x))$ to $A(\Sigma_1(x_1))$ induces a collineation of $PG(2n-1, q)$ sending Q^+ to itself, Σ to Σ_1 and x to x_1 (Kantor [6]). In particular, each collineation of $A(\Sigma(x))$ is the product of a perspectivity with axis at infinity and a collineation of $PG(2n-1, q)$ preserving Q^+ , Σ and x . Thus, the collineation groups of many translation planes can be found simultaneously once the group $G(\Sigma)$ of collineations preserving Q^+ and Σ is known.

Several spreads Σ are described in Kantor [6,7], and some of the resulting translation planes of order q^{n-1} are studied in detail. When $n=2$, or $n=4$ and $q=2$, only desarguesian planes occur. In all other cases, new planes are obtained. Some of the planes are uninteresting, but others have collineation groups behaving in unusual manners (such as flag-transitively, or behaving as described in Johnson's paper in these proceedings). Complete arcs and dual ovals in all of these planes were found by Thas [13].

The simplest example of a hyperbolic spread is obtained as follows. The desarguesian plane $AG(2, q^{n-1})$ arises from the unique spread of $PG(1, q^{n-1})$, which is trivially symplectic. This produces a symplectic spread Σ' in $PG(2n-3, q)$, and hence (if n and q are even) a spread Σ of a hyperbolic quadric in $PG(2n-1, q)$. This spread is called *desarguesian*, for obvious reasons. It was found by Dillon [4] and Dye [4]; $G(\Sigma)$ was determined by Dye [5] and Cohen and Wilbrink [2]. As we will see in §6, an "affine" version of this spread was discovered much earlier by Kerdock [10].

Other hyperbolic spreads arise from the hermitian curve in $PG(2, q^2)$, from triality, and from field changes generalizing that of the preceding paragraph.

For coordinate descri

It seems unlik
On the other hand, if
spreads probably $\rightarrow \infty$ a

3. MORE TRANSLA

Let Q^+ be a hyp

Consider a spread Σ o
 Q^+ . Let T be any mem
empty or a line. Set
spread of $PG(3, q)$, an
translation planes ob
These translation pla

Let τ be a tria
sets: the points of Q^+
 τ sends lines of Q^+ t
both points and 3-spa
else consists of q^3+1

Let Q^+ be a hyp
Thas [14]) to be a se
count shows that each
that, if $x \in Q^+ - \Omega$,
quadric in x^\perp/x .

If $n > 4$, ovoid
 $PG(2n-1, 2)$ (Kantor [9
spondence) to spreads

If $n=4$, Σ is
Moreover, if T is as
by the Klein correspo
with $\Omega(x)$ than $T \cap \Sigma$,
are more complicated
all known Σ in Kantor
recover Ω or Σ from A
related to automorphi

For coordinate descriptions of many of the known examples we refer to Kantor [6,7].

It seems unlikely that hyperbolic spreads can exist when $n > 4$ and q is odd. On the other hand, if q is even and fixed, the number of inequivalent hyperbolic spreads probably $\rightarrow \infty$ as $n \rightarrow \infty$.

3. MORE TRANSLATION PLANES; OVOIDS

Let Q^+ be a hyperbolic quadric in $PG(7,q)$, where q is now even or odd. Consider a spread Σ of Q^+ . Then Σ belongs to one of the families of 3-spaces of Q^+ . Let T be any member of that family not in Σ . If $W \in \Sigma$ then $T \cap W$ is either empty or a line. Set $T \cap \Sigma = \{T \cap W | W \in \Sigma \text{ and } T \cap W \text{ is a line}\}$. Then $T \cap \Sigma$ is a spread of $PG(3,q)$, and hence determines a translation plane of order q^2 . (The translation planes obtained from Σ as in §2 when q is even have order q^3 , not q^2 .) These translation planes are studied in Kantor [8].

Let τ be a triality map. Then τ cyclically permutes the following three sets: the points of Q^+ , and the two families of 3-spaces on Q^+ . At the same time, τ sends lines of Q^+ to lines of Q^+ , while preserving incidence between lines and both points and 3-spaces of Q^+ . It follows that Σ^τ is either another spread, or else consists of q^3+1 pairwise noncollinear points of Q^+ .

Let Q^+ be a hyperbolic quadric in $PG(2n-1,q)$. An *ovoid* of Q^+ is defined (by Thas [14]) to be a set Ω of $q^{n-1}+1$ pairwise noncollinear points of Q^+ . A simple count shows that each $n-1$ -space on Q^+ contains a unique point of Ω . It follows that, if $x \in Q^+ - \Omega$, then $x^\perp \cap \Omega$ projects onto an ovoid $\Omega(x)$ of the obvious quadric in x^\perp/x .

If $n > 4$, ovoids probably do not exist, but this has only been proven in $PG(2n-1,2)$ (Kantor [9, (4.3)]). If $n=3$, ovoids correspond (under the Klein correspondence) to spreads in $PG(3,q)$, and hence to translation planes of order q^2 .

If $n=4$, Σ is a spread, and Σ^τ is not a spread, then Σ^τ is an ovoid Ω . Moreover, if T is as before then T^τ is a point x , and $T \cap \Sigma$ and $\Omega(x)$ are related by the Klein correspondence. When dealing with coordinates, it is easier to work with $\Omega(x)$ than $T \cap \Sigma$, since points of Ω require 8 coordinates while 3-spaces in Σ are more complicated to describe. The translation planes $A(\Omega(x))$ are studied for all known Σ in Kantor [8]. Unlike the situation in §2, it is not clear how to recover Ω or Σ from $A(\Omega(x))$; moreover, some collineations of the plane need not be related to automorphisms of Ω or Σ .

4. STRONGLY REGULAR GRAPH

Let Σ be a spread of a hyperbolic quadric Q^+ in $PG(2n-1, q)$, where $n \geq 4$. Let Ω be the set of all hyperplanes of members of Σ , so that $|\Omega| = |\Sigma|(q^n - 1)/(q - 1)$ is the number of points of Q^+ . If $X, Y \in \Omega$ and $X \neq Y$, write $X \sim Y \iff X^\perp \cap Y \neq \emptyset$. Then (Ω, \sim) is a strongly regular graph having the same parameters as the collinearity graph (Q^+, \perp) of Q^+ (Kantor [9]). If $n=4$, these graphs are isomorphic (an isomorphism being induced by triality). However, if $n > 4$ they are probably never isomorphic. This is known if $q=2$ (see [9]); the proof uses the nonexistence of ovoids of Q^+ when $q=2$. The graphs are also not isomorphic when Σ is the desarguesian spread defined in §2; this was proved in [9] by brute-force calculations.

If $W \in \Sigma$ let W^* be its set of hyperplanes. Then $\Sigma^* = \{W^* | W \in \Sigma\}$ is a partition of Ω into cliques. If $X \in \Omega - W^*$ then X is joined to exactly $(q^{n-1} - 1)/(q - 1)$ members of W^* , namely, $\{Y \in W^* | X^\perp \cap W \in Y\}$. Thus, each clique W^* inherits from (Ω, \sim) the structure of a $PG(n-1, q)$ in a natural manner. Further properties of (Ω, \sim) reminiscent of properties of (Q^+, \perp) are found in [9]. While it is easy to reconstruct Q^+ and Σ given (Ω, \sim) and Σ^* , it is not known whether Σ^* can be determined from (Ω, \sim) alone.

Variations on the construction of (Ω, \sim) are found in [9]. Instead of a spread of a hyperbolic quadric, spreads of other quadrics, of hermitian geometries, and of symplectic geometries could have been used: strongly regular graphs again arise in exactly the same manner, having the same parameters as the underlying geometries. In the symplectic case, new symmetric designs are also obtained having the parameters of $PG(2n-1, q)$.

5. PARTIAL GEOMETRIES

In this section, Q^+ and Σ will be as before, but q will be 2 or 3. Define Ω as in §4.

If $q=2$, let N be the complement of Q^+ . If $q=3$, let N consist of the points of the complement of "length" 1 (where "length" refers to the value at x of the quadratic form defining Q^+).

It is straightforward to check that the incidence structure (N, Ω, \perp) is a partial geometry $pg(\Sigma)$ (DeClerck, Dye, Thas [3,15]). Namely, if $x \in N$ and $X \in \Omega$, then x is perpendicular to $q^{n-1} + 1$ members of Ω (one per member of Σ), X is perpendicular to $q^{n-1} + 1$ members of N (lying in an affine space), and (if $x \notin X^\perp$) there

are exactly $q^{n-2}(q-1)$

Let $x, y \in N$, x and y are perpendicular in $PG(2n-1, q)$ and Q^+ can be different spreads Σ produce $n=4$ then Σ is essential so is $pg(\Sigma)$. If $q=2$ presumably, the same for $n > 4$ it seems likely

Let $X, Y \in \Omega$, X and Y are not adjacent in (Ω, \sim)

6. CODES

Let Q^+ and Σ be as before. Introduce coordinates (x_1, \dots, x_n) in $PG(2n-1, q)$. Fix a basis $E, F \in \Omega$. Fix a basis $\{e_i, f_i\}$ so that the underlying geometry is $pg(\Sigma)$ with respect to the coordinates.

Let $F' \in \Omega - \{E, F\}$

for a uniquely determined I and J are the zero vectors of Q^+ , and is the identity

Thus, Σ and E, F determine a code. This provides an "affine" code and M'' are the corresponding matrices nonsingular. Thus, M is nonsingular (with zero diagonal)

Conversely, any code C of length n (*) Note that this code can produce different Kerdock codes produce the same spread can be found in Kantor [10]

are exactly $q^{n-2}(q-1)$ pairs (y, Y) with $y \in N$, $Y \in \Omega$, $y \in X^\perp \cap Y^\perp$ and $x \in Y^\perp$.

Let $x, y \in N$, $x \neq y$. Then $x, y \in X^\perp$ for some $X \in \Omega$ if and only if $q=2$ and x and y are perpendicular, or $q=3$ and x and y are not perpendicular. Consequently, $PG(2n-1, q)$ and Q^+ can be recovered from $pg(\Sigma)$. Moreover, projectively inequivalent spreads Σ produce nonisomorphic partial geometries (Kantor [9, (7.3)]). If $n=4$ then Σ is essentially unique (cf. Patterson [12]; Kantor [6, §10]), and hence so is $pg(\Sigma)$. If $q=2$ and $n-1$ is composite then Σ is not unique (Kantor [7, (9.12)]); presumably, the same is true for all even $n > 4$. On the other hand, if $q=3$ and $n > 4$ it seems likely that no Σ exists.

Let $X, Y \in \Omega$, $X \neq Y$. Then $X, Y \in x^\perp$ for some $x \in N$ if and only if X and Y are not adjacent in the graph (Ω, \sim) obtained in §4.

6. CODES

Let Q^+ and Σ be as usual. In order to define some codes, we will have to introduce coordinates. The vector space underlying Q^+ can be written $E \oplus F$ with $E, F \in \Omega$. Fix a basis e_1, \dots, e_n of E , and let f_1, \dots, f_n be the dual basis of F , so that the underlying bilinear form has $(e_i, f_j) = \delta_{ij}$. We will write matrices with respect to the ordered basis $e_1, \dots, e_n, f_1, \dots, f_n$.

Let $F' \in \Omega - \{E\}$. Then

$$F' = F \begin{pmatrix} I & O \\ M' & I \end{pmatrix} \quad (*)$$

for a uniquely determined skew symmetric $n \times n$ matrix M' (with zero diagonal), where O and I are the zero and identity $n \times n$ matrices. (The above $2n \times 2n$ matrix preserves Q^+ , and is the identity on E .)

Thus, Σ and E determine a family $K=K(\Sigma, E)$ of skew symmetric $n \times n$ matrices. This provides an "affine" version of Σ . If $F', F'' \in \Sigma - \{E\}$, $F' \neq F''$, and if M' and M'' are the corresponding matrices, then $F' \cap F'' = 0$ implies that $M' - M''$ is nonsingular. Thus, K is a *Kerdock set*: a set of q^{n-1} skew symmetric $n \times n$ matrices (with zero diagonal) such that the difference of any two is nonsingular.

Conversely, any Kerdock set of $n \times n$ matrices defines a hyperbolic spread via (*). Note that this spread is uniquely determined; thus, inequivalent spreads produce different Kerdock sets. Note, however, that different Kerdock sets can produce the same spread. A further discussion of equivalence of Kerdock sets can be found in Kantor [6, §5].

Now let $q=2$, and fix a Kerdock set K of $n \times n$ matrices. We may assume that $0 \in K$. If $M \in K$, there is a quadratic form $Q_M(x)$ associated with M :

$$xMy^t = Q_M(x+y) - Q_M(x) - Q_M(y)$$

for all $x, y \in \mathbb{Z}_2^n$. Consider the following functions $\mathbb{Z}_2^n \rightarrow \mathbb{Z}_2$:

$$Q_M(x) + L(x) + c \quad (**)$$

for $M \in K$, L a linear functional, and $c \in \mathbb{Z}_2$. There are $2^{n-1} \cdot 2^n \cdot 2 = 2^{2n}$ functions.

Let $C = C(K)$ be the set of all sets of zeros of the various functions (**).

Clearly, C contains all hyperplanes of $AG(n, 2)$ as well as \emptyset and \mathbb{Z}_2^n .

If $X, Y \in C$, then their symmetric difference $X \Delta Y$ has size $0, 2^n, 2^{n-1}, 2^{n-1} \pm 2^{\frac{1}{2}n}$, corresponding to whether $X \Delta Y$ is $\{0\}, \mathbb{Z}_2^{2n}$, an affine hyperplane, or a quadric or its complement in \mathbb{Z}_2^{2n} (cf. Cameron and Seidel [1]). If the vectors in \mathbb{Z}_2^{2n} are called v_1, \dots, v_N with $N = 2^{2n}$, then each subset X of \mathbb{Z}_2^{2n} can be written $X = \sum_{i=1}^N \alpha_i v_i$ with $\alpha_i \in \mathbb{Z}_2$, where $\sum_{i=1}^N$ refers to symmetric difference. In this way, C can be regarded as an error-correcting code (MacWilliams and Sloane [11]), having length 2^n , minimum distance $2^{n-1} - 2^{\frac{1}{2}n}$, and size 2^{2n} . It is extremal in a sense discussed on p. 667 of [11].

The codes $C(K)$ were first discovered by Kerdock [10], with K a Kerdock set arising from the desarguesian hyperbolic spread described in §2. Since inequivalent spreads exist whenever $n-1$ is composite, inequivalent codes $C(K)$ exist as well. However, from a coding theoretic point of view, it is not clear how different codes $C(K)$ differ (for a fixed n): those corresponding to desarguesian spreads seem as if they should be "best", but it is not clear what this means.

BIBLIOGRAPHY

1. P.J. Cameron and J.J. Seidel, Quadratic forms over $GF(2)$, *Kon. Ned. Ak. Wet., Proc.* A76 (1973) 1-8.
2. A.M. Cohen and H.A. Wilbrink, The stabilizer of Dye's spread on a hyperbolic quadric in $PG(4n-1, 2)$ within the orthogonal group (to appear).
3. F. DeClerck, R.H. Dye and J.A. Thas, An infinite class of partial geometries associated with the hyperbolic quadric in $PG(4n-1, 2)$ (to appear).
4. J. Dillon, On Pall partitions for quadratic forms (unpublished manuscript).
5. R.H. Dye, Partitions and their stabilizers for line complexes and quadrics. *Annali di Mat.* (4) 114 (1977) 173-194.
6. W.M. Kantor, Spreads, translation planes and Kerdock sets. I (to appear).

7. W.M. Kantor, S
8. W.M. Kantor, C
9. W.M. Kantor, S
10. A.M. Kerdock,
20 (1972)
11. F.J. MacWillia
North Ho
12. N.J. Patterson
(A)20 (1
13. J.A. Thas, Cor
planes,
14. J.A. Thas, Ovc
15. J.A. Thas, Pol
16. J.A. Thas, Som
(to appe

Bell Laborator
Murray Hill
New Jersey 079
Permanent addr
Mathematics De
University of
Eugene, OR 974

7. W.M. Kantor, Spreads, translation planes and Kerdock sets. II (to appear).
8. W.M. Kantor, Ovoids and translation planes (to appear).
9. W.M. Kantor, Strongly regular graphs defined by spreads (to appear).
10. A.M. Kerdock, A class of low-rate non-linear binary codes, *Inform. Control* 20 (1972) 182-187.
11. F.J. MacWilliams and N.J.A. Sloane, *The theory of error-correcting codes*. North Holland, Amsterdam 1977.
12. N.J. Patterson, A four-dimensional Kerdock set over $GF(3)$, *J. Comb. Theory* (A)20 (1976) 365-366.
13. J.A. Thas, Construction of maximal arcs and dual ovals in translation planes, *Europ. J. Comb.* 1 (1980) 189-192.
14. J.A. Thas, Ovoids and spreads of finite classical polar spaces (to appear).
15. J.A. Thas, Polar spaces, generalized hexagons and perfect codes (to appear).
16. J.A. Thas, Some results on quadrics and a new class of partial geometries (to appear).

Bell Laboratories
Murray Hill
New Jersey 07974, U.S.A.

Permanent address

Mathematics Department
University of Oregon
Eugene, OR 97403, U.S.A.