

What is the Dowling–Wilson conjecture?

Tom Braden ^{*} Jacob P. Matherne [†] Nicholas Proudfoot [‡]

March 11, 2024

The Dowling–Wilson conjecture is a fundamental inequality in combinatorial incidence geometry. The simplest special case answers the following question: given n points in the plane which do not all lie on a line, how many unique lines can they determine? Since each pair of points determines a line, the number of lines is clearly at most $\binom{n}{2}$. A lower bound is less obvious: a 1948 theorem of de Bruijn and Erdős [dBE48] shows that there must be at least n lines. In fact, their proof is a clever counting argument that involves no geometry at all: they show that, given subsets A_1, \dots, A_m of the n points with the property that each pair of points is contained in exactly one of the subsets and no subset contains all of the points, it is necessary that $m \geq n$.

For a more general version of this problem, consider a finite subset E of a vector space over a field \mathbb{F} , and let d be the dimension of the vector space V that they span. The k^{th} **Whitney number** $W_k = W_k(E)$ is defined as the number of distinct k -dimensional linear subspaces of V that can be obtained as the linear span of some subset of E . For example, we have $W_0 = W_d = 1$, corresponding to the zero subspace (spanned by the empty set) and V (spanned by all of E), respectively. In addition, note that $W_1(E) \leq |E|$, with equality unless one of our vectors is a multiple of another. In the case $d = 3$, the nonzero elements of

E determine W_1 points in the projective plane $\mathbb{F}\mathbb{P}^2$. These W_1 points span a total of W_2 lines, and the de Bruijn–Erdős theorem is equivalent to the statement that $W_1 \leq W_2$.

Since then, a number of successively more general results have appeared with the theme “ E determines more subspaces of large dimension than of small dimension.” In 1951, Motzkin [Mot51] proved that $W_1 \leq W_{d-1}$ in arbitrary ambient dimension d . In 1968, Basterfield and Kelly [BK68] gave a combinatorial proof of this fact which does not use projective geometry. In 1975, Dowling and Wilson [DW75] showed that

$$W_0 + W_1 + \dots + W_k \leq W_{d-k} + \dots + W_d \quad (1)$$

whenever $k \leq d/2$, and in a related paper [DW74] they made the following stronger conjecture.

Dowling–Wilson Conjecture. For any $k \leq d/2$, we have $W_k \leq W_{d-k}$.

The Dowling–Wilson conjecture is also called the **top heavy conjecture**, because it says that the poset $\mathcal{L} = \mathcal{L}(E)$ of subspaces of V spanned by elements of E , ranked by dimension, has more elements of high rank than of low rank. When $d = 3$ the Dowling–Wilson conjecture, the inequality (1), and the de Bruijn–Erdős theorem are all equivalent.

Example 1. Suppose that $|E| = n \geq d$, and that the vectors in E are in general position in their span V , meaning that any subset of cardinality at most d is linearly independent. In this case, we have $W_k(E) = \binom{n}{k}$ for all $k < d$, and the inequality $W_k \leq W_{d-k}$ is easy to verify algebraically.

Example 2. Let b_1, \dots, b_n be the standard basis for \mathbb{F}^n , and let $E = \{b_i - b_j \mid i < j \in [n]\}$. These vectors

^{*}The first author is a professor of mathematics at the University of Massachusetts. His email address is braden@umass.edu.

[†]The second author is an assistant professor of mathematics at North Carolina State University. His email address is jpmather@ncsu.edu.

[‡]The third author is a professor of mathematics at the University of Oregon. His email address is njp@uoregon.edu. He received support from NSF Grants DMS-1954050, DMS-2039316, and DMS-2053243.

span the $(n - 1)$ -dimensional subspace

$$V := \left\{ (x_1, \dots, x_n) \in \mathbb{F}^n \mid \sum x_i = 0 \right\}.$$

Subspaces of V spanned by subsets of E are in bijection with partitions of the set $[n]$, where a subspace of dimension k corresponds to a partition with $n - k$ parts. More precisely, if $S = \{S_1, \dots, S_r\}$ is an unordered collection of disjoint nonempty subsets of $[n]$ with $[n] = S_1 \sqcup \dots \sqcup S_r$, then we may define

$$V_S := \left\{ (x_1, \dots, x_n) \in \mathbb{F}^n \mid \sum_{i \in S_j} x_i = 0 \text{ for all } j \right\},$$

which has dimension $n - r$ and is spanned by the vectors $\{b_i - b_j \mid i, j \in S_m \text{ for some } m\}$. Thus $W_k(E)$ is equal to the Stirling number $S(n, n - k)$, which counts partitions of $[n]$ into $n - k$ parts. Already here, the Dowling–Wilson conjecture is not obvious; the resulting inequality on Stirling numbers was proved by Dobson and Rennie [RD69, Theorem 2].

Matroids

In fact, Dowling and Wilson conjectured their inequality in a more general setting than the vector configurations we considered above. Just as the theorem of de Bruijn and Erdős can be stated and proved without reference to linear geometry, the Dowling–Wilson conjecture makes sense for arbitrary matroids, which give a combinatorial abstraction of linear independence and incidence geometry. We point to [Oxl11] for a comprehensive treatment of the theory of matroids. Matroids famously have dozens of equivalent definitions, one of which we give below.

Definition. A **matroid** is a pair (E, rk) , where E is a finite set and rk is a function from the power set of E to the natural numbers satisfying the following conditions:

- $\text{rk}(\emptyset) = 0$.
- For all $S \subset E$ and $e \in E$,

$$\text{rk}(S) \leq \text{rk}(S \cup \{e\}) \leq \text{rk}(S) + 1.$$

- If $\text{rk}(S \cup \{e\}) = \text{rk}(S \cup \{f\}) = \text{rk}(S)$, then

$$\text{rk}(S \cup \{e, f\}) = \text{rk}(S).$$

A subset $F \subset E$ is called a **flat** if it is maximal in its rank, meaning that $\text{rk}(F \cup \{e\}) > \text{rk}(F)$ for all $e \in E \setminus F$. We define the k^{th} **Whitney number** $W_k(E, \text{rk})$ to be the number of flats of rank k .

If E is a finite subset of a vector space V , then we may define $\text{rk}(S)$ to be the dimension of the linear span of S . In this case, a flat is a subset of E with the property that no other element of E is contained in its span. Thus flats of (E, rk) correspond bijectively to subspaces of V spanned by subsets of E , and we have an equality $W_k(E, \text{rk}) = W_k(E)$ relating the two different notions of Whitney numbers.

A matroid that arises from a set (or multiset) of vectors is called **realizable**. Although it is somewhat difficult to come up with non-realizable examples (the smallest is called the Vámos matroid, which has $|E| = 8$), a theorem of Nelson [Nel18] says that *almost all* matroids are non-realizable. More precisely, the percentage of matroids with $E = [n]$ that are realizable goes to zero as n goes to infinity.

The Dowling–Wilson conjecture is now a theorem. Huh and Wang [HW17] used techniques from algebraic geometry to show that it holds for realizable matroids, and more recently Huh, Wang, and the authors [BHM⁺] showed that it holds for arbitrary matroids.

The Möbius algebra and the graded Möbius algebra

Dowling and Wilson’s proof of the inequality (1) for arbitrary matroids can be expressed in an instructive way. For a matroid $M = (E, \text{rk})$, the poset of flats \mathcal{L} is a ranked lattice, and in particular it has a join operation sending flats F, G to $F \vee G$, the unique smallest flat containing both F and G . It has the property that

$$\text{rk}(F \vee G) \leq \text{rk}(F) + \text{rk}(G). \quad (2)$$

When the matroid is realized by a vector configuration, this operation corresponds to taking the sum of vector subspaces of V .

Consider the **Möbius algebra** of \mathcal{L} , which is the \mathbb{Q} -vector space with one basis element y_F for each flat $F \in \mathcal{L}$, and with the multiplication $y_F * y_G = y_{F \vee G}$. Define a pairing by putting

$$\langle y_F, y_G \rangle = \begin{cases} 1 & \text{if } y_F * y_G = y_E \\ 0 & \text{otherwise,} \end{cases}$$

and extending linearly. Let $d = \text{rk}(E)$. For any k , the inequality (2) implies that the subspace spanned by the elements $\{y_F \mid \text{rk } F \leq k\}$ pairs trivially with the subspace spanned by $\{y_F \mid \text{rk } F < d - k\}$. Dowling and Wilson deduced their theorem from a result which is equivalent to the statement that this pairing is nondegenerate, meaning that every nonzero element pairs nontrivially with something. This implies that the pairing induces an injection from the subspace spanned by $\{y_F \mid \text{rk } F \leq k\}$ to the linear dual of the subspace spanned by $\{y_F \mid \text{rk } F \geq d - k\}$, which implies the inequality (1).

In passing from the inequality (1) to the full conjecture, it is natural to pass from the Möbius algebra to the **graded Möbius algebra** $H^*(M)$, which has the same underlying vector space, but with the modified multiplication

$$y_F \cdot y_G = \begin{cases} y_{F \vee G} & \text{if } \text{rk}(F \vee G) = \text{rk}(F) + \text{rk}(G) \\ 0 & \text{otherwise.} \end{cases}$$

It is a graded algebra: if $H^k(M)$ is the span of the y_F with $\text{rk } F = k$, then $H^j(M) \cdot H^k(M) \subset H^{j+k}(M)$. (In technical language, it is the associated graded algebra obtained from a filtration of the Möbius algebra.) One can define a new pairing on $H^*(M)$ in the same way as before, this time using the modified multiplication. However, this pairing cannot be nondegenerate because $H^k(M)$ and $H^{d-k}(M)$ do not have the same dimension.

To prove the Dowling–Wilson conjecture it would be enough to show that every nonzero element of $H^k(M)$ pairs nontrivially with some element of $H^{d-k}(M)$ whenever $k \leq d/2$, or equivalently that the pairing induces an injection of $H^k(M)$ into the linear dual of $H^{d-k}(M)$ for all $k \leq d/2$. This was proved by Kung when $k = 1$ [Kun79, Corollary 3.3], but the statement is false when $k = 2$. Instead, [HW17] and

[BHM⁺] deduce the Dowling–Wilson conjecture from a different statement: if $L \in H^1(M)$ is a positive combination of y_F over all rank 1 flats F , then for any $k \leq d/2$, the multiplication

$$L^{d-2k}: H^k(M) \rightarrow H^{d-k}(M)$$

is injective.

The proof in the realizable case

In the realizable case, this injectivity was proved by Huh and Wang by interpreting the graded Möbius algebra $H^*(M)$ as the cohomology ring of an algebraic variety, which we now describe. We assume for simplicity that our matroid is realizable by vectors over the complex numbers \mathbb{C} ; the case of arbitrary fields can be obtained by replacing singular cohomology with ℓ -adic étale cohomology.

Let V be a vector space over \mathbb{C} spanned by a finite set of vectors E . Let V^* be the linear dual of V , consisting of linear maps from V to \mathbb{C} . We define a linear map $i: V^* \rightarrow \mathbb{C}^E$ whose e^{th} coordinate is given by evaluation on the element $e \in E \subset V$. The fact that V is spanned by E implies that i is an injection. Consider the Riemann sphere $\mathbb{CP}^1 = \mathbb{C} \cup \{\infty\}$, and let Y be the closure of $i(V^*)$ inside of $(\mathbb{CP}^1)^E$. The algebraic variety Y is called the **arrangement Schubert variety** of the pair (V, E) , in analogy with classical Schubert varieties in Lie theory. The connection between the geometry of Y and the structure of the matroid M represented by E was first explored by Ardila and Boocher [AB16].

The additive action of V^* on itself extends to an action of V^* on Y . This action has finitely many orbits, indexed by the flats of M , and each orbit is isomorphic to an affine space. More precisely, for any subset $S \subset E$, let $p_S \in (\mathbb{CP}^1)^E$ be the point whose e^{th} coordinate is equal to 0 if $e \in S$ and ∞ otherwise. Then:

- The point p_S lies in Y if and only if S is a flat of the matroid M .
- If $F \subset E$ is a flat, then the stabilizer of p_F in V^* is equal to $\text{Span}(F)^\perp$, and therefore its orbit is isomorphic to $V^*/\text{Span}(F)^\perp \cong \text{Span}(F)^*$. In particular, it is an affine space of dimension $\text{rk } F$.

- Every element of Y lies in the orbit of exactly one point p_F .

For a schematic picture of the orbits for the vector arrangement of Example 2 with $n = 3$, see Figure 1.




Figure 1: The Schubert variety Y associated with the vector arrangement of Examples 2 and 3 with $n = 3$. Points are labeled by partitions of the set $[3]$, which correspond to flats. They are also labeled by triples of elements of $\mathbb{C}\mathbb{P}^1$ corresponding to the three vectors $b_1 - b_2, b_1 - b_3, b_2 - b_3 \in V$. For example, the $b_1 - b_2$ and $b_2 - b_3$ coordinates of the point $p_{\{13,2\}}$ are equal to ∞ , while the $b_1 - b_3$ coordinate is equal to 0 because 1 and 3 lie in the same block of the partition.

Example 3. Consider the vector configuration from Example 2. We may regard V^* as the space of ordered n -tuples of points in \mathbb{C} up to simultaneous translation. Under this identification, the element $b_i - b_j \in E$ is identified with the linear functional that takes an n -tuple of complex numbers to the difference between the i^{th} and j^{th} points, and the compactification Y of V^* is obtained by allowing these distances to go to ∞ . Recall from Example 2 that flats are in bijection with partitions of the set $[n]$. If F is the flat corresponding to the partition $[n] = S_1 \sqcup \cdots \sqcup S_r$, then the orbit containing p_F consists of all tuples for

which the distance between the i^{th} and j^{th} points is finite if and only if i and j lie in the same part of the partition.

The decomposition of Y into affine spaces is a topological cell decomposition with all cells of even dimension. In particular, it implies that the cohomology $H^*(Y; \mathbb{Q})$ vanishes in odd degrees, and that the dimension of $H^{2k}(Y; \mathbb{Q})$ is the k^{th} Whitney number $W_k(M)$. In fact, we have a stronger statement: as a ring, $H^*(Y; \mathbb{Q})$ is isomorphic to the graded Möbius algebra $H^*(M)$, with degrees doubled. (This is not needed to prove the Dowling–Wilson conjecture in the realizable case, but it is key to generalizing it to all matroids.)

Because the variety Y is singular, it is natural to consider its **intersection cohomology** $\text{IH}^*(Y; \mathbb{Q})$, which is a graded module over $H^*(Y; \mathbb{Q})$. For smooth algebraic varieties, intersection cohomology is isomorphic to ordinary cohomology, while for singular varieties the intersection cohomology retains many of the important properties of the cohomology of smooth varieties. In particular, since Y is a projective complex algebraic variety, it satisfies the **hard Lefschetz property**: If $L \in H^2(Y; \mathbb{Q})$ is ample, then for any $j \leq d = \dim Y$, the multiplication map

$$L^{d-j} : \text{IH}^j(Y; \mathbb{Q}) \rightarrow \text{IH}^{2d-j}(Y; \mathbb{Q}) \quad (3)$$

is an isomorphism.

Because Y is a proper variety which has a decomposition into affine spaces, an argument of Björner and Ekedahl [BE09] implies that the graded $H^*(Y; \mathbb{Q})$ -module $\text{IH}^*(Y; \mathbb{Q})$ has a graded submodule isomorphic to $H^*(Y; \mathbb{Q})$, regarded as a module over itself. This is the last ingredient that we needed to prove the Dowling–Wilson conjecture! Indeed, let L be an ample class in $H^2(Y; \mathbb{Q})$. Restricting the isomorphism (3) to the submodule $H^*(Y; \mathbb{Q})$ gives an injection

$$H^{2k}(Y; \mathbb{Q}) \rightarrow H^{2d-2k}(Y; \mathbb{Q}).$$

Since the source and target have dimension $W_k(M)$ and $W_{d-k}(M)$, respectively, we obtain the inequality $W_k(M) \leq W_{d-k}(M)$.

The proof for general matroids

The proof of the full Dowling–Wilson conjecture in [BHM⁺] follows the same basic plan as the proof in the realizable case. Although there is no analogue of the variety Y for general matroids, we still have an analogue of its cohomology ring $H^*(Y; \mathbb{Q})$, namely the graded Möbius algebra $H^*(M)$. What is needed is to define a graded module $IH^*(M)$ over the graded ring $H^*(M)$, called the **intersection cohomology module** of the matroid M , and to show that it satisfies the conditions needed for the argument in the previous section:

- $IH^*(M)$ has a submodule isomorphic to $H^*(M)$.
- $IH^*(M)$ satisfies the hard Lefschetz property.

The first condition is immediate from the definition of $IH^*(M)$ (which we will not give here). The second condition is proved via an elaborate induction in which the hard Lefschetz property is one of fifteen different properties of the module $IH^*(M)$ that are simultaneously proved [BHM⁺, Theorem 3.16]. For a diagram depicting the structure of this induction, see [BHM⁺, Figure 1].

Many of the steps of the induction are motivated by statements that are known geometrically in the realizable case. There are also strong similarities with two other notable examples where “intersection cohomology” has been defined and Hodge-theoretic statements such as hard Lefschetz have been proved for a non-existent variety: the intersection cohomology of non-rational fans of Karu [Kar04], and Elias and Williamson’s Hodge theory of Soergel bimodules [EW14].

References

- [AB16] Federico Ardila and Adam Boocher, *The closure of a linear space in a product of lines*, J. Algebraic Combin. **43** (2016), no. 1, 199–235. MR3439307
- [BE09] Anders Björner and Torsten Ekedahl, *On the shape of Bruhat intervals*, Ann. of Math. (2) **170** (2009), no. 2, 799–817. MR2552108
- [BHM⁺] Tom Braden, June Huh, Jacob Matherne, Nicholas Proudfoot, and Botong Wang, *Singular Hodge theory for combinatorial geometries*. arXiv:2010.06088.
- [BK68] J. G. Basterfield and L. M. Kelly, *A characterization of sets of n points which determine n hyperplanes*, Proc. Cambridge Philos. Soc. **64** (1968), 585–588. MR233719
- [dBE48] N. G. de Bruijn and P. Erdős, *On a combinatorial problem*, Indagationes Math. **10** (1948), 421–423. MR28289
- [DW74] Thomas A. Dowling and Richard M. Wilson, *The slimmest geometric lattices*, Trans. Amer. Math. Soc. **196** (1974), 203–215. MR345849
- [DW75] ———, *Whitney number inequalities for geometric lattices*, Proc. Amer. Math. Soc. **47** (1975), 504–512. MR354422
- [EW14] Ben Elias and Geordie Williamson, *The Hodge theory of Soergel bimodules*, Ann. of Math. (2) **180** (2014), no. 3, 1089–1136. MR3245013
- [HW17] June Huh and Botong Wang, *Enumeration of points, lines, planes, etc.*, Acta Math. **218** (2017), no. 2, 297–317. MR3733101
- [Kar04] Kalle Karu, *Hard Lefschetz theorem for nonrational polytopes*, Invent. Math. **157** (2004), no. 2, 419–447. MR2076929
- [Kun79] Joseph P. S. Kung, *The Radon transforms of a combinatorial geometry. I*, J. Combin. Theory Ser. A **26** (1979), no. 2, 97–102. MR530281
- [Mot51] Theodore Motzkin, *The lines and planes connecting the points of a finite set*, Trans. Amer. Math. Soc. **70** (1951), 451–464. MR41447
- [Nel18] Peter Nelson, *Almost all matroids are nonrepresentable*, Bull. Lond. Math. Soc. **50** (2018), no. 2, 245–248. MR3830117
- [Oxl11] James Oxley, *Matroid theory*, Second, Oxford Graduate Texts in Mathematics, vol. 21, Oxford University Press, Oxford, 2011. MR2849819
- [RD69] B. C. Rennie and A. J. Dobson, *On Stirling numbers of the second kind*, J. Combinatorial Theory **7** (1969), 116–121. MR241310